

Published in final edited form as:

Conf Proc IEEE Eng Med Biol Soc. 2018 July; 2018: 2374–2377. doi:10.1109/EMBC.2018.8512848.

Microbiota of Inflammatory Bowel Disease Models

Ziyang Gao,

Department of Biomedical Engineering, Duke University, Durham, NC 27708 USA (ziyang.gao@duke.edu).

Kai-Yuan Chen,

Department of Biomedical Engineering, Duke University, Durham, NC 27708 USA (phone: 607-793-0198; kai.yuan.chen@duke.edu).

Olaf Mueller.

Duke Center for Genomic and Computational Biology, Duke University, Durham, NC 27708 (olaf.mueller@duke.edu)

Han Zhang,

School of Medicine, Johns Hopkins University, Baltimore, MD 21218 USA (hzhang94@jhmi.edu)

Nikolai Rakhilin,

School of Electrical and Computational Engineering, Cornell University, Ithaca, NY 14850 USA (nr233@cornell.edu)

Jiande Chen,

School of Medicine, Johns Hopkins University, Baltimore, MD 21218 USA (jchen184@jhmi.edu)

Xiling Shen

Department of Biomedical Engineering, Duke University, Durham, NC 27708 USA (xs37@duke.edu).

Abstract

Gut microbiome plays an important role in inflammatory bowel disease (IBD), a group of intestinal chronic inflammation conditions that affect a large population. The animal models of IBD have long been established on basis of pathological features, but their ability to recapitulate patient gut microbiota is unknown. We investigated and compared the composition and biodiversity of bacterial population in the fecal samples from rat models of the two IBD subtypes, and compared them with patient samples. Our analyses revealed that inflammation reduces overall microbiome diversity and increased variation between individuals. We identified specific microbial signatures associated with the two IBD subtypes that were consistent between the animal models and human IBD patients, suggesting that the animal models can partially recapitulate the microbiota in human diseases. Furthermore, metagenome prediction analysis suggested microbial functions that were likely altered by host-microbiota interactions in IBD models.

I. Introduction

IBD is a group of intestinal disorders entailing chronic and relapsing inflammation of the lower digestive tract[1]. There are two major types of human IBDs: Crohn's disease (CD)

and ulcerative colitis (UC), with very different signs and symptoms[2]. Gut microbiota plays an important role in maintaining the integrity of the intestinal mucosa, the alteration of which contributes to the chronic inflammation conditions[3–5].

Dextran Sulfate Sodium (DSS) and 2,4,6-trinitrovebzebe sulfonic acid (TNBS) are widely used to induce UC and CD in animal model respectively[6, 7]. Although the DSS and TNBS treated animals present similar pathological features as human IBD specimens, whether they recapitulate patients' microbiota patterns have not been examined.

We performed 16S ribosome RNA sequencing (16S rRNA-seq) to examine the microbiomes of DSS and TNBS models and compared them with those of UC and CD patients. The biodiversity and microbial signature profiling analyses revealed that microbiota patterns in disease models were significantly altered from health control, and partially overlapped with human IBD.

II. Material and methods

A. IBD rats model

Male rats weighed around 250 grams were used. They were fed with drinking water containing 4% DSS to induce UC (n=10). 22.5mg TNBS in 40% ethanol (total volume 0.75ml) was given via enema once at baseline to induce CD (n=13). Fecal samples were collected at baseline (Day 0) and the end points of chemical induction (TNBS: Day5, DSS: Day7).

B. Sequencing and statistical analysis

The 16s rRNA V3 region of each sample was sequenced via Illumina MiSeq[8, 9]. We processed the raw sequencing data by Qiime 1.9.1, with open reference OTUs picked against Greengenes 13_5[10, 11]. The R package ggplot2 was utilized to visualized the results of taxonomy, alpha and beta diversity analyzed by Qiime[12]. Differential analysis of taxonomies was performed using LEfSe[13]. Morpheus[14] was used to process heatmaps showing relative abundance of corresponding bacterial.

C. Metagenome prediction

The functional contents were computed through metagenome prediction analysis via Phylogenetic Investigation of Communities by Reconstruction of Unobserved States (PICRUSt)[15]. The predicted functions were assigned against Kyoto Encyclopedia of Genes and Genomes (KEGG) Orthology database[16]. Differential functions analysis was done by applying limma package at KO level 2[17].

III. Results

A. a-Diversity significantly altered in inflammation models

We first performed α -diversity analysis on 16s rRNA data to investigate the microbiota diversity within individual samples (Figure 1). In UC and CD models, the shannon index

decrease remarkably after the treatment of DSS and TNBS, which suggests the microbiota diversity was greatly reduced by inflammation.

B. Gut inflammation induces shift in biodiversity

We calculated the unifrac distance matrix from beta diversity to display the distance between samples[18]. Diversity describes the extent that individuals differ within each condition, while variation represents for the difference among samples. The distance within each condition increased after inflammation was induced, suggesting greater microbiome variation between individual rat. The distances between baseline and inflammation were highest (Figure 2 A,B). The principle coordinate analysis (PCoA) plots show how the beta-diversity patterns vary in the two IBD models. The region of the fitted ellipse expanded after DSS or TNBS treatment, and the distribution of samples shift to a non-overlapping pattern when IBDs induced (Figure 2 C,D). These results suggest that inflammation causes global changes of microbiome in guts, reducing diversity in each sample and increasing s between samples.

C. Taxonomy abundance analysis

We next performed taxonomy analysis to investigate the abundance of microbiota in control and DSS/TNBC treated rats (Figure 3). The abundance of *Clostridia* and *Bacteroidia*, which constitutes majority of the microbiota (approximately 80%), altered significantly after drug treatment in both the UC and CD models. The rest of the classes made up nearly 20% of the whole microbiota. The ratio of the majority to the minority remained consistent after induction of inflammation in both models. Although most of the shifted patterns are overlapped by two models, some abundance alternations were specific to each treatment. In the UC model, *Alphaproteobacteria* was enriched after DSS treatment, while in the CD's disease model, *Erysipelotrichi, Deltaproteobactria* and *Actinobacteria* were enriched after TNBS treatment.

D. Reconstitution of microbiota pattern

To find out the key species that contribute to the microbiome pattern shifting in IBD rat's model, we did the differential analysis to compare the microbiome signature between the health and induced IBD status. Linear differential analysis (LDA) is a method for characterizing two or more categories of objects. It was applied to the abundance of microbiome at genus or the lowest level by LDA Effect Size (LEfSe) after filtering out the least 0.001% genus. Using LDA scrore >=3.0 as the cutoff (p<=0.05), the remaining taxonomies are considered as the biomarkers for their corresponding condition (Baseline and the end day at DSS and TNBS treatment). Ranked by the LDA score of the biomarkers under health and inflammation status, the heatmaps show clear signature that biomarkers for health status get more enriched at baseline, and biomarkers for induced IBD become more abundant after DSS or TNBS treated.

Bacteriodes and Sutterella were enriched by DSS or TNBS treatments (Figure 4), consistent with previous reports that they were enriched in IBD patients.[19, 20] The abundance of RF39, Lachonospiraceae, Ruminococcus, Rikenellaceae, and Clodtridiales commonly reduced on the two IBD models. (Figure 4). In the DSS model, the abundance signature of

Bacteroides, Peptostreptococcaceae, Enterobaceriaceae, Prevotella, Akkermansia, Lachnospiraceae, Ruminococcus, Rikenellaceae is identical to that of UC patients.[19, 21–28] In the TNBS model, the abundance signature of Blautia, Fusobacterium, Odoribacter, Lactobacillus, Lactobacillus, Rikenellaceae, Ruminococcus, Lachnospiraceae, Oscillospira, Clostridiales is similar to that of the CD patients.[21, 26, 27, 29–33]

E. Functional Content Prediction

We performed metagenome prediction analysis based on the close-reference picked up OTUs to infer the microbial functions in IBD, testing 7 functional categories from KEGG Orthology (KO) level 1 and 40 from level 2. The abundance of functional contents on level one only shows slight difference between the two models. Metabolism consist to near half of the whole contents, while organismal systems take the smallest part. Differential analysis was applied to the KO level 2 groups by limma.

We selected the two most significantly different functions of each condition to demonstrate how the functions were regulated with the alteration of microbiota. (Figure 5) In both TNBS and DSS models, membrane transport (Level1: Environmental Information Processing) and cell motility (Level 1: Cellular Processes) were downregulated while glycan biosynthesis and metabolism (Level1: Metabolism) was upregulated compared with the healthy controls. In the DSS (UC) model, the level of carbohydrate metabolism (Level 1: Metabolism) increased most significantly among all the functions at KO Level 2, while in the TNBS (CD) model, energy metabolism had the most significant increase.

IV. Discussion

Animal models have been widely used to study human IBDs, and it is critical to understand how the models recapitulate the pathological features from IBDs. DSS (UC) and TNBS (CD) treatment are two of the most common models for IBD. Our study characterized the microbiome in the DSS and TNBS induced IBD rat models. The α -diversity of gut microbiome in individual samples was significantly reduced by inflammation. DSS and TNBS rat models recapitulate microbiome signatures consistent with clinical samples from human IBD patients. The functional analyses suggested possible microbial functional changes in response to gut inflammation. Therefore, IBD might cause profound changes in host-microbiome interactions.

Acknowledgment

We thank Duke Center for Genomic and Computational Biology and other members of the Shen lab for useful discussion and generous support. We thank DARPA (N66001-15-2-4059) for supporting the grant on this project.

References

- 1. Endo K, et al., Inflammatory bowel disease: IBD. Rinsho Byori, 2009 57(6): p. 527–32. [PubMed: 19621784]
- 2. Baumgart DC and Carding SR, Inflammatory bowel disease: cause and immunobiology. Lancet, 2007 369(9573): p. 1627–40. [PubMed: 17499605]

^{*} Research supported by R35GM122465 and DARPA N66001-15-2-4059.

3. Loh G and Blaut M, Role of commensal gut bacteria in inflammatory bowel diseases. Gut Microbes, 2012 3(6): p. 544–55. [PubMed: 23060017]

- 4. Mukhopadhya I, et al., IBD-what role do Proteobacteria play? Nat Rev Gastroenterol Hepatol, 2012 9(4): p. 219–30. [PubMed: 22349170]
- Rosshart SP, et al., Wild Mouse Gut Microbiota Promotes Host Fitness and Improves Disease Resistance. Cell, 2017 171(5): p. 1015-+. [PubMed: 29056339]
- 6. Antoniou E, et al., The TNBS-induced colitis animal model: An overview. Ann Med Surg (Lond), 2016 11: p. 9–15. [PubMed: 27656280]
- 7. Martin JC, Beriou G, and Josien R, Dextran Sulfate Sodium (DSS)-Induced Acute Colitis in the Rat. Methods Mol Biol, 2016 1371: p. 197–203. [PubMed: 26530802]
- 8. Chakravorty S, et al., A detailed analysis of 16S ribosomal RNA gene segments for the diagnosis of pathogenic bacteria. J Microbiol Methods, 2007 69(2): p. 330–9. [PubMed: 17391789]
- Janda JM and Abbott SL, 16S rRNA gene sequencing for bacterial identification in the diagnostic laboratory: pluses, perils, and pitfalls. J Clin Microbiol, 2007 45(9): p. 2761–4. [PubMed: 17626177]
- Caporaso JG, et al., QIIME allows analysis of high-throughput community sequencing data. Nat Methods, 2010 7(5): p. 335–6. [PubMed: 20383131]
- 11. DeSantis TZ, et al., Greengenes, a chimera-checked 16S rRNA gene database and workbench compatible with ARB. Appl Environ Microbiol, 2006 72(7): p. 5069–72. [PubMed: 16820507]
- 12. Wickham H, ggplot2: Elegant Graphics for Data Analysis. Ggplot2: Elegant Graphics for Data Analysis, 2009: p. 1–212.
- Segata N, et al., Metagenomic biomarker discovery and explanation. Genome Biol, 2011 12(6): p. R60. [PubMed: 21702898]
- 14. Starruss J, et al., Morpheus: a user-friendly modeling environment for multiscale and multicellular systems biology. Bioinformatics, 2014 30(9): p. 1331–2. [PubMed: 24443380]
- 15. Langille MGI, et al., Predictive functional profiling of microbial communities using 16S rRNA marker gene sequences. Nature Biotechnology, 2013 31(9): p. 814—+.
- 16. Kanehisa M, et al., KEGG: new perspectives on genomes, pathways, diseases and drugs. Nucleic Acids Res, 2017 45(D1): p. D353–D361. [PubMed: 27899662]
- 17. Ritchie ME, et al., limma powers differential expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Research, 2015 43(7).
- Lozupone C, et al., UniFrac: an effective distance metric for microbial community comparison.
 ISME J, 2011 5(2): p. 169–72. [PubMed: 20827291]
- Lucke K, et al., Prevalence of Bacteroides and Prevotella spp. in ulcerative colitis. Journal of Medical Microbiology, 2006 55(5): p. 617–624. [PubMed: 16585651]
- 20. Mukhopadhya I, et al., A comprehensive evaluation of colonic mucosal isolates of Sutterella wadsworthensis from inflammatory bowel disease. PLoS One, 2011 6(10): p. e27076. [PubMed: 22073125]
- 21. Dalal SR and Chang EB, The microbial basis of inflammatory bowel diseases. J Clin Invest, 2014 124(10): p. 4190–6. [PubMed: 25083986]
- 22. Kabeerdoss J, et al., Alterations of mucosal microbiota in the colon of patients with inflammatory bowel disease revealed by real time polymerase chain reaction amplification of 16S ribosomal ribonucleic acid. Indian J Med Res, 2015 142(1): p. 23–32. [PubMed: 26261163]
- 23. Lavelle A, et al., Spatial variation of the colonic microbiota in patients with ulcerative colitis and control volunteers. Gut, 2015 64(10): p. 1553–61. [PubMed: 25596182]
- 24. Matsuoka K and Kanai T, The gut microbiota and inflammatory bowel disease. Semin Immunopathol, 2015 37(1): p. 47–55. [PubMed: 25420450]
- 25. Nagao-Kitamoto H and Kamada N, Host-microbial Cross-talk in Inflammatory Bowel Disease. Immune Netw, 2017 17(1): p. 1–12. [PubMed: 28261015]
- 26. Papa E, et al., Non-invasive mapping of the gastrointestinal microbiota identifies children with inflammatory bowel disease. PLoS One, 2012 7(6): p. e39242. [PubMed: 22768065]
- 27. Pascal V, et al., A microbial signature for Crohn's disease. Gut, 2017 66(5): p. 813–822. [PubMed: 28179361]

28. Png CW, et al., Mucolytic bacteria with increased prevalence in IBD mucosa augment in vitro utilization of mucin by other bacteria. Am J Gastroenterol, 2010 105(11): p. 2420–8. [PubMed: 20648002]

- 29. Ghosh S, van Heel D, and Playford RJ, Probiotics in inflammatory bowel disease: is it all gut flora modulation? Gut, 2004 53(5): p. 620–622. [PubMed: 15082574]
- 30. Hold GL, et al., Role of the gut microbiota in inflammatory bowel disease pathogenesis: What have we learnt in the past 10 years? World Journal of Gastroenterology, 2014 20(5): p. 1192–1210. [PubMed: 24574795]
- 31. Jonkers D and Stockbrugger R, Probiotics and inflammatory bowel disease. Journal of the Royal Society of Medicine, 2003 96(4): p. 167–171. [PubMed: 12668702]
- 32. Morgan XC, et al., Dysfunction of the intestinal microbiome in inflammatory bowel disease and treatment. Genome Biol, 2012 13(9): p. R79. [PubMed: 23013615]
- 33. Santoru ML, et al., Cross sectional evaluation of the gut-microbiome metabolome axis in an Italian cohort of IBD patients. Sci Rep, 2017 7(1): p. 9523. [PubMed: 28842640]




Figure 1. α -diversity was calculated through shannon index of each sample. The boxplots depicted the shannon index within each group.




Figure 2. Beta-diversity of microbiota in IBD model rats.

- **A, B.** Boxplots of the unifrac distances within and between baseline and inflammation groups. **(p-value <0.01), *** (p-value <0.001)
- **C, D.** Positions of paired samples in UC and CD model are shown by the PCoA plots. The statistical ellipses are fitted to the range of scatters in each condition. The paired samples at baseline and inflammation condition are linked by purple lines.




Figure 3.
Relative abundance of taxonomy composition at class level. Each color of the stacked bar represents a class. with the least 0.5% of the total removed. Each column represents the sample at baseline or after IBD induced, the paired samples are in the same order under both conditions.




Figure 4. Microbiome signature of significantly differential taxonomies

The results of differential analysis on taxonomies are displayed as LDA score (threshold is set as 3, p<=0.05). Positive values represent the taxonomies are enriched when inflammation induced, while negative values represent that at baseline. The relative abundance of each taxonomy is quantile normalized by rows, shown in heatmaps as the color range between row min and row max. The taxonomy names assigned to red or green are intersected by inflammation or baseline enhanced bacteria respectively. In addition, the taxonomy names marked with '*' are consistent with clinical reports, sharing the same signature with the corresponding disease (UC and CD).




Figure 5.Metagenome of DSS and TNBS models were predicted by KO functions at level 1 and 2.
The pie charts stand for the composition of KO level 1 functions at inflammation status. The right panels of boxplots indicate the most 2 different level 2 functions under health or IBD-induced condition. The colors filled in boxes are corresponding to the colors of level 1 function slides in pie charts.