


Inflammatory bowel disease and immunonutrition: novel therapeutic approaches through modulation of diet and the gut microbiome

Larissa S. Celiberto,* Franziska A. Graef,* Genelle R. Healey,* Else S. Bosman,* Kevan Jacobson, Laura M. Sly and Bruce A. Vallance 

Department of Paediatrics, BC Children's Hospital, University of British Columbia, Vancouver, BC, Canada

doi:10.1111/imm.12939

Received 13 February 2018; revised 28 March 2018; accepted 18 April 2018.

*These authors contributed equally.

Correspondence: Dr Bruce A. Vallance, Department of Paediatrics, Division of Gastroenterology, University of British Columbia and BC Children's Hospital, CH.I.L.D. Foundation Chair in Paediatric Gastroenterology, Rm. K4-188, 4480 Oak Street Vancouver, BC V6H 3V4, Canada.
Email: bvallance@cw.bc.ca

or

Dr Laura M. Sly, Department of Paediatrics, Division of Gastroenterology, University of British Columbia and BC Children's Hospital, Rm. A5-142, 950 West 28th Avenue, Vancouver, BC V5Z 4H4, Canada.
Email: laurasly@mail.ubc.ca
Senior author: Bruce A. Vallance

Summary

Inflammatory bowel disease (IBD) is a chronic inflammatory condition of the gastrointestinal tract, thought to at least in part reflect an aberrant immune response to gut bacteria. IBD is increasing in incidence, particularly in populations that have recently immigrated to western countries. This suggests that environmental factors are involved in its pathogenesis. We hypothesize that the increase in IBD rates might reflect the consumption of an unhealthy Western diet, containing excess calories and lacking in key nutritional factors, such as fibre and vitamin D. Several recent studies have determined that dietary factors can dramatically influence the activation of immune cells and the mediators they release through a process called immunonutrition. Moreover, dietary changes can profoundly affect the balance of beneficial versus pathogenic bacteria in the gut. This microbial imbalance can alter levels of microbiota-derived metabolites that in turn can influence innate and adaptive intestinal immune responses. If the diet–gut microbiome disease axis does indeed underpin much of the ‘western’ influence on the onset and progression of IBD, then tremendous opportunity exists for therapeutic changes in lifestyle, to modulate the gut microbiome and to correct immune imbalances in individuals with IBD. This review highlights four such therapeutic strategies – probiotics, prebiotics, vitamin D and caloric restriction – that have the potential to improve and add to current IBD treatment regimens.

Keywords: fasting; gut; immunonutrition; microbiome; prebiotics; probiotics; vitamin D.

Introduction to inflammatory bowel disease

Inflammatory bowel disease (IBD) is an immune-mediated disease characterized by inflammation of the gastrointestinal (GI) tract. IBD encompasses both Crohn's disease and ulcerative colitis; conditions that are chronic and often progressive, but are most commonly associated with intermittent disease flares. The exact aetiology of IBD remains elusive. However, it is well accepted that IBD occurs at the intersection between genetic, immune and environmental factors. Genetically, IBD is one of the best-described complex diseases with > 200 gene variants associated with disease.¹ Many of these risk variants, for genes such as *NOD2* and *ATG16L1*, are predicted to cause defects in epithelial barrier function and/or bacterial recognition and clearance,² thereby implicating gut

microbes as drivers of IBD. On their own the currently identified gene risk variants only predispose individuals to disease. Clearly, modifiable factors such as gut microbes and diet play an important role in disease development, considering that IBD incidence has dramatically increased in developed countries over the second half of the twentieth century, pointing to a role for the ‘western’ lifestyle. Over the past decade, studies have repeatedly identified an imbalance in the gut microbiota, referred to as dysbiosis, in patients with IBD.³ Characterized by reduced microbial diversity and increased numbers of potential pathobionts, such as adherent/invasive *Escherichia coli*, it remains unclear whether these shifts in microbial communities are a result of inflammation, or instead reflect the high-sugar, high-fat, low-fibre diets common in developed countries. In either case, the

microbial dysbiosis seen in patients with IBD leads to altered levels of microbial metabolites within the gut, thereby, along with diet, inducing changes in host metabolism and function. Hence, although the current paradigm is that IBD occurs in genetically susceptible individuals due to an inappropriate immune response to enteric commensal microbes, an alternative hypothesis is that diet/aberrant microbiota initiates and/or promotes the damaging immune responses seen in IBD. Through this review, we will address how immunonutrition, referring to the effect of nutrients and metabolites on the immune system, is altered in IBD, and how manipulation of diet and the gut microbiome may offer new therapeutic strategies to treat IBD.

Current therapies for IBD

At present, treatments for individuals with IBD predominantly target their pathological immune responses rather than any potential causal factors. The traditional step-up approach to therapy moves sequentially through immunosuppressives (5-aminosalicylic acid), immunomodulators (azathioprine and methotrexate), and finally to biologics.^{4,5} Biologics used to treat IBD include antibodies that block inflammation by targeting the pro-inflammatory cytokines tumour necrosis factor- α (TNF- α) or interleukin-12 (IL-12)/23p40, or by preventing immune cell recruitment to the gut. The anti-TNF- α antibody (infliximab) revolutionized the treatment for Crohn's disease upon its introduction in 1998,^{6,7} and was subsequently found to be effective for the treatment of ulcerative colitis.⁸ A fully humanized version of the same antibody (adalimumab) is also used to treat both Crohn's disease⁹ and ulcerative colitis¹⁰ with similar efficacy to infliximab treatment.¹¹ Other newer biologics include antibodies that block $\alpha_4\beta_7$ (vedolizumab) or $\alpha_e\beta_7$ (etrolizumab) integrin-mediated immune cell recruitment into the gut,^{12–14} or antibodies that target the pro-inflammatory effects of IL-12/23p40 (ustekinumab).¹⁵ Additional biological therapies including anti-cytokine antibodies, anti-immune cell trafficking therapies, inhibitors of the Janus kinase pathway, and antisense oligonucleotides are all in various stages of clinical trials (reviewed in ref. 16).

Although representing the most widely accepted IBD therapies, these approaches have important limitations. For instance, most biologics only work in subsets of patients,¹⁷ and although they can prove highly effective for that subset, they can also induce substantial adverse effects.¹⁷ For example, steroids or biologics increase susceptibility to infections and cancer,^{18,19} and evidence suggests that their efficacy decreases over time in a substantial number of patients.²⁰ Moreover, it can be argued that steroids²¹ are not specifically targeting the underlying, causal abnormalities in immune cells or in the gut microbiome. Hence, we propose that it is

worthwhile examining non-genetic, modifiable contributors to IBD. As our understanding of the mechanisms underlying IBD expands, rational alternative approaches targeting the gut microbiome and immune system through dietary modulation are receiving increased attention.

Inflammatory bowel disease and the dysbiotic gut microbiome

The human GI tract harbours a vast array of microorganisms that provide many benefits to the health and well-being of the host.²² These microbes include bacteria, viruses, fungi and eukaryotes; collectively referred to as the microbiome (reviewed in ref. 23–26). Development of the microbiome begins in early infancy and is influenced by several factors including route of delivery (vaginal versus caesarean section), environment, breast feeding, diet, genetics, infections, antibiotic use, age and hygiene.^{27,28} Humans and their gut microbiota have co-evolved to possess a symbiotic relationship, wherein the microbes obtain a place to dwell, as well as nutrition, while they help their hosts by aiding in food digestion, out-competing potential pathogens, and releasing beneficial metabolites such as short-chain fatty acids (SCFA), notably acetate, propionate and butyrate. In contrast, when an imbalance in bacterial composition and a lack of bacterial diversity (termed dysbiosis) occurs, microbial functionality may become perturbed. Dysbiotic microbiota have been implicated in a number of diseases including colon cancer, obesity and type 2 diabetes.^{29–32} The gut microbiota of individuals with IBD is characterized by low microbial diversity,^{33,34} a reduced abundance of *Bifidobacterium* spp.,^{33,35} *Lactobacillus* spp.³⁴ and *Faecalibacterium prausnitzii*,^{33,35,36} and a higher abundance of pathobionts such as adherent/invasive *E. coli*^{37,38} and *Clostridium difficile*,³⁹ resulting in lower SCFA concentrations⁴⁰ compared with healthy individuals (Fig. 1).

At present, it remains unclear whether the dysbiotic microbiota found in many IBD patients truly plays a causative role or alternatively, is simply a reflection of the inflammatory and antimicrobial responses elicited during the course of disease. The truth probably lies somewhere in between. Several observations have indicated that intestinal dysbiosis might promote IBD pathogenesis, as inflammation is usually located in the distal ileum or colon, which are also the sites of highest bacterial abundance in the intestine. Moreover, studies using spontaneous and induced animal models of IBD have shown that animals develop little if any inflammation under germ-free (GF) conditions.^{41–43} Nonetheless, inflammation on its own seems to favour the typical dysbiosis seen in patients with IBD, i.e. depletion of *Firmicutes* and the expansion of *Enterobacteriaceae*, especially *E. coli* strains.^{44,45} This dysbiotic state is probably due to




Figure 1. Complex interactions that exist between diet, gut microbiota, colonocytes and immune cells during intestinal homeostasis and inflammation. (a) Intestinal homeostasis is associated with a healthier dietary pattern, microbial diversity, higher abundance of beneficial bacteria such as *Bifidobacterium* spp., *Faecalibacterium prausnitzii* and *Lactobacillus* spp., and increased short-chain fatty acid production, particularly butyrate. Butyrate possesses powerful anti-inflammatory properties as it promotes regulatory T (Treg) cell proliferation and enhances intestinal barrier function. Dendritic cells within the lamina propria sample commensal microbiota antigens within the intestinal lumen. In response to these commensal antigens, dendritic cells release transforming growth factor- β (TGF- β) which activates Treg cells to release interleukin-10 (IL-10) and TGF- β , leading to a more tolerant immune phenotype. A thicker mucus layer is another characteristic of intestinal homeostasis, providing a protective barrier between luminal bacteria and epithelial cells. Lastly, aryl hydrocarbon receptor (AhR) ligands, derived from fruits and vegetables, induce innate lymphoid cell 3 (ILC3) to produce IL-22, which helps maintain intestinal barrier function. (b) Conversely, intestinal inflammation is associated with a Western dietary pattern and bacterial dysbiosis (i.e. lower microbial diversity, reduced short-chain fatty acid production, lower abundance of beneficial bacteria, and higher abundance of pathobionts such as *Clostridium difficile* and *Escherichia coli*). A thinner, more patchy mucus layer provides less of a protective barrier between the luminal bacteria and colonocytes. Along with a reduction in expression of cellular tight junctions this leads to impaired intestinal barrier function resulting in leakage of bacterial products, such as lipopolysaccharides (LPS), from the intestinal lumen into the lamina propria. LPS bind to toll-like receptors (TLR) activating macrophages to produce tumour necrosis factor- α (TNF- α) which promotes T helper cell type 1 (Th1) to proliferate and release pro-inflammatory cytokines [i.e. TNF- α and interferon- γ (IFN- γ)] leading to inflammation, which also compromises intestinal barrier function. Additionally, a reduction in IL-10-producing Treg cells further contributes to intestinal inflammation.

changes in oxygen levels and other environmental features of the gut including the development of new nutritional niches only suitable for this group of bacteria.^{38,44,45} Considering the unique ability of *Enterobacteriaceae* to thrive proximal to inflamed tissues, these examples support the concept that microbial dysbiosis might be a consequence rather than a cause of inflammation in patients with IBD.

Inflammatory bowel disease and the Western diet: calorically dense, nutritionally scarce

Nutrition is a critical component of health that can impact the onset and progression of IBD.⁴⁶ Although healthy diets are thought to provide the body's caloric requirements from whole foods that are nutritiously diverse, modern (Western) dietary patterns consist of sugar- and fat-rich, high-calorie processed foods generally devoid of fibre. As such, Western diets typically contain few fruits, vegetables, legumes and whole grains.⁴⁷ Aside from lacking fibre, it is also notable that several foods present in the Western diet have been shown to have detrimental effects on the gut and the residing microbes or are associated with an increased risk of developing IBD. These foods include non-caloric artificial sweeteners, alcohol, preservatives and stabilizers, refined carbohydrates and high amounts of processed animal products, especially meat (as reviewed in ref. 48,49). Interestingly, most foodstuffs considered detrimental in the western style diet are also calorically dense foods, whereas beneficial high-fibre foods contain far fewer calories per gram. Western diets therefore lead to increased rates of obesity, in concert with a rise in the occurrence of colon cancer and inflammatory conditions such as type 2 diabetes and IBD.⁵⁰ Indeed, it is well documented that excess body fat leads to a state of systemic low-grade inflammation in humans,⁵¹ as excess nutrients can activate and fuel immune cells to drive inflammation^{52,53} as well as promote endoplasmic reticulum stress. Consistent with that, recent research suggests that limiting nutrients or even fasting can have the opposite effect on immune cells and thereby dampen inflammation.^{54,55}

Interestingly, the clinical course of IBD itself can impair nutrient uptake, as chronic gut inflammation, intestinal surgery and the adverse effects of IBD-specific medications can predispose individuals to deficiencies in certain micronutrients such as iron, calcium, zinc, magnesium, folic acid, vitamin B12, vitamin A and vitamin D.⁵⁶ Other dietary factors of importance and at risk of deficiency in some individuals with IBD are natural plant flavonoids and indoles.⁵⁷ Recent reviews provide additional insight into the other IBD-related micronutrient deficiencies mentioned above.^{58,59} However, among these deficiencies, the role of vitamin D (also known as the sunshine vitamin) in gut immunity is one of the best-characterized relationships in the literature, and is described later in this review.

Immunonutrition and its potential impact on IBD

As described above, immunonutrition refers to the effects that dietary factors can have on different aspects of the immune system as well as the microbiome. Within the GI tract, nutrients are likely to affect mucosal barrier function and cellular defence, as well as modulate local inflammation. For example, intestinal homeostasis in individuals consuming a healthy, balanced diet is normally maintained through the release of a variety of microbial metabolites, including SCFA (e.g. acetate, propionate and butyrate), which improve intestinal barrier function by providing energy for colonic epithelial cells (colonocytes), and by promoting regulatory T cell (Treg) function. SCFA are rapidly used by colonocytes or absorbed into the systemic circulation where they bind to the G protein-coupled receptors GPR41 and GPR43, mediating protective immunity by promoting epithelial cell production of cytokines and chemokines.⁶⁰ Butyrate is the best studied of the SCFA, acting as a key source of energy for colonocytes as well as possessing powerful anti-inflammatory properties. Anti-inflammatory activities include altering the maturation of dendritic cells (DC), increasing the number of Treg cells and levels of the anti-inflammatory cytokine IL-10, while inhibiting production of pro-inflammatory cytokines such as interferon- γ and IL-2.^{61–63} A healthy gut microbiome can also direct the production of the anti-inflammatory mediators transforming growth factor- β , retinoic acid and thymic stromal lymphopoietin, all of which help to maintain normal mucosal immune function and intestinal homeostasis. The intestinal immune system can also be directly modulated by products such as aryl hydrocarbon receptor (AhR) ligands, derived from dietary tryptophan as well as flavonoids and indoles from fruits and vegetables. The AhR ligands induce immune cells [innate lymphoid cells 3 and T helper type 17 (Th17) cells] to produce mediators such as IL-22 that help to maintain the epithelial barrier (Fig. 1).

In contrast, the typical western style diet that is high in sugar and saturated fats and low in fibre can lead to systemic low-grade inflammation, as a well-characterized consequence of obesity.⁶⁴ Although inflammation in adipose tissues and in the liver is a more established feature of obesity, the intestine is emerging as a key site for immunological changes that affect whole-body metabolism.⁶⁵ Specifically, microbial and dietary factors influence underlying innate and adaptive responses of the intestinal immune system. These responses can lead to disrupted intestinal barrier function, system inflammation, impaired glucose metabolism and can promote bacterial dysbiosis (as reviewed in ref. 66). These changes in the gut microbiota are sufficient to promote a state of low-grade chronic mucosal inflammation, thereby disrupting intestinal homeostasis. A lack of fruit and vegetables reduces

the levels of AhR ligands, reducing IL-22 production by innate lymphoid cells 3 and Th17 cells, and thereby weakening intestinal barrier integrity.⁶⁷ Microbial dysbiosis can also lead to reduced levels of IL-10, coupled with increased local levels of TNF- α , thereby promoting inflammation.⁶⁸ Impaired gut barrier function results in the leakage of luminal bacterial products such as lipopolysaccharides out of the intestine, which drives a shift in T-cell polarity towards interferon- γ -releasing Th1 cells.⁶⁹ The ensuing low-grade inflammation further compromises intestinal barrier function,⁷⁰ resulting in the characteristic leaky gut associated with diet-induced obesity (Fig. 1).

Although metabolic disorders and IBD are distinct conditions, it is notable that many of the pathological features seen in the intestine in response to high-fat diets (baseline inflammation, barrier dysfunction) are similar to those seen in, or that predispose to, IBD. For example, although nucleotide-binding oligomerization domain sensors are best known as innate receptors involved in recognizing intracellular bacteria, they also respond to endoplasmic reticulum stress,⁷¹ highlighting the potential for alterations in immunonutrition to play a role in IBD development. A reduction in vitamin D levels as well as AhR-activating compounds are also seen in patients with IBD, in concert with impaired mucosal healing.⁷²

Aside from immune cells, recent studies have found a key role for epithelial cell metabolism in controlling the gut microbiota, and specifically the overgrowth of *E. coli* pathobionts. As noted above, under healthy conditions, colonocytes use butyrate as a key energy source. Metabolizing butyrate consumes oxygen, thereby rendering surface colonocytes hypoxic and promoting the luminal growth of strict anaerobic bacteria such as *Firmicutes*.^{73,74} In contrast, intestinal inflammation typically results in a loss of *Firmicutes* (i.e. *F. prausnitzii*), leading to a reduction in butyrate production. The absence of butyrate forces colonocytes to obtain energy through the fermentation of glucose to lactate (fermentative metabolism), a process that does not consume oxygen and so increases oxygen levels in and near these cells to 3–10%. This increase in oxygen dramatically affects the make-up of local microbes, depleting strict anaerobes while permitting a bloom of facultative anaerobic bacteria such as the *E. coli* pathobionts^{73,74} that have been implicated in IBD. Taken together, the similarity between the intestinal pathophysiology seen in metabolic disorders, and those seen in IBD raises the possibility that some IBD patients develop their disease because of heightened susceptibility to environmentally driven dysregulated intestinal immunity, such as that caused by western-style diet or other factors that promote microbial dysbiosis in the GI tract. As counter-regulatory measures, we propose and discuss dietary immunomodulation – or immunonutrition

strategies that could help to alleviate and re-balance an inflamed gut (summarized in Fig. 2).

Changing the gut microbiome – probiotics as therapy

One approach to overcome the microbial dysbiosis seen in patients with IBD is through the oral or per enema delivery of beneficial gut microbes (probiotics) known to be lacking in IBD patients. As defined by the World Health Organization, probiotics are microorganisms that confer a health benefit to the host when administered in adequate amounts.^{75,76} Probiotics can be easily incorporated into the diet through the consumption of fermented foods (i.e. yoghurt, kefir, kimchi, sauerkraut), or consumed on a daily basis as a probiotic supplement. Studies attribute several health benefits to probiotics including direct effects such as producing SCFA (e.g. butyrate) and excluding pathogens from the gut by competition for space and nutrients, as well as indirect effects such as enhancing epithelial barrier function and promoting antimicrobial peptide production and secretory IgA. Moreover, probiotics have been shown to increase mucin secretion from intestinal goblet cells and to beneficially modulate the host immune system through the stimulation of anti-inflammatory cytokines such as IL-10 and transforming growth factor- β as well as stimulate the induction of Treg cells.^{77,78} The exact mechanism(s) by which probiotics exert these positive effects are unclear; however, it is clear that the efficacy of probiotics varies depending on the microbial strain used and the dose administered.

Microorganisms from the genera *Lactobacillus* and *Bifidobacterium* as well as the yeast *Saccharomyces boulardii* are among the most common probiotic candidates (Tables 1 and 2). Extensive *in vitro* research has shown that several *Lactobacillus* spp. exhibit anti-inflammatory effects, as primarily assessed by Toll-like receptor (TLR) activation.⁷⁹ For example, *Lactobacillus casei* Shirota treatment restores the normal stimulatory capacity of DC from patients with ulcerative colitis by reducing TLR2 and TLR4 expression.^{80,81} *Lactobacillus plantarum* CGMCC1258 increases tight junction protein levels and decreases permeability in the intestinal epithelial cell line, IPEC-J2. Moreover, this probiotic reduces IL-8 and TNF- α expression in intestinal porcine epithelial cells challenged by *E. coli* K88, possibly through a decrease in TLR expression, nuclear factor- κ B activation, and mitogen-activated protein kinase pathways.⁸²

In mouse models of intestinal inflammation, *Lactobacillus acidophilus* Bar 13 and *Bifidobacterium longum* Bar 33 promote the expansion of Treg cells and reduce the number of intraepithelial lymphocytes in 2,4,6-trinitrobenzene sulphonic acid-induced colitis.⁸³ In a similar model of murine colitis (2,4-dinitrobenzene sulphonic acid),




Figure 2. Summary of immunonutrition strategies that modulate the gut microbiota and help alleviate and re-balance an inflamed gut. DAI, disease activity index; DC, dendritic cells; HB, hydroxybutyrate; IFN, interferon; IL, interleukin; mTOR, mechanistic target of rapamycin; NF, nuclear factor; SCFA, short-chain fatty acids; Th, T helper cell; TLR, toll-like receptors; TNF, tumour necrosis factor; Treg, regulatory T cell.

L. casei DN-114 001 ameliorates disease severity through the induction and expansion of colonic CD4⁺ FoxP3⁺ Treg cells.⁸⁴ Other studies using mice with dextran sulphate sodium (DSS) colitis show that a combination of eight different probiotic strains (VSL#3) effectively reduces disease activity and colon inflammation including a significant reduction in inflammatory markers such as IL-1 β , nuclear factor- κ B and the neutrophil marker, myeloperoxidase.^{85–87} Similarly, administration of *L. plantarum* 299V prevents spontaneous colitis development in IL-10-deficient (*Il10*^{−/−}) mice,⁸⁸ and treatment with VSL#3 ameliorates colitis and overall disease activity in *Il10*^{−/−} mice.

Curiously, despite the broad success of probiotics in animal models of colitis, their effects in clinical IBD trials have been less successful, with only small subsets of

treated patients showing beneficial effects (reviewed in ref. 89–91). One reason behind this limited effect in patients with IBD may stem from the ‘one size fits all’ approach that has been commonly employed with probiotics. It is strongly believed that as an infant’s immune system matures, it develops a mutualistic relationship with the resident microbes in the intestine. This ensures that these resident gut microbes establish an environmental niche, as well as an immunological niche, that is recognized by the immune system as a long-term part of the host. In contrast, new microbes encountered after this relationship has developed are typically seen as foreign and are expelled. Hence, giving exogenous probiotic microbes to patients without defining whether there are environmental/immunological niches for those gut microbes may mean that the probiotics will be seen as

Table 1. Effect of prebiotic and/or probiotic interventions on gut immunity and microbiota in animal models of colitis

Animal model	Prebiotic/probiotic	Dose and duration	Effect on gut immunity	Effect on gut microbiota	Ref
HLA-B27 transgenic rats	IN/FOS (Synergy 1)	5 g/kg in drinking water; 7 weeks	↓ Histological & gross caecal scores, & IL-1 β . ↓ mucosal inflammation. ↑ TGF- β . ϕ IL-10	↑ <i>Bifidobacterium</i> spp. & <i>Lactobacillus</i> spp. ϕ SCFA conc.	107
	IN or FOS	8 g/kg; 12 weeks	↓ Histology score (particularly FOS) & IL-1 β in caecal tissue	↓ Total bacteria & <i>Bacteroides</i> spp. (FOS & IN). ↓ <i>Clostridium</i> cluster XI & XIVa, <i>Enterobacteriaceae</i> & <i>Clostridium difficile</i> & ↑ <i>Bifidobacterium</i> spp. & <i>Clostridium</i> cluster I (FOS). ↓ <i>Lactobacillus</i> spp. & <i>Clostridium</i> cluster IV (IN). ↑ total SCFA conc. (FOS & IN)	108
	IN and <i>Lactobacillus acidophilus</i> La-5 and <i>Bifidobacterium lactis</i> Bb	Unspecified; 8 weeks	↓ Histological score	↑ Microbial diversity & <i>Bifidobacterium animalis</i>	109
DSS-induced colitis in rats	scFOS or RS (type 3)	63 (FOS) or 115 (RS) g/kg; 14 days during DSS	↓ Macroscopic & histological scores (RS)	ϕ SCFA conc.	113
	OF/Inulin (Synergy 1) + <i>Bifidobacterium infantis</i> or OF/inulin only	0.5 g/day via oral gavage; 7 days before DSS & 7 days during DSS	↓ DAI, MPO activity & colonic IL-1 β levels (both groups)	↑ Caecal <i>Bifidobacterium</i> spp. & <i>Lactobacillus</i> spp. counts. ϕ in <i>Enterobacteriaceae</i> . ↑ succinate conc.	105
	Inulin	1% of total diet in drinking water; 9 days before DSS & 5 days during DSS	↓ Mucosal damage scores & MPO activity	↑ Counts of <i>Lactobacillus</i> spp. ϕ in <i>Bifidobacterium</i> spp.	112
	VSL#3 + wortmannin (PI3k/Akt inhibitor)	15 mg VSL#3 + 1.4 mg/kg wortmannin daily after DSS for 7 days	↓ DAI and MPO activity ↓ iNOS, Cox-2, NF-kB, TNF- α , IL-6, p-Akt ↑ IL-10		85

Table 1 (Continued)

Animal model	Prebiotic/probiotic	Dose and duration	Effect on gut immunity	Effect on gut microbiota	Ref
DSS-induced colitis in mice	FOS	1.5 g/mL bd via oral gavage; 7 days before DSS & 7 days during DSS	↓ DAI & ↑ colon length		106
	Fibre mix- GOS, FOS, oligofructose, gum arabic, soy polysaccharides, cellulose & RS	1.5% of total diet; 14 days before DSS & 7 days during DSS	↑ Colon length/weight ratio. ↑ in colonic IL-10 levels. ↑ in Treg/Th2+ Treg/Th17 ratio		163
	β-glucan: glucan, zymosan & curdian	25 mg/kg by oral gavage; 14 days before DSS	↑ Histological evidence of colitis (all β-glucan types). ↑ TNF-α, CCL-2 and IL-6 (curdian and zymosan). ↑ IL-10 (glucan)		164
	β-glucan	500 mg/kg or 1000 mg/kg by oral gavage; 7 days before DSS & 7 days during DSS	↓ DAI, spleen weight & colon shortening. ↓ MPO, malondialdehyde & nitrate (both doses). ↓ TNF-α, IL-6, iNOS & IL-1β		165
	Yeast β-glucan or β-glucan	5% of total diet; 7 days during DSS	↓ DAI, colon shortening & histological evidence of colitis. ↓ apoptosis rate & serum IgA, IgG and IgM. ↓ infiltration of macrophages & neutrophils. ↓ TNF-α, IL-6, IL-8, iNOS & COX-2. ↓ EPO & MPO in colonic tissue. ↑ expression of ZO-1, occludin, claudin-1 & JAM-1		166
IL-10-deficient mice	NRS, SFD-t, SFD-c or SCF	4% of total diet; Weaning until end of experiment (12 weeks)	↓ IL-6 & CXCL1 (NRS). ↓ TNF-α, IL-1β & IL-23 (SFD-t). ↓ IL-12p70, IL-6 & CXCL1 ($P < 0.05$). φ cytokines (SCF)	↓ <i>Prevotellaceae</i> , <i>Incertae sedis</i> XIV & <i>Lachnospiraceae</i> (NRS). ↑ <i>Incertae sedis</i> XIV, <i>Lachnospiraceae</i> & <i>Ruminococcaceae</i> , & ↓ <i>Lactobacillaceae</i> (SFD-t). ↑ <i>Porphyromonadaceae</i> & <i>Prevotellaceae</i> , ↓ <i>Lactobacillaceae</i> , <i>Incertae sedis</i> XIV, <i>Lachnospiraceae</i> & <i>Ruminococcaceae</i> (SFD-c). ↑ <i>Porphyromonadaceae</i> & <i>Prevotellaceae</i> (SCF)	167

Table 1 (Continued)

Animal model	Prebiotic/probiotic	Dose and duration	Effect on gut immunity	Effect on gut microbiota	Ref
TNBS-induced colitis in rats	FOS	1 g bd via intragastric catheter; 2 days before TNBS & 7 or 14 days after TNBS	↓ Gross inflammation score & MPO activity (14 days)	↑ Butyrate (7 days only), lactate & total lactic acid bacteria	110
	scFOS (2 types: DP4 and DP8)	60 g/kg/day; 7 days before TNBS & 3 or 10 days after TNBS	↑ Cecal weight and contents. Significant ↓ in macroscopic damage score. ↓ MPO activity (DP8 only)	φ SCFA conc.	111
	GOS	4 g/kg/day by oral gavage; 10 days before TNBS & 3 days after TNBS	φ Colonic damage score, oedema or myeloperoxidase activity	↑ <i>Bifidobacterium</i> spp. & total bacteria conc.	168
TNBS-induced colitis in mice	Mix1 (<i>Lactobacillus acidophilus</i> Bar 13 + <i>Bifidobacterium longum</i> Bar 33), Mix2 (<i>L. plantarum</i> Bar 10, <i>Streptococcus thermophilus</i> Bar 20, and <i>B. animalis</i> subsp. <i>lactis</i> Bar 30)	10 ⁹ CFU/day; 3 weeks	Significant ↓ in histological damage. ↓ CD4 ⁺ cells of IELs and LPLs (Mix1). ↑ Tregs and IL-10 (Mix1 and Mix2) ↓ TNF-α and (MCP)-1 (Mix1 and Mix2)		83
DNBS-induced colitis in mice	<i>Lactobacillus casei</i> DN – 114 001	200 μL of 10 ⁸ CFU/mL by oral gavage; 14 days before DNBS & 5 days after DNBS	↑ Foxp3 ⁺ CD4 ⁺ T cells		84

bd, twice daily; CCL, chemokine ligand; COX, cyclooxygenase; DAI, disease activity index; DNBS, 2,4-dinitrobenzene sulphonic acid; DSS, dextran sulphate sodium; EPO, eosinophil peroxidase; scFOS, short-chain fructo-oligosaccharides; G-CSF, granulocyte colony-stimulating factor; IELs, intraepithelial lymphocytes; IFN, interferon; Ig, immunoglobulin; IL, interleukin; IN, inulin; IP, interferon-γ-induced protein; KC, keratinocyte chemoattractant; LPLs, lamina propria lymphocytes; MCP-1, monocyte chemoattractant protein 1; MPO, myeloperoxidase; NRS, Corn-derived hydroxypropylated new resistant starch; iNOS, inducible nitric oxide synthase; RS, resistant starch; SFD-c, soluble fibre dextrin from corn; SFD-t, soluble fibre dextrin from tapioca; SCF, soluble corn fibre; SCEA, short-chain fatty acids; TGF, transforming growth factor; TLR, toll-like receptor; TNBS, 2,4,6-trinitrobenzene sulphonic acid; TNF, tumour necrosis factor. φ, no change, ↓, decrease/lower, ↑, increase/higher.

Table 2. Effect of prebiotic and/or probiotic interventions on gut immunity and microbiota in individuals with inflammatory bowel disease

Disease	Participants and duration	Probiotic/Prebiotic and dose	Effect on gut immunity	Effect on gut microbiota	Ref
CD	10 adults; 3 weeks	FOS; 15 g/day	↓ HBI. ↑ IL-10 positive CD11c ⁺ intestinal DC. ↓ IL-6 or IL-12 positive CD11c ⁺ intestinal DC. ↑ DC expression TLR2 & TLR4	↑ Faecal but not mucosal <i>Bifidobacterium</i> spp. ↓ total bacteria, <i>Bacteroides-Prevotella</i> or <i>Clostridium coccoides-Eubacterium rectale</i>	114
	103 adults; 4 weeks	IN/FOS (Synergy 1); 7.5 g bd	↑ DC IL-10 staining of rectal tissue. ↓ IL-12p40 production by intestinal DC or IL-6 positive DC. ↓ CDAI	↓ <i>Bifidobacterium</i> spp. or <i>Faecalibacterium prausnitzii</i>	121
	35 adults; 6 months	<i>Bifidobacterium longum</i> with IN/FOS (Synergy 1); 6 g bd	↓ CDAI, histological scores & TNF-α	↑ Mucosal <i>Bifidobacterium</i> spp.	115
	40 adults; 4 weeks	OF-IN; 10 g bd	Positive correlation between ↓ HBI & in ↑ <i>B. longum</i>	↓ <i>Ruminococcus gnavus</i> . ↑ <i>B. longum</i> . ↓ <i>Faecalibacterium prausnitzii</i>	116
	165 adults; 52 weeks	<i>Saccharomyces boulardii</i> ; 1 g/day	↓ CDAI, C-reactive protein, remission rate		169
	70 adults; 12 weeks	<i>Lactobacillus johnsonii</i> (LA1); 10 ¹⁰ CFU/day	↓ Endoscopic score, severe recurrence, CDAI		170
	19 adults; 2 weeks	IN/FOS (Synergy 1); 4 g tds	↓ Rachmilewitz scores. ↓ faecal calprotectin		119
	18 adults; 4 weeks	IN and FOS with <i>B. longum</i> ; 6 g bd	↓ Sigmoidoscopy score, mRNA levels for human β-defensins 2, 3 and 4, TNF-α & IL-1α. Biopsies showed ↓ inflammation & regeneration of epithelial tissue	↑ <i>Bifidobacterium</i> spp. in rectal mucosa	117
	16 adults; 8 weeks	Arabinogalactan/xilo-oligosaccharides with <i>Lactobacillus paracasei</i> (B21060); 6 g bd	↓ Serum IL-6 & IL-8. ↓ in general well-being, rectal bleeding & number of bowel motions		118
	39 adults; 12 months	GOS with <i>Bifidobacterium breve</i> fermented milk (Yakult); 5.5 g/day	↓ Endoscopic score & MPO levels	↓ <i>Bacteroidaceae</i> . ↓ <i>Bifidobacterium</i> spp. ↓ faecal pH levels	120
UC	42 adults; 8 weeks	VSL#3; 10 ¹⁰ CFU bd	↓ DC and TLR2 expression. ↑ IL-10 and ↓ IL-12p40		171
	30 adults; 8 weeks	BIFICO; 1.26 g/day	↓ NF-κB, TNF-α and IL-1β, ↑ IL-10	↑ <i>Lactobacillus</i> spp. and <i>Bifidobacterium</i> spp.	172
	52 adults; 4 months	Lactulose; 10 g/day	↓ Clinical activity indices, endoscopic score or immunohistochemical parameters. ↑ QoL index in UC		122

bd, twice daily; CD, Crohn's disease; CDAI, Crohn's disease activity index; DC, dendritic cells; FOS, fructo-oligosaccharide; GOS, galacto-oligosaccharide; HBI, Harvey-Bradshaw index; IL, interleukin; IN, inulin; MPO, myeloperoxidase; NF-κB, nuclear factor-κB; OF, oligofructose; QoL, quality of life; tds, three times a day; TLR, Toll-like receptor; TNF, tumour necrosis factor; UC, ulcerative colitis. ↓, decrease/lower; ↑, increase/higher.

foreign and will be unable to take up permanent residence in the GI tracts of those patients. Similarly, the inflamed intestines of patients with IBD are inhospitable to probiotic microbes because of the exaggerated inflammatory and antimicrobial responses seen during disease. These responses clear new microbes, including potentially beneficial bacteria, rapidly from the intestine, often before they have the opportunity to work.

Clearly, new approaches to designing probiotics, and promoting their survival will be key to the future success

of this potential therapy. Additionally, engineered probiotics have been developed that produce and release the anti-inflammatory cytokine IL-10^{92,93} or trefoil factor as strategies to locally suppress intestinal inflammation and promote healing.⁹⁴ Moreover, recent insights regarding the make-up of the human microbiome should allow us to identify potential next-generation probiotic species with improved potential for colonizing the human GI tract. Recently, Maldonado-Gómez *et al.*⁹⁵ demonstrated that the microbe *B. longum* AH1206 was able to persist in

the intestines of a subset of individuals for at least 6 months after the administration without causing adverse effects or overtly altering the resident microbiota composition. This microbe's extended colonization is attributed to its ability to establish a nutritional niche related to genes involved in carbohydrate use. This finding suggests that the establishment of new probiotic microbes will depend on an individual's baseline microbiota as well as on the availability of nutritional resources, supporting the critical role of dietary substrates such as fermentable carbohydrates and prebiotics in permitting the long-term persistence of a probiotic strain.

Prebiotics – creating new niches for beneficial gut microbes

One strategy that has shown promise in promoting a healthier gut microbiome is the use of prebiotics. A prebiotic is 'a substrate that is selectively used by host microorganisms conferring a health benefit'.⁹⁶ Established prebiotics include inulin-type fructans (i.e. fructo-oligosaccharides, inulin and oligofructose), galacto-oligosaccharides and lactulose. Other fermentable carbohydrates that have shown prebiotic potential include resistant starch, β -glucans, arabinoxylan oligosaccharides, xylo-oligosaccharides, soy bean oligosaccharides, isomalto-oligosaccharides and pectin. Prebiotics are found naturally in foods (i.e. inulin is found in breads and cereals, onions, garlic and artichokes), are added to foods to increase their fibre content (i.e. inulin-containing yoghurts), or can be added to the diet in the form of powdered supplements. Prebiotics have the potential to create a new nutritional niche within the human GI tract, providing microbes with sufficient nutrients to establish residence. For example, inulin-type fructan prebiotics have been shown to increase the numbers of beneficial bacteria such as *Bifidobacterium* spp., *F. prausnitzii* and *Lactobacillus* spp.;^{97,98} while reducing pathobiont (*E. coli*) adherence to epithelial cells.⁹⁹ Prebiotic interventions can also lead to the increased production of SCFA that possess many beneficial properties as noted above.^{100,101} Prebiotics also regulate gut immunity, largely due to the indirect effects they have on gut microbiota and the metabolites they produce. However, prebiotics have also been shown to have direct effects on immune function *in vitro*, largely through their activation of innate immune pathways within DC and epithelial cells. As DC are constantly sampling the contents of the gut lumen, it has been proposed that prebiotics ligate the TLR found on the surface of DC¹⁰² leading to a cascade of immune responses. They could also bind to TLR on epithelial cells, thereby modulating barrier function,¹⁰³ and signalling to adjacent immune cells (i.e. DC and intraepithelial lymphocytes).¹⁰⁴ As data suggesting that prebiotics directly influence gut immunity have only been acquired from *in vitro* studies, this topic requires further investigation.

Prebiotic intervention studies using murine models of colitis have shown promise in modulating gut microbiota and host immune response (Table 1). The most frequently studied prebiotics in these studies are inulin-type fructans, which have been shown to significantly reduce disease activity index,^{105,106} histological and macroscopic damage scores,^{107–112} pro-inflammatory cytokine production (e.g. IL-1 β)^{105,107,108} and neutrophil myeloperoxidase levels,^{105,110–112} while increasing colon length/weight ratios.^{106,111} In some studies, the amelioration of gut inflammation was accompanied by changes in gut microbiota composition and SCFA production with increases in microbial diversity,¹⁰⁹ *Bifidobacterium* spp.,^{105,107–109} *Lactobacillus* spp.,^{105,107,112} total SCFA¹⁰⁸ and butyrate concentrations¹¹⁰ being observed. Notably, the effects of providing these prebiotics have not always been consistent, with some studies showing no changes in *Bifidobacterium* spp. concentrations¹¹² or SCFA production.^{107,111,113} The heterogeneity in study results may reflect differences in prebiotic type, dose, intervention length, the colitis model used, and the microbial environment of the respective animal facility. The immunomodulatory capacity of other prebiotics, such as galacto-oligosaccharides and β -glucans, has also been investigated using experimental models of colitis; the effects that these and other prebiotics have on gut immunity and microbiota are summarized in Table 1.

Compared with the number of prebiotic studies performed in mice, far fewer intervention studies have been conducted in patients with IBD (Table 2). However, studies have demonstrated that prebiotics facilitate reduced disease activity scores (Crohn's disease activity index and Harvey Bradshaw index),^{114–116} inflammation on biopsy,¹¹⁷ pro-inflammatory cytokine production (e.g. TNF- α , IL-6 and IL-8),^{115,117,118} faecal calprotectin¹¹⁹ and myeloperoxidase,¹²⁰ and an increase in IL-10-positive CD11c⁺ DC.^{114,121} Conversely, some studies have demonstrated no change in IL-6- and IL-12-producing CD11c⁺ DC¹¹⁴ or disease activity indices.^{119,121,122} Additionally, inconsistencies in the gut microbiota modulating effects of prebiotics have been observed in patients with IBD. Some studies have shown that prebiotics increase *Bifidobacterium* spp. concentrations in patients with IBD,^{114,115,117} whereas others showed no effect on *Bifidobacterium* spp.^{120,121} or *F. prausnitzii*.^{116,121} Marked differences in prebiotic type, dose and intervention length, as well as the participant's disease severity, type and location, and characteristics such as age, gender, habitual dietary intakes and baseline gut microbiota composition, are likely to have contributed to the different effects observed between studies.

Although prebiotic interventions generally do not lead to adverse outcomes in patients with IBD, the totality of evidence does not suggest that prebiotics consistently improve IBD outcomes. It is possible that a personalized

approach to prebiotic interventions is required as it is likely that a universal approach may be responsible, in part, for the inconsistencies in results. Prebiotic interventions during IBD relapse show great promise. However, additional, well-designed, randomized controlled trials are required before this alternative, complementary treatment option can be routinely recommended. Future research is necessary to define the ideal prebiotic type/s, dose and administration regimen (i.e. in combination with a habitual or exclusion diet, and/or probiotics) for use in patients with IBD.

Other dietary interventions – vitamin D

Vitamin D, best known as the sunshine vitamin, is naturally present in a limited number of foods, such as fatty fish and eggs. It is also often provided as a supplement within milk and other common food products. In part based on dietary choices, as well as limited sun exposure due to an indoor lifestyle, use of sunscreen and/or geographic location, it is estimated that one-third of the healthy population in northern latitudes is vitamin D insufficient (30–75 nmol/L) or deficient (<30 nmol/L). Moreover, low vitamin D levels are among the most important nutrient deficiencies associated with the onset and progression of IBD. Roughly 68% of the patients with IBD have insufficient serum vitamin D levels, more than half of these patients are vitamin D deficient.¹²³ Vitamin D deficiency can result in increased IBD disease activity, as well as a higher chance of relapses and an increased risk of hospitalization and surgery during the course of disease. Moreover, vitamin D supplementation studies suggest that increasing vitamin D levels reduces disease activity while increasing the quality of life of patients with IBD (reviewed in ref. 123).

The beneficial effects of vitamin D appear to work through multiple mechanisms including regulating host gene transcription and modulating the gut microbiome. Vitamin D regulates a number of genes involved in maintaining intestinal homeostasis, including barrier function and antimicrobial peptide production. Several studies have also shown that vitamin D signalling can be a major determinant of an individual's gut microbiota composition, especially with regards to diversity. Compared with wild-type mice, the gut microbiota of vitamin D receptor (VDR) deficient mice show a loss of *Lactobacillus* spp. and increases in *Proteobacteria* spp. and *Bacteroidetes* spp.,¹²⁴ a composition that is markedly dysbiotic and similar to that in patients with IBD.¹²⁵ It is therefore not surprising that mice lacking VDR signalling or that are vitamin-D-deficient are highly susceptible to chemically or bacterially induced intestinal inflammation such as DSS colitis,¹²⁶ *Citrobacter rodentium*-induced colitis,^{127,128} *S. Typhimurium* infections¹²⁹ and inflammation-associated malignancies, such as colon cancer.¹³⁰

An example of how vitamin D signalling is pivotal for healthy interactions between the gut microbiota and intestinal immunity involves the SCFA butyrate. Studies have shown that butyrate increases the expression of VDR in epithelial cells *in vitro*, and unexpectedly, butyrate is also thought to bind directly to the VDR, thereby inducing the differentiation of Caco2 epithelial cells,¹³¹ and down-regulating TNF- α production in HT-29 cells.¹³² This implies that bacterially derived butyrate could be used to restore VDR-dependent transcriptional regulation, which is severely lacking in patients with IBD. Similar results have been obtained following administration of the probiotic strains *Lactobacillus rhamnosus* GG ATCC 53103 and *L. plantarum*, with these microbes increasing VDR protein levels in both mouse and human intestinal epithelial cells. Moreover, the role of probiotics in regulating VDR signalling was assessed *in vivo* using the *Salmonella*-induced colitis model.¹³³ Probiotic treatment reduced colitis manifestations in infected wild-type mice, whereas probiotics had no effect on *Vdr*^{-/-} mice¹³³ suggesting a role for the VDR in probiotic efficacy. Since patients with IBD are often vitamin D insufficient (or deficient), these findings may clarify why probiotics often do not show beneficial effects in clinical trials. The ability of probiotics to inhibit intestinal inflammation and bacterial infection may depend on the VDR signalling pathway, so future studies should evaluate if there is a link between VDR expression and probiotic efficacy in a clinical setting.

Besides its actions on the gut microbiota, vitamin D also impacts host immunity, including immune cell differentiation, migration and anti-inflammatory functions. The immunomodulatory effects of vitamin D supplementation in patients with IBD are reviewed by Mohammed *et al.*¹³⁴ Human monocytes and macrophages treated with vitamin D show enhanced antimicrobial ability through improved autophagy and increased production of antimicrobial peptides, while also exhibiting reduced TLR-mediated inflammatory responses upon recognition of microbial products.¹³⁵ Additionally, vitamin D is involved in DC differentiation and maturation, resulting in a tolerogenic phenotype.¹³⁶ Treatment of human DC with the active form of vitamin D (1,25(OH)₂D₃) inhibits the production of the cytokines IL-12 and IL-23, which are responsible for driving Th1/Th17 responses, respectively. On the other hand, immunosuppressive IL-10 production by DC is enhanced following stimulation with active vitamin D, which results in DC developing a tolerogenic phenotype and subsequent induction of tolerogenic Treg cell immunity (increase of FOXP3 Treg cells and IL-10 secretion). Vitamin D also impacts adaptive immunity by inhibiting the proliferation of naive CD4⁺ T cells as well as altering their functionality by suppressing the transcription of pro-inflammatory Th1 and Th17 cytokines like interferon- γ , TNF- α , IL-17 and IL-21. Vitamin D also

displays synergistic effects with antibodies targeting TNF- α in patients with IBD. Adequate levels of vitamin D before the start of therapy were shown to drastically increase the chance of patients reaching remission,¹³⁷ not only reducing the risk of the biological therapy failing, but also limiting the formation of antibodies that can neutralize the drug.¹³⁸

Taken together, it appears clear that deficiency in vitamin D, or defects in its receptor can predispose to overt inflammatory responses, dysbiosis, and a worsened clinical outcome in patients with IBD. Moreover, vitamin D appears indispensable for maintaining intestinal mucosal homeostasis through its effects promoting intestinal barrier function, mucosal immunity, and subsequently shaping the intestinal microbiota. Vitamin D is clearly an important player in the emerging field of immunonutrition with exciting potential to improve IBD therapies and patient quality of life. Taken together, this suggests that an increased emphasis on treating vitamin D deficiency could help to reduce the onset and severity of IBD.¹³⁹ Hence supplementation strategies based on the severity of vitamin D deficiency in the patient should be incorporated into the treatment plan of IBD patients.¹⁴⁰

Calorie restriction – taking fuel from the fire

Specific nutrients may be important in controlling IBD, but another major issue already mentioned is that excessive caloric intake on a daily basis creates metabolic pressure that negatively impacts whole body energy homeostasis, resulting in systemic inflammation.⁵¹ This can give rise to various pathological conditions including insulin resistance, type 2 diabetes and cardiovascular disease.¹⁴¹ Energy homeostasis, however, is not only important on an organismal level, but also and especially on a cellular level.¹⁴² Perpetual activation of intracellular nutrient sensors leads to the activation of intracellular pro-inflammatory pathways. For example, continuous activation of biosynthetic pathways congests the protein folding capacity of the endoplasmic reticulum, thereby promoting the unfolded protein response, which can in turn induce the pro-inflammatory transcription factor nuclear factor- κ B.¹⁴³ These mechanisms are not confined to a specific cell type, but it is in immune cells where activation of pro-inflammatory pathways due to nutrient excess has the most serious implications. It is now becoming increasingly clear that immune cells respond directly to differences in nutrient availability by modulating their effector functions. All major cell types of the immune system have been shown to undergo a switch in their metabolism dependent on the type of inflammatory stimulus and the availability of nutrients (mainly glucose), in their environment (reviewed in ref. 144,145).

Intracellular energy sensors determine cell fate through the integration of nutrient signals. One of the central

intracellular energy sensors is the mechanistic target of rapamycin (mTOR). Primarily as part of the mTORC1 protein complex, mTOR regulates cellular energy metabolism, autophagy and proliferation as well as differentiation and effector functions of immune cells. For example, strong and incessant mTOR activation in concert with strong T-cell receptor binding favours Th1 and Th17 cell differentiation whereas low or oscillatory mTOR activation with weak T-cell receptor binding leads to increased Treg cell differentiation. Hence, if caloric intake is excessive, overactive mTOR signalling leaves fewer opportunities for Treg cell differentiation.¹⁴⁶ This, together with many other immunomodulatory effects of a high calorie diet, can lead to an imbalance between pro-inflammatory and anti-inflammatory cell types with chronic inflammation as a result. This imbalance is characteristic of patients with IBD, who have been shown to have more circulating Th17 and fewer Treg cells compared with healthy controls.¹⁴⁷

Based on what is known about the impact of cellular metabolism on immune function, targeting mTOR could be a promising strategy to re-establish immunological homeostasis. Several pharmacological mTOR inhibitors exist, i.e. rapamycin and its analogues or metformin.¹⁴⁸ In a DSS model of colitis, treatment with an mTOR inhibitor alleviated intestinal inflammation by reducing the number of Th1 and Th17 cells and increasing the number of Treg cells in the lamina propria.¹⁴⁹ Furthermore, rapamycin has been successfully used in a paediatric population with refractory IBD.¹⁵⁰ However, these drugs have severe adverse effects.¹⁵¹ Therefore, it is worth noting that one of the most natural and potent suppressors of mTOR is nutrient scarcity. Indeed, periods of fasting or caloric restriction decrease mTOR activity, with the potential to create a more anti-inflammatory environment.^{54,152}

Different types of caloric restriction exist, including intermittent or every-other-day fasting and time-restricted feeding. Yet, the most likely to prove beneficial is a prolonged fast with minimal caloric consumption or a calorically restricted diet containing nutrients that mimic fasting (also known as a fasting mimicking diet). During a prolonged fast, an organism undergoes a metabolic switch from mainly carbohydrate to predominantly (stored) fat utilization. This changes the metabolism of most cells, from using glucose as a central nutrient, to instead using ketone bodies, the most abundant of which is β -hydroxybutyrate (β -HB).¹⁵³ In an animal model of multiple sclerosis, three cycles of a 3-day fasting mimicking diet were found to not only induce the apoptosis of autoimmune Th1 and Th17 effector T cells, but also to promote the expansion of the Treg cell population as well as the oligodendrocyte precursor pool. This supports the concept that caloric restriction is able to shift the immune environment from pro- to anti-inflammatory cell types.¹⁵⁴ Interestingly, mTOR suppression is also one of

the mechanisms of action of the anti-inflammatory cytokine IL-10. Interleukin-10 achieves mTOR suppression via DNA damage inducible transcript 4 (DDIT4) activation. Through this axis, IL-10 can metabolically reprogramme inflammatory macrophages towards a more anti-inflammatory phenotype.¹⁵⁵ Intriguingly, prolonged fasting has also been shown to dramatically increase DDIT4 expression,¹⁵⁶ indicating another possible link for fasting-mediated immunomodulation.

Additionally, the fasting-induced ketone body β -HB exhibits a host of beneficial effects, including anti-inflammatory properties. For example, β -HB inhibits NLRP3 inflammasome activation in response to danger-associated molecular patterns.¹⁵⁷ This reduces the release of the pro-inflammatory cytokine IL-1 β both *in vitro* and *in vivo*, which may in part be facilitated through SIRT3-mediated activation of superoxide dismutase 2.¹⁵⁸ β -HB also acts as a histone deacetylase inhibitor, so promoting histone hyperacetylation and the subsequent induction of stress response genes, such as the transcription factor forkhead box O3 (FOXO3).¹⁵⁹ Moreover, β -HB up-regulates autophagy, which can be defective in patients with IBD, through FOXO3-mediated regulation of important autophagy genes such as *ATG4*, *ATG5* or *LC3* but also by indirectly inhibiting mTOR.^{160,161} Moreover, fasting is known to change the composition of the gut microbiome in mice and most likely in humans. Although it still needs to be determined on a larger scale how fasting modulates the intestinal microbiome, isolated studies have shown an increase in *F. prausnitzii* and *Akkermansia muciniphila*, both of which have been characterized as reflective of an anti-inflammatory intestinal environment.¹⁶²

Taken together, it seems likely that any substantial reduction in calories will dampen inflammation. Hence, interventions that reduce caloric intake could prove a beneficial complementary treatment for autoimmune and auto-inflammatory diseases such as IBD, where the immune system is chronically activated. As suppression of immunity could however impair protective responses to pathogens, more research is needed to establish if prolonged periods of fasting or caloric restriction have any negative effects on susceptibility or clearance of infectious organisms. It is also important to assess each patient's nutritional status before recommending a prolonged fast. Malnourished patients as well as children with intestinal diseases who are at risk of developing an inadequate nutritional status that would negatively impact their growth and development should not be fasted. Clinical studies should determine which types of caloric restriction are the safest and promise the best clinical outcomes.

Conclusions

Current IBD therapies focus on suppressing immune responses and restoring intestinal barriers, but we propose

that nutritional and microbiological interventions that are able to protect the gut against damaging inflammation offer the next source of IBD therapies. Elucidating the impact of nutrient intake on gut immunity, whether through direct effects on host immune cells or indirect effects on the gut microbiome, will provide us with critical information essential to promote gut health. Moreover, this understudied and underexploited area of research provides a tractable system that can be used to reduce damaging intestinal inflammation. Herein, we describe current paradigms as well as recent advances demonstrating that nutritional interventions, including modification of the gut microbiome by prebiotics and probiotics, as well as caloric restriction and supplementation with vitamin D, can modulate the mucosal immune response and suppress the damaging inflammation that characterizes IBD.

Disclosures

The authors declare no competing interests.

References

- Brant SR, Okou DT, Simpson CL, Cutler DJ, Haritunians T, Bradfield JP *et al.* Genome-wide association study identifies african-specific susceptibility loci in African Americans with inflammatory bowel disease. *Gastroenterology* 2017; **152**:206–17. e2.
- Knights D, Lassen KG, Xavier RJ. Advances in inflammatory bowel disease pathogenesis: linking host genetics and the microbiome. *Gut* 2013; **62**:1505–10.
- Sartor RB. Mechanisms of disease: pathogenesis of Crohn's disease and ulcerative colitis. *Nat Clin Pract Gastroenterol Hepatol* 2006; **3**:390–407.
- Strober W, Fuss I, Mannon P. The fundamental basis of inflammatory bowel disease. *J Clin Invest* 2007; **117**:514–21.
- Kaser A, Zeissig S, Blumberg RS. Inflammatory bowel disease. *Annu Rev Immunol* 2010; **28**:573–621.
- Hanauer SB, Feagan BG, Lichtenstein GR, Mayer LE, Schreiber S, Colombel JF *et al.* Maintenance infliximab for Crohn's disease: the ACCENT I randomised trial. *Lancet* 2002; **359**:1541–9.
- Colombel JF, Sandborn WJ, Reinisch W, Mantzaris GJ, Kornbluth A, Rachmilewitz D *et al.* Infliximab, azathioprine, or combination therapy for Crohn's disease. *N Engl J Med* 2010; **362**:1383–95.
- Colombel JF, Rutgeerts P, Reinisch W, Esser D, Wang Y, Lang Y *et al.* Early mucosal healing with infliximab is associated with improved long-term clinical outcomes in ulcerative colitis. *Gastroenterology* 2011; **141**:1194–201.
- Rutgeerts P, Van Assche G, Sandborn WJ, Wolf DC, Geboes K, Colombel J *et al.* Adalimumab induces and maintains mucosal healing in patients with Crohn's disease: data from the EXTEND trial. *Gastroenterology* 2012; **142**:1102–11. e2.
- Sandborn WJ, van Assche G, Reinisch W, Colombel J, D'Haens G, Wolf DC *et al.* Adalimumab induces and maintains clinical remission in patients with moderate-to-severe ulcerative colitis. *Gastroenterology* 2012; **142**:257–65. e3.
- Da W, Zhu J, Wang L, Lu Y. Adalimumab for Crohn's disease after infliximab treatment failure. *Eur J Gastroenterol Hepatol* 2013; **25**:885–91.
- Sandborn WJ, Feagan BG, Rutgeerts P, Hanauer S, Colombel J-F, Sands BE *et al.* Vedolizumab as induction and maintenance therapy for Crohn's disease. *N Engl J Med* 2013; **369**:711–21.
- Feagan BG, Rutgeerts P, Sands BE, Hanauer S, Colombel J-F, Sandborn WJ *et al.* Vedolizumab as induction and maintenance therapy for ulcerative colitis. *N Engl J Med* 2013; **369**:699–710.
- Wei X, Gibiansky L, Wang Y, Fuh F, Erickson R, O'Byrne S *et al.* Pharmacokinetic and pharmacodynamic modeling of serum etrolizumab and circulating β 7 receptor occupancy in patients with ulcerative colitis. *J Clin Pharmacol* 2018; **58**:386–98.
- Feagan BG, Sandborn WJ, Gasink C, Jacobstein D, Lang Y, Friedman JR *et al.* Ustekinumab as induction and maintenance therapy for Crohn's disease. *N Engl J Med* 2016; **375**:1946–60.

- 16 Atreya R, Neurath MF. Current and future targets for mucosal healing in inflammatory bowel disease. *Visc Med* 2017; **33**:82–8.
- 17 Ben-Horin S, Mao R, Chen M. Optimizing biologic treatment in IBD: objective measures, but when, how and how often? *BMC Gastroenterol* 2015; **15**:178.
- 18 Renna S, Cottone M, Orlando A. Optimization of the treatment with immunosuppressants and biologics in inflammatory bowel disease. *World J Gastroenterol* 2014; **20**:9675.
- 19 Stallmach A, Hagel S, Bruns T. Adverse effects of biologics used for treating IBD. *Best Pract Res Clin Gastroenterol* 2010; **24**:167–82.
- 20 Roda G, Jharap B, Neeraj N, Colombel J-F. Loss of response to anti-TNFs: definition, epidemiology, and management. *Clin Transl Gastroenterol* 2016; **7**:e135.
- 21 Bielefeldt K, Davis B, Binion DG. Pain and inflammatory bowel disease. *Inflamm Bowel Dis* 2009; **15**:778–88.
- 22 Cénit MC, Matzaraki V, Tighelela EF, Zhernakova A. Rapidly expanding knowledge on the role of the gut microbiome in health and disease. *Biochim Biophys Acta* 2014; **1842**:1981–92.
- 23 Caballero S, Pamer EG. Microbiota-mediated inflammation and antimicrobial defense in the intestine. *Annu Rev Immunol* 2015; **33**:227–56.
- 24 Pfeiffer JK, Virgin HW. Viral immunity. Transkingdom control of viral infection and immunity in the mammalian intestine. *Science* 2016; **351**:aad5872.
- 25 Underhill DM, Pearlman E. Immune interactions with pathogenic and commensal fungi: a two-way street. *Immunity* 2015; **43**:845–58.
- 26 Belkaid Y, Hand TW. Role of the microbiota in immunity and inflammation. *Cell* 2014; **157**:121–41.
- 27 Martin R, Nauta AJ, Amor Ben K, Knippels LMJ, Knol J, Garssen J. Early life: gut microbiota and immune development in infancy. *Beneficial Microbes* 2010; **1**:367–82.
- 28 Dominguez-bello MG, Costello EK, Contreras M, Magris M, Hidalgo G. Delivery mode shapes the acquisition and structure of the initial microbiota across multiple body habitats in newborns. *Proc Natl Acad Sci* 2010; **107**:11971–5.
- 29 Ley RE, Turnbaugh PJ, Klein S, Gordon JL. Microbial ecology: human gut microbes associated with obesity. *Nature* 2006; **444**:1022–3.
- 30 Walters WA, Xu Z, Knight R. Meta-analyses of human gut microbes associated with obesity and IBD. *FEBS Lett* 2014; **588**:4223–33.
- 31 Wu X, Ma C, Han L, Nawaz M, Gao F, Zhang X *et al.* Molecular characterisation of the faecal microbiota in patients with type II diabetes. *Curr Microbiol* 2010; **61**:69–78.
- 32 Ohgishi S, Sudo K, Kobayashi D, Takahashi O, Takahashi T, Asahara T *et al.* Changes of the intestinal microbiota, short chain fatty acids, and fecal pH in patients with colorectal cancer. *Dig Dis Sci* 2013; **58**:1717–26.
- 33 Andoh A, Kuzuoka H, Tsujikawa T, Nakamura S, Hirai F, Suzuki Y *et al.* Multicenter analysis of fecal microbiota profiles in Japanese patients with Crohn's disease. *J Gastroenterol* 2012; **47**:1298–307.
- 34 Ott SJ, Musfeldt M, Wenderoth DF, Hampe J, Brant O, Fölsch UR *et al.* Reduction in diversity of the colonic mucosa associated bacterial microflora in patients with active inflammatory bowel disease. *Gut* 2004; **53**:685–93.
- 35 Joossens M, Huys G, Cnockaert M, De Preter V, Verbeke K, Rutgeerts P *et al.* Dysbiosis of the faecal microbiota in patients with Crohn's disease and their unaffected relatives. *Gut* 2011; **60**:631–7.
- 36 Sokol H, Seksik P, Furet JP, Firmesse O, Nion-Larmurier I, Beaugerie L *et al.* Low counts of *Faecalibacterium prausnitzii* in colitis microbiota. *Inflamm Bowel Dis* 2009; **15**:1183–9.
- 37 Sokol H, Lepage P, Seksik P, Doré J, Marteau P. Temperature gradient gel electrophoresis of fecal 16S rRNA reveals active *Escherichia coli* in the microbiota of patients with ulcerative colitis. *J Clin Microbiol* 2006; **44**:3172–7.
- 38 Darfeuille-Michaud A, Boudeau J, Bulois P, Neut C, Glasser AL, Barnich N *et al.* High prevalence of adherent-invasive *Escherichia coli* associated with ileal mucosa in Crohn's disease. *Gastroenterology* 2004; **127**:412–21.
- 39 Rodemann JF, Dubberke ER, Reske KA, Seo DH, Stone CD. Incidence of *Clostridium difficile* infection in inflammatory bowel disease. *Clin Gastroenterol Hepatol* 2007; **5**:339–44.
- 40 Huda-Faujan N, Abdulamir AS, Fatimah AB, Anas OM, Shuhaimi M, Yazid AM *et al.* The impact of the level of the intestinal short chain fatty acids in inflammatory bowel disease patients versus healthy subjects. *Open Biochem J* 2010; **4**:53–8.
- 41 Taurog JD, Richardson JA, Croft JT, Simmons WA, Zhou M, Fernández-Sueiro JL *et al.* The germfree state prevents development of gut and joint inflammatory disease in HLA-B27 transgenic rats. *J Exp Med* 1994; **180**:2359–64.
- 42 Dianda L, Hanby AM, Wright NA, Sebesteny A, Hayday AC, Owen MJ. T cell receptor- $\alpha\beta$ -deficient mice fail to develop colitis in the absence of a microbial environment. *Am J Pathol* 1997; **150**:91–7.
- 43 Hudcovic T, Stěpánková R, Cebra J, Tlaskalová-Hogenová H. The role of microflora in the development of intestinal inflammation: acute and chronic colitis induced by dextran sulfate in germ-free and conventionally reared immunocompetent and immunodeficient mice. *Folia Microbiol (Praha)* 2001; **46**:565–72.
- 44 Gophna U, Sommerfeld K, Gophna S, Doolittle WF, Veldhuyzen van Zanten SJO. Differences between tissue-associated intestinal microfloras of patients with Crohn's disease and ulcerative colitis. *J Clin Microbiol* 2006; **44**:4136–41.
- 45 Frank DN, St Amand AL, Feldman RA, Boedeker EC, Harpaz N, Pace NR. Molecular-phylogenetic characterization of microbial community imbalances in human inflammatory bowel diseases. *Proc Natl Acad Sci* 2007; **104**:13780–5.
- 46 Nguyen DL, Limketkai B, Medici V, Saire Mendoza M, Palmer L, Bechtold M. Nutritional strategies in the management of adult patients with inflammatory bowel disease: dietary considerations from active disease to disease remission. *Curr Gastroenterol Rep* 2016; **18**:55.
- 47 Briefel RR, Johnson CL. Secular trends in dietary intake in the United States. *Annu Rev Nutr* 2004; **24**:401–31.
- 48 Knight-Sepulveda K, Kais S, Santaolalla R, Abreu MT. Diet and inflammatory bowel disease. *Gastroenterol Hepatol (N Y)* 2015; **11**:511–20.
- 49 Statovci D, Aguilera M, MacSharry J, Melgar S. The impact of western diet and nutrients on the microbiota and immune response at mucosal interfaces. *Front Immunol* 2017; **8**:838.
- 50 Alwan A, MacLean DR, Riley LM, D'Espaignet ET, Mathers CD, Stevens GA *et al.* Monitoring and surveillance of chronic non-communicable diseases: progress and capacity in high-burden countries. *Lancet* 2010; **376**:1861–8.
- 51 Reilly SM, Saltiel AR. Adapting to obesity with adipose tissue inflammation. *Nat Rev Endocrinol* 2017; **13**:633–43.
- 52 Lyons C, Kennedy E, Roche H. Metabolic inflammation – differential modulation by dietary constituents. *Nutrients* 2016; **8**:247.
- 53 Cohen S, Danzaki K, MacIver NJ. Nutritional effects on T-cell immunometabolism. *Eur J Immunol* 2017; **47**:225–35.
- 54 Longo VD, Mattson MP. Fasting: molecular mechanisms and clinical applications. *Cell Metab* 2014; **19**:181–92.
- 55 Longo VD, Panda S. Fasting, circadian rhythms, and time-restricted feeding in healthy lifespan. *Cell Metab* 2016; **23**:1048–59.
- 56 Owczarek D, Rodacki T, Domagala-Rodacka R, Cibor D, Mach T. Diet and nutritional factors in inflammatory bowel diseases. *World J Gastroenterol* 2016; **22**:895–905.
- 57 Joeris T, Müller-Luda K, Agace WW, Mowat AM. Diversity and functions of intestinal mononuclear phagocytes. *Mucosal Immunol* 2017; **10**:845–64.
- 58 Massironi S, Rossi RE, Cavalcoli FA, Della Valle S, Fraquelli M, Conte D. Nutritional deficiencies in inflammatory bowel disease: therapeutic approaches. *Clin Nutr* 2013; **32**:904–10.
- 59 Hwang C, Ross V, Mahadevan U. Micronutrient deficiencies in inflammatory bowel disease: from A to zinc. *Inflamm Bowel Dis* 2012; **18**:1961–81.
- 60 Kim MH, Kang SG, Park JH, Yanagisawa M, Kim CH. Short-chain fatty acids activate GPR41 and GPR43 on intestinal epithelial cells to promote inflammatory responses in mice. *Gastroenterology* 2013; **145**:396–406.
- 61 Meijer K, de Vos P, Priebe MG. Butyrate and other short-chain fatty acids as modulators of immunity: what relevance for health? *Curr Opin Clin Nutr Metab Care* 2010; **13**:715–21.
- 62 Millard AL, Mertes PM, Ittelet D, Villard F, Jeannesson P, Bernard J. Butyrate affects differentiation, maturation and function of human monocyte-derived dendritic cells and macrophages. *Clin Exp Immunol* 2002; **130**:245–55.
- 63 Furusawa Y, Obata Y, Fukuda S, Endo TA, Nakato G, Takahashi D *et al.* Commensal microbe-derived butyrate induces the differentiation of colonic regulatory T cells. *Nature* 2013; **504**:446–50.
- 64 Manzel A, Muller DN, Hafler DA, Erdman SE, Linker RA, Kleinewietfeld M. Role of “western diet” in inflammatory autoimmune diseases. *Curr Allergy Asthma Rep* 2014; **14**:404.
- 65 Winer DA, Winer S, Dranse HJ, Lam TKT. Immunologic impact of the intestine in metabolic disease. *J Clin Invest* 2017; **127**:33–42.
- 66 Winer DA, Luck H, Tsai S, Winer S. The intestinal immune system in obesity and insulin resistance. *Cell Metab* 2016; **23**:413–26.
- 67 Spencer SP, Belkaid Y. Dietary and commensal derived nutrients: shaping mucosal and systemic immunity. *Curr Opin Immunol* 2012; **24**:379–84.
- 68 Thevaranjan N, Puchta A, Schulz C, Naidoo A, Szamosi JC, Verschoor CP *et al.* Age-associated microbial dysbiosis promotes intestinal permeability, systemic inflammation, and macrophage dysfunction. *Cell Host Microbe* 2017; **21**:455–66. e4.
- 69 Ilan Y. Leaky gut and the liver: a role for bacterial translocation in nonalcoholic steatohepatitis. *World J Gastroenterol* 2012; **18**:2609.
- 70 Cox AJ, West NP, Cripps AW. Obesity, inflammation, and the gut microbiota. *Lancet Diabetes Endocrinol* 2015; **3**:207–15.
- 71 Keestra-Gounder AM, Byndloss MX, Seyffert N, Young BM, Chávez-Arroyo A, Tsai AY *et al.* NOD1 and NOD2 signalling links ER stress with inflammation. *Nature* 2016; **532**:394–7.
- 72 Basson A, Trotter A, Rodriguez-Palacios A, Cominelli F. Mucosal interactions between genetics, diet, and microbiome in inflammatory bowel disease. *Front Immunol* 2016; **7**:290.

- 73 Byndloss MX, Olsan EE, Rivera-Chávez F, Tiffany CR, Cevallos SA, Lokken KL *et al.* Microbiota-activated PPAR- γ signaling inhibits dysbiotic *Enterobacteriaceae* expansion. *Science* 2017; **357**:570–5.
- 74 Litvak Y, Byndloss MX, Tsois RM, Bäumlér AJ. Dysbiotic Proteobacteria expansion: a microbial signature of epithelial dysfunction. *Curr Opin Microbiol* 2017; **39**:1–6.
- 75 Probiotics in food Health and nutritional properties and guidelines for evaluation FAO FOOD AND NUTRITION PAPER [WWW document]. URL <http://www.fao.org/3/a-a0512e.pdf> [accessed on 6 January 2018].
- 76 Hill C, Guarner F, Reid G, Gibson GR, Merenstein DJ, Pot B *et al.* The international scientific association for probiotics and prebiotics consensus statement on the scope and appropriate use of the term probiotic. *Nat Rev Gastroenterol Hepatol* 2014; **11**:506–14.
- 77 Lebeer S, Vanderleyden J, De Keersmaecker SCJ. Genes and molecules of lactobacilli supporting probiotic action. *Microbiol Mol Biol Rev* 2008; **72**:728–64.
- 78 Borchers AT, Selmi C, Meyers FJ, Keen CL, Gershwin ME. Probiotics and immunity. *J Gastroenterol* 2009; **44**:26–46.
- 79 Wells JM. Immunomodulatory mechanisms of lactobacilli. *Microb Cell Fact* 2011; **10** (Suppl 1):S17.
- 80 Mann ER, You J, Horneffer-van der Sluis V, Bernardo D, Omar Al-Hassi H, Landy J *et al.* Dysregulated circulating dendritic cell function in ulcerative colitis is partially restored by probiotic strain *Lactobacillus casei* shirota. *Mediators Inflamm* 2013; **2013**:1–12.
- 81 Mann ER, Bernardo D, Ng SC, Rigby RJ, Al-Hassi HO, Landy J *et al.* Human gut dendritic cells drive aberrant gut-specific T-cell responses in ulcerative colitis, characterized by increased IL-4 production and loss of IL-22 and IFN γ . *Inflamm Bowel Dis* 2014; **20**:2299–307.
- 82 Wu Y, Zhu C, Chen Z, Chen Z, Zhang W, Ma X *et al.* Protective effects of *Lactobacillus plantarum* on epithelial barrier disruption caused by enterotoxigenic *Escherichia coli* in intestinal porcine epithelial cells. *Vet Immunol Immunopathol* 2016; **172**:55–63.
- 83 Roselli M, Finamore A, Nuccitelli S, Carnevali P, Brigidi P, Vitali B *et al.* Prevention of TNBS-induced colitis by different *Lactobacillus* and *Bifidobacterium* strains is associated with an expansion of $\gamma\delta$ T and regulatory T cells of intestinal intraepithelial lymphocytes. *Inflamm Bowel Dis* 2009; **15**:1526–36.
- 84 Hacini-Rachinel F, Nancey S, Boschetti G, Sardi F, Doucet-Ladeveze R, Durand P-Y *et al.* CD4 $^{+}$ T cells and *Lactobacillus casei* control relapsing colitis mediated by CD8 $^{+}$ T cells. *J Immunol* 2009; **183**:5477–86.
- 85 Dai C, Zheng C-Q, Meng F, Zhou Z, Sang L, Jiang M. VSL#3 probiotics exerts the anti-inflammatory activity via PI3k/Akt and NF- κ B pathway in rat model of DSS-induced colitis. *Mol Cell Biochem* 2013; **374**:1–11.
- 86 Salim SY, Young PY, Lukowski CM, Madsen KL, Sis B, Churchill TA *et al.* VSL#3 probiotics provide protection against acute intestinal ischaemia/reperfusion injury. *Benef Microbes* 2013; **4**:357–65.
- 87 Talero E, Bolivar S, Ávila-Román J, Alcaide A, Fiorucci S, Motilva V. Inhibition of chronic ulcerative colitis-associated adenocarcinoma development in mice by VSL#3. *Inflamm Bowel Dis* 2015; **21**:1027–37.
- 88 Schultz M, Veltkamp C, Dieleman LA, Grenther WB, Wyrick PB, Tonkonogy SL *et al.* *Lactobacillus plantarum* 299V in the treatment and prevention of spontaneous colitis in interleukin-10-deficient mice. *Inflamm Bowel Dis* 2002; **8**:71–80.
- 89 DuPont A, Richards DM, Jelinek KA, Krill J, Rahimi E, Ghouri Y. Systematic review of randomized controlled trials of probiotics, prebiotics, and synbiotics in inflammatory bowel disease. *Clin Exp Gastroenterol* 2014; **7**:473.
- 90 Whelan K, Quigley EMM. Probiotics in the management of irritable bowel syndrome and inflammatory bowel disease. *Curr Opin Gastroenterol* 2013; **29**:184–9.
- 91 Veerappan GR, Betteridge J, Young PE. Probiotics for the treatment of inflammatory bowel disease. *Curr Gastroenterol Rep* 2012; **14**:324–33.
- 92 Steidler L, Hans W, Schotte L, Neirynek S, Obermeier F, Falk W *et al.* Treatment of murine colitis by *Lactococcus lactis* secreting interleukin-10. *Science* 2000; **289**:1352–5.
- 93 Braat H, Rottiers P, Hommes DW, Huyghebaert N, Remaut E, Remon J *et al.* A phase I trial with transgenic bacteria expressing interleukin-10 in Crohn's disease. *Clin Gastroenterol Hepatol* 2006; **4**:754–9.
- 94 Vandenbroucke K, Hans W, Van Huysse J, Neirynek S, Demetter P, Remaut E *et al.* Active delivery of trefoil factors by genetically modified *Lactococcus lactis* prevents and heals acute colitis in mice. *Gastroenterology* 2004; **127**:502–13.
- 95 Maldonado-Gómez MX, Martínez I, Bottacini F, O'Callaghan A, Ventura M, van Sinderen D *et al.* Stable engraftment of *Bifidobacterium longum* AH1206 in the human gut depends on individualized features of the resident microbiome. *Cell Host Microbe* 2016; **20**:515–26.
- 96 Gibson GR, Hutkins R, Sanders ME, Prescott SL, Reimer RA, Salminen SJ *et al.* The international scientific association for probiotics and prebiotics (ISAPP) consensus statement on the definition and scope of prebiotics. *Nat Rev Gastroenterol Hepatol* 2017; **14**:491–502.
- 97 Dewulf EM, Cani PD, Claus SP, Fuentes S, Puylaert PGB, Neyrinck AM *et al.* Insight into the prebiotic concept: lessons from an exploratory, double blind intervention study with inulin-type fructans in obese women. *Gut* 2013; **62**:1112–21.
- 98 Costabile A, Kolida S, Klinder A, Gietl E, Bäuerlein M, Froberg C *et al.* A double-blind, placebo-controlled, cross-over study to establish the bifidogenic effect of a very-long-chain inulin extracted from globe artichoke (*Cynara scolymus*) in healthy human subjects. *Br J Nutr* 2010; **104**:1007–17.
- 99 Shoaf K, Mulvey GL, Armstrong GD, Hutkins RW. Prebiotic galactooligosaccharides reduce adherence of enteropathogenic *Escherichia coli* to tissue culture cells. *Infect Immun* 2006; **74**:6920–8.
- 100 Tarini J, Wolever TMS. The fermentable fibre inulin increases postprandial serum short-chain fatty acids and reduces free-fatty acids and ghrelin in healthy subjects. *Appl Physiol Nutr Metab* 2010; **35**:9–16.
- 101 Jung T-H, Jeon W-M, Han K-S. *In vitro* effects of dietary inulin on human fecal microbiota and butyrate production. *J Microbiol Biotechnol* 2015; **25**:1555–8.
- 102 MacPherson G, Milling S, Yrlid U, Cousins L, Turnbull E, Huang FP. Uptake of antigens from the intestine by dendritic cells. *Ann N Y Acad Sci* 2004; **1029**:75–82.
- 103 de Kivit S, Kraneveld AD, Garssen J, Willemsen LE. Glycan recognition at the interface of the intestinal immune system: target for immune modulation via dietary components. *Eur J Pharmacol* 2011; **668**:26–50.
- 104 Ortega-González M, Ocón B, Romero-Calvo I, Anzola A, Guadix E, Zarzuelo A *et al.* Nondigestible oligosaccharides exert nonprebiotic effects on intestinal epithelial cells enhancing the immune response via activation of TLR4-NF κ B. *Mol Nutr Food Res* 2014; **58**:384–93.
- 105 Osman N, Adawi D, Molin G, Ahrne S, Berggren A, Jeppsson B. *Bifidobacterium infantis* strains with and without a combination of oligofructose and inulin (OFI) attenuate inflammation in DSS-induced colitis in rats. *BMC Gastroenterol* 2006; **6**:31.
- 106 Winkler J, Butler R, Symonds E. Fructo-oligosaccharide reduces inflammation in a dextran sodium sulphate mouse model of colitis. *Dig Dis Sci* 2007; **52**:52–8.
- 107 Hoentjen F, Welling GW, Harmsen HJM, Zhang X, Snart J, Tannock GW *et al.* Reduction of colitis by prebiotics in HLA-B27 transgenic rats is associated with microflora changes and immunomodulation. *Inflamm Bowel Dis* 2005; **11**:977–85.
- 108 Koleva PT, Valcheva RS, Sun X, Gänzle MG, Dieleman LA. Inulin and fructo-oligosaccharides have divergent effects on colitis and commensal microbiota in HLA-B27 transgenic rats. *Br J Nutr* 2012; **108**:1633–43.
- 109 Schultz M, Munro K, Tannock GW, Melchner I, Göttl C, Schwietz H *et al.* Effects of feeding a probiotic preparation (SIM) containing inulin on the severity of colitis and on the composition of the intestinal microflora in HLA-B27 transgenic rats. *Clin Diagn Lab Immunol* 2004; **11**:581–7.
- 110 Cherbut C, Michel C, Lecannu G. The prebiotic characteristics of fructooligosaccharides are necessary for reduction of TNBS-induced colitis in rats. *J Nutr* 2003; **133**:21–7.
- 111 Ito H, Tanabe H, Kawagishi H, Wada T, Tomono Y, Sugiyama K *et al.* Short-chain inulin-type fructans reduce endotoxin and bacterial translocations and attenuate development of TNBS-induced colitis in rats. *Dig Dis Sci* 2009; **54**:2100–8.
- 112 Videla S, Vilaseca J, Antolin M, Garcia-Lafuente A, Guarner F, Crespo E *et al.* Dietary inulin improves distal colitis induced by dextran sodium sulfate in the rat. *Am J Gastroenterol* 2001; **96**:1486–93.
- 113 Moreau NM, Martin LJ, Toquet CS, Labois CL, Nguyen PG, Siliart BS *et al.* Restoration of the integrity of rat caeco-colonic mucosa by resistant starch, but not by fructo-oligosaccharides, in dextran sulfate sodium-induced experimental colitis. *Br J Nutr* 2003; **90**:75–85.
- 114 Lindsay JO, Whelan K, Stagg AJ, Gobin P, Al-Hassi HO, Rayment N *et al.* Clinical, microbiological, and immunological effects of fructo-oligosaccharide in patients with Crohn's disease. *Gut* 2006; **55**:348–56.
- 115 Steed H, MacFarlane GT, Blackett KL, Bahrami B, Reynolds N, Walsh SV *et al.* Clinical trial: the microbiological and immunological effects of synbiotic consumption – a randomized double-blind placebo-controlled study in active Crohn's disease. *Aliment Pharmacol Ther* 2010; **32**:872–83.
- 116 Joossens M, De Preter V, Ballet V, Verbeke K, Rutgeerts P, Vermeire S. Effect of oligofructose-enriched inulin (OF-IN) on bacterial composition and disease activity of patients with Crohn's disease: results from a double-blinded randomised controlled trial. *Gut* 2011; **61**:958.
- 117 Furrie E, Macfarlane S, Kennedy A, Cummings JH, Walsh SV, O'neil DA *et al.* Synbiotic therapy (*Bifidobacterium longum*/Synergy 1) initiates resolution of inflammation in patients with active ulcerative colitis: a randomised controlled pilot trial. *Gut* 2005; **54**:242–9.
- 118 Federico A, Tuccillo C, Grossi E, Abbiati R, Garbagna N, Romano M *et al.* The effect of a new symbiotic formulation on plasma levels and peripheral blood mononuclear cell expression of some pro-inflammatory cytokines in patients with ulcerative colitis: a pilot study. *Eur Rev Med Pharmacol Sci* 2009; **13**:285–93.
- 119 Casellas F, Borruel N, Torrejon A, Varela E, Antolin M, Guarner F *et al.* Oral oligofructose-enriched inulin supplementation in acute ulcerative colitis is well tolerated and associated with lowered faecal calprotectin. *Aliment Pharmacol Ther* 2007; **25**:1061–7.

- 120 Ishikawa H, Matsumoto S, Ohashi Y, Imaoka A, Setoyama H, Umesaki Y *et al.* Beneficial effects of probiotic *Bifidobacterium* and galacto-oligosaccharide in patients with ulcerative colitis: a randomized controlled study. *Digestion* 2011; **84**:128–33.
- 121 Benjamin JL, Hedin CRH, Koutsoumpas A, Ng SC, McCarthy NE, Hart AL *et al.* Randomised, double-blind, placebo-controlled trial of fructo-oligosaccharides in active Crohn's disease. *Gut* 2011; **60**:923–9.
- 122 Hafer A, Krämer S, Duncker S, Krüger M, Manns MP, Bischoff SC. Effect of oral lactulose on clinical and immunohistochemical parameters in patients with inflammatory bowel disease: a pilot study. *BMC Gastroenterol* 2007; **7**:1–11.
- 123 White JH. Vitamin D deficiency and the pathogenesis of Crohn's disease. *J Steroid Biochem Mol Biol Pergamon* 2018; **175**:23–8.
- 124 Jin D, Wu S, Zhang Y, Lu R, Xia Y, Dong H *et al.* Lack of vitamin D receptor causes dysbiosis and changes the functions of the murine intestinal microbiome. *Clin Ther* 2015; **37**:996–1009. e7.
- 125 Sokol H, Seksik P, Rigottier-Gois L, Lay C, Lepage P, Podglajen I *et al.* Specificities of the fecal microbiota in inflammatory bowel disease. *Inflamm Bowel Dis* 2006; **12**:106–11.
- 126 Ooi JH, Li Y, Rogers CJ, Cantorna MT. Vitamin D regulates the gut microbiome and protects mice from dextran sodium sulfate-induced colitis. *J Nutr* 2013; **143**:1679–86.
- 127 Ryz NR, Patterson SJ, Zhang Y, Ma C, Huang T, Bhinder G *et al.* Active vitamin D (1,25-dihydroxyvitamin D₃) increases host susceptibility to *Citrobacter rodentium* by suppressing mucosal Th17 responses. *Am J Physiol Liver Physiol* 2012; **303**:G1299–311.
- 128 Ryz NR, Lochner A, Bhullar K, Ma C, Huang T, Bhinder G *et al.* Dietary vitamin D3 deficiency alters intestinal mucosal defense and increases susceptibility to *Citrobacter rodentium*-induced colitis. *Am J Physiol Liver Physiol* 2015; **309**:G730–42.
- 129 Wu S, Liao AP, Xia Y, Chun Li Y, Li J-D, Sartor RB *et al.* Vitamin D receptor negatively regulates bacterial-stimulated NF- κ B activity in intestine. *Am J Pathol* 2010; **177**:686–97.
- 130 Takada I, Makishima M. Control of inflammatory bowel disease and colorectal cancer by synthetic vitamin D receptor ligands. *Curr Med Chem* 2017; **24**:868–75.
- 131 Gaschott T, Werz O, Steinmeyer A, Steinhilber D, Stein J. Butyrate-induced differentiation of caco-2 cells is mediated by vitamin D receptor. *Biochem Biophys Res Commun* 2001; **288**:690–6.
- 132 Schwab M, Reynders V, Loitsch S, Steinhilber D, Stein J, Schröder O. Involvement of different nuclear hormone receptors in butyrate-mediated inhibition of inducible NF κ B signalling. *Mol Immunol* 2007; **44**:3625–32.
- 133 Wu S, Yoon S, Zhang Y-G, Lu R, Xia Y, Wan J *et al.* Vitamin D receptor pathway is required for probiotic protection in colitis. *Am J Physiol Liver Physiol* 2015; **309**:G341–9.
- 134 Alhassan Mohammed H, Mirshafiey A, Vahedi H, Hemmasi G, Moussavi Nasl Khameneh A, Parastouei K *et al.* Immunoregulation of inflammatory and inhibitory cytokines by vitamin D3 in patients with inflammatory bowel diseases. *Scand J Immunol* 2017; **85**:386–94.
- 135 Chen Y, Liu W, Sun T, Huang Y, Wang Y, Deb DK *et al.* 1,25-dihydroxyvitamin D promotes negative feedback regulation of TLR signaling via targeting MicroRNA-155-SOCS1 in macrophages. *J Immunol* 2013; **190**:3687–95.
- 136 Ferreira GB, Vanherwegen A-S, Eelen G, Gutiérrez ACF, Van Lommel L, Marchal K *et al.* Vitamin D3 induces tolerance in human dendritic cells by activation of intracellular metabolic pathways. *Cell Rep* 2015; **10**:711–25.
- 137 Winter RW, Collins E, Cao B, Carrellas M, Crowell AM, Korzenik JR. Higher 25-hydroxyvitamin D levels are associated with greater odds of remission with anti-tumour necrosis factor- α medications among patients with inflammatory bowel diseases. *Aliment Pharmacol Ther* 2017; **45**:653–9.
- 138 Santos-Antunes J, Nunes AC-R, Lopes S, Macedo G. The relevance of vitamin D and antinuclear antibodies in patients with inflammatory bowel disease under anti-TNF treatment. *Inflamm Bowel Dis* 2016; **22**:1101–6.
- 139 Ananthakrishnan AN, Cagan A, Gainer VS, Cai T, Cheng S-C, Savova G *et al.* Normalization of plasma 25-hydroxy vitamin D is associated with reduced risk of surgery in Crohn's disease. *Inflamm Bowel Dis* 2013; **19**:1.
- 140 Barbalho SM, Goulart RdeA, Gasparini RG. Associations between inflammatory bowel diseases and vitamin D. *Crit Rev Food Sci Nutr* 2017; **00**:1–10.
- 141 Shah A, Mehta N, Reilly MP. Adipose inflammation, insulin resistance, and cardiovascular disease. *JPEN J Parenter Enteral Nutr* 2008; **32**:638–44.
- 142 DeBerardinis RJ, Thompson CB. Cellular metabolism and disease: what do metabolic outliers teach us? *Cell* 2012; **148**:1132–44.
- 143 Osborn O, Olefsky JM. The cellular and signaling networks linking the immune system and metabolism in disease. *Nat Med* 2012; **18**:363–74.
- 144 Buck MD, Sowell RT, Kaech SM, Pearce EL. Metabolic instruction of immunity. *Cell* 2017; **169**:570–86.
- 145 O'Neill LAJ, Kishton RJ, Rathmell J. A guide to immunometabolism for immunologists. *Nat Rev Immunol* 2016; **16**:553–65.
- 146 De Rosa V, La Cava A, Matarese G. Metabolic pressure and the breach of immunological self-tolerance. *Nat Immunol* 2017; **18**:1190–6.
- 147 Eastaff-Leung N, Mabarrack N, Barbour A, Cummins A, Barry S. Foxp3⁺ regulatory T cells, Th17 effector cells, and cytokine environment in inflammatory bowel disease. *J Clin Immunol* 2010; **30**:80–9.
- 148 Ben Sahra I, Regazzetti C, Robert G, Laurent K, Le Marchand-Brustel Y, Auberger P *et al.* Metformin, independent of AMPK, induces mTOR inhibition and cell-cycle arrest through REDD1. *Cancer Res* 2011; **71**:4366–72.
- 149 Hu S, Chen M, Wang Y, Wang Z, Pei Y, Fan R *et al.* mTOR inhibition attenuates dextran sulfate sodium-induced colitis by suppressing T cell proliferation and balancing TH1/TH17/Treg profile. *PLoS ONE* 2016; **11**:e0154564.
- 150 Mutalib M, Borrelli O, Blackstock S, Kiparissi F, Elawad M, Shah N *et al.* The use of sirolimus (rapamycin) in the management of refractory inflammatory bowel disease in children. *J Crohn's Colitis* 2014; **8**:1730–4.
- 151 Pallet N, Legendre C. Adverse events associated with mTOR inhibitors. *Expert Opin Drug Saf* 2013; **12**:177–86.
- 152 Ma L, Dong W, Wang R, Li Y, Xu B, Zhang J *et al.* Effect of caloric restriction on the SIRT1/mTOR signaling pathways in senile mice. *Brain Res Bull* 2015; **116**:67–72.
- 153 Owen OE, Reichard GA, Patel MS, Boden G. Energy metabolism in feasting and fasting. *Adv Exp Med Biol* 1979; **111**:169–88.
- 154 Choi IY, Piccio L, Childress P, Bollman B, Ghosh A, Brandhorst S *et al.* A diet mimicking fasting promotes regeneration and reduces autoimmunity and multiple sclerosis symptoms. *Cell Rep* 2016; **15**:2136–46.
- 155 Ip WKE, Hoshi N, Shouval DS, Snapper S, Medzhitov R. Anti-inflammatory effect of IL-10 mediated by metabolic reprogramming of macrophages. *Science* 2017; **356**:513 LP-519.
- 156 Schupp M, Chen F, Briggs ER, Rao S, Pelzmann HJ, Pessentheiner AR *et al.* Metabolite and transcriptome analysis during fasting suggest a role for the p53-Ddit4 axis in major metabolic tissues. *BMC Genom* 2013; **14**:758.
- 157 Youm Y-H, Nguyen KY, Grant RW, Goldberg EL, Bodogai M, Kim D *et al.* The ketone metabolite β -hydroxybutyrate blocks NLRP3 inflammasome-mediated inflammatory disease. *Nat Med* 2015; **21**:263–9.
- 158 Traba J, Geiger SS, Kwarteng Siaw M, Han K, Ra OH, Siegel RM *et al.* Prolonged fasting suppresses mitochondrial NLRP3 inflammasome assembly and activation via SIRT3 mediated activation of superoxide dismutase 2. *J Biol Chem* 2017; **292**:12153–64.
- 159 Shimazu T, Hirschey MD, Newman J, He W, Shirakawa K, Le Moan N *et al.* Suppression of oxidative stress by β -hydroxybutyrate, an endogenous histone deacetylase inhibitor. *Science* 2013; **339**:211–4.
- 160 Rojas-Morales P, Tapia E, Pedraza-Chaverri J. β -Hydroxybutyrate: a signaling metabolite in starvation response? *Cell Signal* 2016; **28**:917–23.
- 161 Iida T, Onodera K, Nakase H. Role of autophagy in the pathogenesis of inflammatory bowel disease. *World J Gastroenterol* 2017; **23**:1944–53.
- 162 Remely M, Hippe B, Geretschlaeger I, Stegmayer S, Hoefinger I, Haslberger A. Increased gut microbiota diversity and abundance of *Faecalibacterium prausnitzii* and *Akkermansia* after fasting: a pilot study. *Wien Klin Wochenschr* 2015; **127**:394–8.
- 163 Hartog A, Belle FN, Bastiaans J, De Graaff P, Garssen J, Harthoorn LF *et al.* A potential role for regulatory T-cells in the amelioration of DSS induced colitis by dietary non-digestible polysaccharides. *J Nutr Biochem* 2015; **26**:227–33.
- 164 Heinsbroek SEM, Williams DL, Welting O, Meijer SL, Gordon S, de Jonge WJ. Orally delivered β -glucans aggravate dextran sulfate sodium (DSS)-induced intestinal inflammation. *Nutr Res* 2015; **35**:1106–12.
- 165 Liu B, Lin Q, Yang T, Zeng L, Shi L, Chen Y *et al.* Oat β -glucan ameliorates dextran sulfate sodium (DSS)-induced ulcerative colitis in mice. *Food Funct* 2015; **6**:3454–63.
- 166 Han F, Fan H, Yao M, Yang S, Han J. Oral administration of yeast β -glucan ameliorates inflammation and intestinal barrier in dextran sodium sulfate-induced acute colitis. *J Funct Foods* 2017; **35**:115–26.
- 167 Valcheva R, Hotte N, Gillevet P, Sikaroodi M, Thiessen A, Madsen KL. Soluble dextrin fibers alter the intestinal microbiota and reduce proinflammatory cytokine expression in male IL-10-deficient mice. *J Nutr* 2015; **145**:2060–6.
- 168 Holma R, Juvonen P, Asmawi MZ, Vapaatalo H, Korpela R. Galacto-oligosaccharides stimulate the growth of bifidobacteria but fail to attenuate inflammation in experimental colitis in rats. *Scand J Gastroenterol* 2002; **37**:1042–7.
- 169 Bourreille A, Cadiot G, Le Dreau G, Laharie D, Beaugerie L, Dupas J *et al.* *Saccharomyces boulardii* does not prevent relapse of Crohn's disease. *Clin Gastroenterol Hepatol* 2013; **11**:982–7.
- 170 Van Gossom A, Dewit O, Louis E, de Hertogh G, Baert F, Fontaine F *et al.* Multicenter randomized-controlled clinical trial of probiotics (*Lactobacillus johnsonii*, LA1) on early endoscopic recurrence of Crohn's disease after ileo-caecal resection. *Inflamm Bowel Dis* 2007; **13**:135–42.
- 171 Ng SC, Plamondon S, Kamm MA, Hart AL, Al-Hassi HO, Guenther T *et al.* Immunosuppressive effects via human intestinal dendritic cells of probiotic bacteria and steroids in the treatment of acute ulcerative colitis. *Inflamm Bowel Dis* 2010; **16**:1286–98.
- 172 Cui H-H, Chen C-L, Wang J-D, Yang Y-J, Cun Y, Wu J-B *et al.* Effects of probiotic on intestinal mucosa of patients with ulcerative colitis. *World J Gastroenterol* 2004; **10**:1521–5.