

An abstract network diagram with nodes and edges, rendered in blue, red, and yellow colors, set against a dark blue background with a bokeh effect.

COMP SCI 2201-7201

Algorithm and Data Structure Analysis

Asymptotic Analysis

Dr. Muhammad Uzair

School of Computer and Mathematical Sciences

**make
history.**



Asymptotic Analysis

- *Asymptotic analysis* is a branch of mathematics that studies the behavior of mathematical functions as their input become very large



Asymptotic Analysis

- *Asymptotic analysis* is a branch of mathematics that studies the behavior of mathematical functions as their input become very large
 - Functions are not studied alone but are studied relative to other functions

Asymptotic Analysis

- *Asymptotic analysis* is a branch of mathematics that studies the behavior of mathematical functions as their input become very large
 - Functions are not studied alone but are studied relative to other functions
 - Usually, the *growth rate of functions relative to other functions* is studied (in computer science)



Asymptotic Analysis

- *Asymptotic analysis* is a branch of mathematics that studies the behavior of mathematical functions as their input become very large
 - Functions are not studied alone but are studied relative to other functions
 - Usually, the *growth rate of functions relative to other functions* is studied (in computer science)
- *Asymptotic analysis* or *Asymptotics* is the calculus of approximations
 - Approximation of functions by simpler functions



Asymptotic Analysis

- *Asymptotic analysis* is a branch of mathematics that studies the behavior of mathematical functions as their input become very large
 - Functions are not studied alone but are studied relative to other functions
 - Usually, the *growth rate of functions relative to other functions* is studied (in computer science)
- *Asymptotic analysis* or *Asymptotics* is the calculus of approximations
 - Approximation of functions by simpler functions
- The term "asymptotic" can also be used more broadly to describe situations where two things approach each other or become more similar, but never quite reach a point of perfect equality.

Example

$$f(n) = n^2 + 4n + 7$$

$$g(n) = n^2$$



Example

$$f(n) = n^2 + 4n + 7 \qquad g(n) = n^2$$

- All the below statements mean the same

$f(n)$ is "asymptotically similar" to $g(n)$ as $n \rightarrow \infty$

$f(n)$ is "asymptotically same" to $g(n)$ as $n \rightarrow \infty$

$f(n)$ is "asymptotically equivalent" to $g(n)$ as $n \rightarrow \infty$

$f(n)$ is "asymptotically equal" to $g(n)$ as $n \rightarrow \infty$

$f(n)$ is "asymptotic" to $g(n)$ as $n \rightarrow \infty$

$$f(n) \sim n^2$$



Example

$$f(n) = n^2 + 4n + 7 \qquad g(n) = n^2$$

- All the below statements mean the same

$f(n)$ is "asymptotically similar" to $g(n)$ as $n \rightarrow \infty$

$f(n)$ is "asymptotically same" to $g(n)$ as $n \rightarrow \infty$

$f(n)$ is "asymptotically equivalent" to $g(n)$ as $n \rightarrow \infty$

$f(n)$ is "asymptotically equal" to $g(n)$ as $n \rightarrow \infty$

$f(n)$ is "asymptotic" to $g(n)$ as $n \rightarrow \infty$

$$f(n) \sim n^2$$

\sim is an asymptotic notation used to convey the message that a function is asymptotically similar/same/equivalent/equal to another function

Asymptotic Notations

- A way to communicate the relationship between the behaviors of different functions
 - Usually, the relationship between the *growth rates* of different functions (in computer science)



Asymptotic Notations (Relation formulas)

重要

$f(n) = O(g(n))$ $\exists c > 0, \exists n_0 > 0$ such that $\forall n \geq n_0$ $|f(n)| \leq cg(n)$ Asymptotic upper bound

$f(n) = \Omega(g(n))$ $\exists c > 0, \exists n_0 > 0$ such that $\forall n \geq n_0$ $|f(n)| \geq cg(n)$ Asymptotic lower bound

$f(n) = \Theta(g(n))$ $\exists c_1, c_2 > 0, \exists n_0 > 0$ such that $\forall n \geq n_0$ $c_1g(n) \leq |f(n)| \leq c_2g(n)$ Asymptotic tight bound

$f(n) = o(g(n))$ $\forall c > 0, \exists n_0 > 0$ such that $\forall n \geq n_0$ $|f(n)| \leq cg(n)$
upper bound that is not asymptotically tight

$f(n) = \omega(g(n))$ $\forall c > 0, \exists n_0 > 0$ such that $\forall n \geq n_0$ $|f(n)| \geq cg(n)$
lower bound that is not asymptotically tight

$f(n) \sim g(n)$ $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 1$

$f(n) = O(g(n))$ can also be written as $f(n) \in O(g(n))$



Asymptotic Notations (Limits)

$$f(n) = O(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq \infty$$

$$f(n) = \Omega(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq 0$$

$$f(n) = \Theta(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq 0, \infty$$

$$f(n) = o(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0$$

$$f(n) = \omega(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \infty$$

$$f(n) \sim g(n) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 1$$

How to perform asymptotic analysis

- Identify the dominant term $f(n) = 3n^3 + 2n^2 + 5n + 7$



How to perform asymptotic analysis

- Identify the dominant term $f(n) = \underline{3n^3} + 2n^2 + 5n + 7$



How to perform asymptotic analysis

- Identify the dominant term $f(n) = \underline{3n^3} + 2n^2 + 5n + 7$

- Drop the lower order terms $3n^3$



How to perform asymptotic analysis

- Identify the dominant term $f(n) = \underline{3n^3} + 2n^2 + 5n + 7$

- Drop the lower order terms $3n^3$

- Drop the constant factors n^3

How to perform asymptotic analysis

- Identify the dominant term $f(n) = \underline{3n^3} + 2n^2 + 5n + 7$
- Drop the lower order terms $3n^3$
- Drop the constant factors n^3 Let's call it $g(n) = n^3$



How to perform asymptotic analysis

- Identify the dominant term $f(n) = \underline{3n^3} + 2n^2 + 5n + 7$
- Drop the lower order terms $3n^3$
- Drop the constant factors n^3 Let's call it $g(n) = n^3$
- Identify the asymptotic notations

Analyze $f(n)$ and $g(n)$ using relation or limits formulas

How to perform asymptotic analysis

1. Using relation formulas

$$f(n) = 3n^3 + 2n^2 + 5n + 7 \quad g(n) = n^3$$

$$f(n) = O(g(n)) \quad \exists c > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad |f(n)| \leq cg(n)$$

$$f(n) = \Omega(g(n)) \quad \exists c > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad |f(n)| \geq cg(n)$$

$$f(n) = \Theta(g(n)) \quad \exists c_1, c_2 > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad c_1 g(n) \leq |f(n)| \leq c_2 g(n)$$

$$f(n) = o(g(n)) \quad \forall c > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad |f(n)| \leq cg(n)$$

$$f(n) = \omega(g(n)) \quad \forall c > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad |f(n)| \geq cg(n)$$



How to perform asymptotic analysis

1. Using relation formulas

$$f(n) = 3n^3 + 2n^2 + 5n + 7 \quad g(n) = n^3$$

$$|f(n)| \leq cg(n) \quad \text{for a } c > 0 \text{ and } n_0 > 0$$

$$f(n) = O(g(n)) \quad \exists c > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad |f(n)| \leq cg(n)$$

$$f(n) = \Omega(g(n)) \quad \exists c > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad |f(n)| \geq cg(n)$$

$$f(n) = \Theta(g(n)) \quad \exists c_1, c_2 > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad c_1g(n) \leq |f(n)| \leq c_2g(n)$$

$$f(n) = o(g(n)) \quad \forall c > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad |f(n)| \leq cg(n)$$

$$f(n) = \omega(g(n)) \quad \forall c > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad |f(n)| \geq cg(n)$$



How to perform asymptotic analysis

1. Using relation formulas

$$f(n) = 3n^3 + 2n^2 + 5n + 7 \quad g(n) = n^3$$

$$|f(n)| \leq cg(n) \quad \text{for a } c > 0 \text{ and } n_0 > 0$$

$$3n^3 + 2n^2 + 5n + 7 \leq 17n^3, \quad c = 17, n_0 = 1$$

$$f(n) = O(g(n)) \quad \exists c > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad |f(n)| \leq cg(n)$$

$$f(n) = \Omega(g(n)) \quad \exists c > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad |f(n)| \geq cg(n)$$

$$f(n) = \Theta(g(n)) \quad \exists c_1, c_2 > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad c_1g(n) \leq |f(n)| \leq c_2g(n)$$

$$f(n) = o(g(n)) \quad \forall c > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad |f(n)| \leq cg(n)$$

$$f(n) = \omega(g(n)) \quad \forall c > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad |f(n)| \geq cg(n)$$



How to perform asymptotic analysis

1. Using relation formulas

$$f(n) = 3n^3 + 2n^2 + 5n + 7 \quad g(n) = n^3$$

$$|f(n)| \leq cg(n) \quad \text{for a } c > 0 \text{ and } n_0 > 0$$

$$3n^3 + 2n^2 + 5n + 7 \leq 17n^3, \quad c = 17, n_0 = 1$$

$$f(n) = O(n^3) \quad \text{YES}$$

$$f(n) = O(g(n)) \quad \exists c > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad |f(n)| \leq cg(n)$$

$$f(n) = \Omega(g(n)) \quad \exists c > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad |f(n)| \geq cg(n)$$

$$f(n) = \Theta(g(n)) \quad \exists c_1, c_2 > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad c_1g(n) \leq |f(n)| \leq c_2g(n)$$

$$f(n) = o(g(n)) \quad \forall c > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad |f(n)| \leq cg(n)$$

$$f(n) = \omega(g(n)) \quad \forall c > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad |f(n)| \geq cg(n)$$



How to perform asymptotic analysis

1. Using relation formulas

$$f(n) = 3n^3 + 2n^2 + 5n + 7 \quad g(n) = n^3$$

$$|f(n)| \leq cg(n) \quad \text{for a } c > 0 \text{ and } n_0 > 0$$

$$3n^3 + 2n^2 + 5n + 7 \leq 17n^3, \quad c = 17, n_0 = 1$$

$$f(n) = O(n^3) \quad \text{YES}$$

$$|f(n)| \geq cg(n) \quad \text{for a } c > 0 \text{ and } n_0 > 0$$

$$3n^3 + 2n^2 + 5n + 7 \geq 3n^3, \quad c = 3, n_0 = 1$$

$$f(n) = \Omega(n^3) \quad \text{YES}$$

$$f(n) = O(g(n)) \quad \exists c > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad |f(n)| \leq cg(n)$$

$$f(n) = \Omega(g(n)) \quad \exists c > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad |f(n)| \geq cg(n)$$

$$f(n) = \Theta(g(n)) \quad \exists c_1, c_2 > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad c_1 g(n) \leq |f(n)| \leq c_2 g(n)$$

$$f(n) = o(g(n)) \quad \forall c > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad |f(n)| \leq cg(n)$$

$$f(n) = \omega(g(n)) \quad \forall c > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad |f(n)| \geq cg(n)$$



How to perform asymptotic analysis

1. Using relation formulas

$$f(n) = 3n^3 + 2n^2 + 5n + 7 \quad g(n) = n^3$$

$$|f(n)| \leq cg(n) \quad \text{for a } c > 0 \text{ and } n_0 > 0$$

$$3n^3 + 2n^2 + 5n + 7 \leq 17n^3, \quad c = 17, n_0 = 1$$

$$f(n) = O(n^3) \quad \text{YES}$$

$$|f(n)| \geq cg(n) \quad \text{for a } c > 0 \text{ and } n_0 > 0$$

$$3n^3 + 2n^2 + 5n + 7 \geq 3n^3, \quad c = 3, n_0 = 1$$

$$f(n) = \Omega(n^3) \quad \text{YES}$$

We have already found two c's above $c_1 = 3, c_2 = 17$

$$3n^3 \leq f(n) \leq 17n^3$$

$$f(n) = \Theta(n^3) \quad \text{YES}$$

$$f(n) = O(g(n)) \quad \exists c > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad |f(n)| \leq cg(n)$$

$$f(n) = \Omega(g(n)) \quad \exists c > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad |f(n)| \geq cg(n)$$

$$f(n) = \Theta(g(n)) \quad \exists c_1, c_2 > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad c_1 g(n) \leq |f(n)| \leq c_2 g(n)$$

$$f(n) = o(g(n)) \quad \forall c > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad |f(n)| \leq cg(n)$$

$$f(n) = \omega(g(n)) \quad \forall c > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad |f(n)| \geq cg(n)$$



How to perform asymptotic analysis

1. Using relation formulas

$$f(n) = 3n^3 + 2n^2 + 5n + 7 \quad g(n) = n^3$$

$$|f(n)| \leq cg(n) \quad \text{for a } c > 0 \text{ and } n_0 > 0$$

$$3n^3 + 2n^2 + 5n + 7 \leq 17n^3, \quad c = 17, n_0 = 1$$

$$f(n) = O(n^3) \quad \text{YES}$$

$$|f(n)| \geq cg(n) \quad \text{for a } c > 0 \text{ and } n_0 > 0$$

$$3n^3 + 2n^2 + 5n + 7 \geq 3n^3, \quad c = 3, n_0 = 1$$

$$f(n) = \Omega(n^3) \quad \text{YES}$$

We have already found two c 's above $c_1 = 3, c_2 = 17$

$$3n^3 \leq f(n) \leq 17n^3$$

$$f(n) = \Theta(n^3) \quad \text{YES}$$

$$f(n) = O(g(n)) \quad \exists c > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad |f(n)| \leq cg(n)$$

$$f(n) = \Omega(g(n)) \quad \exists c > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad |f(n)| \geq cg(n)$$

$$f(n) = \Theta(g(n)) \quad \exists c_1, c_2 > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad c_1 g(n) \leq |f(n)| \leq c_2 g(n)$$

$$f(n) = o(g(n)) \quad \forall c > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad |f(n)| \leq cg(n)$$

$$f(n) = \omega(g(n)) \quad \forall c > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad |f(n)| \geq cg(n)$$

$$|f(n)| \leq cg(n) \quad \forall c > 0$$

$$3n^3 + 2n^2 + 5n + 7 \leq cn^3, \text{ not for all } c > 0$$

$$f(n) = o(n^3) \quad \text{NO}$$

$$|f(n)| \geq cg(n) \quad \forall c > 0$$

$$3n^3 + 2n^2 + 5n + 7 \geq cn^3, \text{ not for all } c > 0$$

$$f(n) = \omega(n^3) \quad \text{NO}$$



How to perform asymptotic analysis

1. Using relation formulas

$$f(n) = 3n^3 + 2n^2 + 5n + 7 \quad g(n) = n^3$$

$$|f(n)| \leq cg(n) \quad \text{for a } c > 0 \text{ and } n_0 > 0$$

$$3n^3 + 2n^2 + 5n + 7 \leq 17n^3, \quad c = 17, n_0 = 1$$

$$f(n) = O(n^3) \quad \text{YES}$$

$$|f(n)| \geq cg(n) \quad \text{for a } c > 0 \text{ and } n_0 > 0$$

$$3n^3 + 2n^2 + 5n + 7 \geq 3n^3, \quad c = 3, n_0 = 1$$

$$f(n) = \Omega(n^3) \quad \text{YES}$$

We have already found two c 's above $c_1 = 3, c_2 = 17$

$$3n^3 \leq f(n) \leq 17n^3$$

$$f(n) = \Theta(n^3) \quad \text{YES}$$

$$f(n) = O(g(n)) \quad \exists c > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad |f(n)| \leq cg(n)$$

$$f(n) = \Omega(g(n)) \quad \exists c > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad |f(n)| \geq cg(n)$$

$$f(n) = \Theta(g(n)) \quad \exists c_1, c_2 > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad c_1 g(n) \leq |f(n)| \leq c_2 g(n)$$

$$f(n) = o(g(n)) \quad \forall c > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad |f(n)| \leq cg(n)$$

$$f(n) = \omega(g(n)) \quad \forall c > 0, \exists n_0 > 0 \text{ such that } \forall n \geq n_0 \quad |f(n)| \geq cg(n)$$

$$|f(n)| \leq cg(n) \quad \forall c > 0$$

$$3n^3 + 2n^2 + 5n + 7 \leq cn^3, \text{ not for all } c > 0$$

$$f(n) = o(n^3) \quad \text{NO}$$

$$|f(n)| \geq cg(n) \quad \forall c > 0$$

$$3n^3 + 2n^2 + 5n + 7 \geq cn^3, \text{ not for all } c > 0$$

$$f(n) = \omega(n^3) \quad \text{NO}$$

If a function $f(n)$ is both $O(g(n))$ and $\Omega(g(n))$, then $f(n)$ is also $\Theta(g(n))$



How to perform asymptotic analysis

1. Using limits formulas

$$f(n) = 3n^3 + 2n^2 + 5n + 7$$

$$g(n) = n^3$$

$$f(n) = O(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq \infty$$

$$f(n) = \Omega(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq 0$$

$$f(n) = \Theta(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq 0, \infty$$

$$f(n) = o(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0$$

$$f(n) = \omega(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \infty$$

$$f(n) \sim g(n) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 1$$



How to perform asymptotic analysis

1. Using limits formulas

$$f(n) = 3n^3 + 2n^2 + 5n + 7$$

$$g(n) = n^3$$

$$\lim_{n \rightarrow \infty} \frac{3n^3 + 2n^2 + 5n + 7}{n^3}$$

$$f(n) = O(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq \infty$$

$$f(n) = \Omega(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq 0$$

$$f(n) = \Theta(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq 0, \infty$$

$$f(n) = o(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0$$

$$f(n) = \omega(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \infty$$

$$f(n) \sim g(n) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 1$$



How to perform asymptotic analysis

1. Using limits formulas

$$f(n) = 3n^3 + 2n^2 + 5n + 7$$

$$g(n) = n^3$$

$$\lim_{n \rightarrow \infty} \frac{3n^3 + 2n^2 + 5n + 7}{n^3} = \frac{\infty}{\infty} \quad \text{indeterminate form}$$

$$f(n) = O(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq \infty$$

$$f(n) = \Omega(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq 0$$

$$f(n) = \Theta(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq 0, \infty$$

$$f(n) = o(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0$$

$$f(n) = \omega(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \infty$$

$$f(n) \sim g(n) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 1$$



How to perform asymptotic analysis

1. Using limits formulas

$$f(n) = 3n^3 + 2n^2 + 5n + 7$$

$$g(n) = n^3$$

$$\lim_{n \rightarrow \infty} \frac{3n^3 + 2n^2 + 5n + 7}{n^3} = \frac{\infty}{\infty} \quad \text{indeterminate form}$$

To get out of indeterminate situation, take the first derivate and then apply the limits

Keep taking the derivative until you get out of the indeterminate situation

$$f(n) = O(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq \infty$$

$$f(n) = \Omega(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq 0$$

$$f(n) = \Theta(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq 0, \infty$$

$$f(n) = o(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0$$

$$f(n) = \omega(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \infty$$

$$f(n) \sim g(n) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 1$$



How to perform asymptotic analysis

1. Using limits formulas

$$f(n) = 3n^3 + 2n^2 + 5n + 7$$

$$g(n) = n^3$$

$$\lim_{n \rightarrow \infty} \frac{3n^3 + 2n^2 + 5n + 7}{n^3} = \frac{\infty}{\infty} \quad \text{indeterminate form}$$

To get out of indeterminate situation, take
the first derivate and then apply the limits
Keep taking the derivative until
you get out of the indeterminate situation

L'Hôpital's rule for $\frac{\infty}{\infty}$ and $\frac{0}{0}$ indeterminate forms

$$f(n) = O(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq \infty$$

$$f(n) = \Omega(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq 0$$

$$f(n) = \Theta(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq 0, \infty$$

$$f(n) = o(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0$$

$$f(n) = \omega(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \infty$$

$$f(n) \sim g(n) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 1$$



How to perform asymptotic analysis

1. Using limits formulas

$$f(n) = 3n^3 + 2n^2 + 5n + 7$$

$$g(n) = n^3$$

$$\lim_{n \rightarrow \infty} \frac{3n^3 + 2n^2 + 5n + 7}{n^3} = \frac{\infty}{\infty} \quad \text{indeterminate form}$$

To get out of indeterminate situation, take
the first derivate and then apply the limits
Keep taking the derivative until
you get out of the indeterminate situation

L'Hôpital's rule for $\frac{\infty}{\infty}$ and $\frac{0}{0}$ indeterminate forms

$$\lim_{n \rightarrow \infty} \frac{9n^2 + 4n + 5}{3n^2}$$

$$f(n) = O(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq \infty$$

$$f(n) = \Omega(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq 0$$

$$f(n) = \Theta(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq 0, \infty$$

$$f(n) = o(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0$$

$$f(n) = \omega(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \infty$$

$$f(n) \sim g(n) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 1$$



How to perform asymptotic analysis

1. Using limits formulas

$$f(n) = 3n^3 + 2n^2 + 5n + 7$$

$$g(n) = n^3$$

$$\lim_{n \rightarrow \infty} \frac{3n^3 + 2n^2 + 5n + 7}{n^3} = \frac{\infty}{\infty} \quad \text{indeterminate form}$$

To get out of indeterminate situation, take
the first derivate and then apply the limits
Keep taking the derivative until
you get out of the indeterminate situation

L'Hôpital's rule for $\frac{\infty}{\infty}$ and $\frac{0}{0}$ indeterminate forms

$$\lim_{n \rightarrow \infty} \frac{9n^2 + 4n + 5}{3n^2}$$

$$\lim_{n \rightarrow \infty} \frac{18n + 4}{6n}$$

$$f(n) = O(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq \infty$$

$$f(n) = \Omega(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq 0$$

$$f(n) = \Theta(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq 0, \infty$$

$$f(n) = o(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0$$

$$f(n) = \omega(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \infty$$

$$f(n) \sim g(n) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 1$$



How to perform asymptotic analysis

1. Using limits formulas

$$f(n) = 3n^3 + 2n^2 + 5n + 7$$

$$g(n) = n^3$$

$$\lim_{n \rightarrow \infty} \frac{3n^3 + 2n^2 + 5n + 7}{n^3} = \frac{\infty}{\infty} \quad \text{indeterminate form}$$

To get out of indeterminate situation, take
the first derivate and then apply the limits
Keep taking the derivative until
you get out of the indeterminate situation

L'Hôpital's rule for $\frac{\infty}{\infty}$ and $\frac{0}{0}$ indeterminate forms

$$\lim_{n \rightarrow \infty} \frac{9n^2 + 4n + 5}{3n^2}$$

$$\lim_{n \rightarrow \infty} \frac{18n + 4}{6n}$$

$$\lim_{n \rightarrow \infty} \frac{18}{6} = 3$$

$$f(n) = O(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq \infty$$

$$f(n) = \Omega(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq 0$$

$$f(n) = \Theta(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq 0, \infty$$

$$f(n) = o(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0$$

$$f(n) = \omega(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \infty$$

$$f(n) \sim g(n) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 1$$



How to perform asymptotic analysis

1. Using limits formulas

$$f(n) = 3n^3 + 2n^2 + 5n + 7$$

$$g(n) = n^3$$

$$\lim_{n \rightarrow \infty} \frac{3n^3 + 2n^2 + 5n + 7}{n^3} = \frac{\infty}{\infty} \quad \text{indeterminate form}$$

To get out of indeterminate situation, take the first derivate and then apply the limits
Keep taking the derivative until you get out of the indeterminate situation

L'Hôpital's rule for $\frac{\infty}{\infty}$ and $\frac{0}{0}$ indeterminate forms

$$\lim_{n \rightarrow \infty} \frac{9n^2 + 4n + 5}{3n^2}$$

$$f(n) = O(n^3) \quad \text{YES}$$

$$\lim_{n \rightarrow \infty} \frac{18n + 4}{6n}$$

$$\lim_{n \rightarrow \infty} \frac{18}{6} = 3$$

$$f(n) = O(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq \infty$$

$$f(n) = \Omega(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq 0$$

$$f(n) = \Theta(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq 0, \infty$$

$$f(n) = o(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0$$

$$f(n) = \omega(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \infty$$

$$f(n) \sim g(n) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 1$$



How to perform asymptotic analysis

1. Using limits formulas

$$f(n) = 3n^3 + 2n^2 + 5n + 7$$

$$g(n) = n^3$$

$$\lim_{n \rightarrow \infty} \frac{3n^3 + 2n^2 + 5n + 7}{n^3} = \frac{\infty}{\infty} \quad \text{indeterminate form}$$

To get out of indeterminate situation, take the first derivate and then apply the limits
Keep taking the derivative until you get out of the indeterminate situation

L'Hôpital's rule for $\frac{\infty}{\infty}$ and $\frac{0}{0}$ indeterminate forms

$$\lim_{n \rightarrow \infty} \frac{9n^2 + 4n + 5}{3n^2}$$

$$f(n) = O(n^3) \quad \text{YES}$$

$$f(n) = \Omega(n^3) \quad \text{YES}$$

$$\lim_{n \rightarrow \infty} \frac{18n + 4}{6n}$$

$$\lim_{n \rightarrow \infty} \frac{18}{6} = 3$$

$$f(n) = O(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq \infty$$

$$f(n) = \Omega(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq 0$$

$$f(n) = \Theta(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq 0, \infty$$

$$f(n) = o(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0$$

$$f(n) = \omega(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \infty$$

$$f(n) \sim g(n) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 1$$



How to perform asymptotic analysis

1. Using limits formulas

$$f(n) = 3n^3 + 2n^2 + 5n + 7$$

$$g(n) = n^3$$

$$\lim_{n \rightarrow \infty} \frac{3n^3 + 2n^2 + 5n + 7}{n^3} = \frac{\infty}{\infty} \quad \text{indeterminate form}$$

To get out of indeterminate situation, take the first derivate and then apply the limits
Keep taking the derivative until you get out of the indeterminate situation

L'Hôpital's rule for $\frac{\infty}{\infty}$ and $\frac{0}{0}$ indeterminate forms

$$\lim_{n \rightarrow \infty} \frac{9n^2 + 4n + 5}{3n^2}$$

$$f(n) = O(n^3) \quad \text{YES}$$

$$\lim_{n \rightarrow \infty} \frac{18n + 4}{6n}$$

$$f(n) = \Omega(n^3) \quad \text{YES}$$

$$\lim_{n \rightarrow \infty} \frac{18}{6} = 3$$

$$f(n) = \Theta(n^3) \quad \text{YES}$$

$$f(n) = O(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq \infty$$

$$f(n) = \Omega(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq 0$$

$$f(n) = \Theta(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq 0, \infty$$

$$f(n) = o(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0$$

$$f(n) = \omega(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \infty$$

$$f(n) \sim g(n) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 1$$

How to perform asymptotic analysis

1. Using limits formulas

$$f(n) = 3n^3 + 2n^2 + 5n + 7$$

$$g(n) = n^3$$

$$\lim_{n \rightarrow \infty} \frac{3n^3 + 2n^2 + 5n + 7}{n^3} = \frac{\infty}{\infty} \quad \text{indeterminate form}$$

To get out of indeterminate situation, take the first derivate and then apply the limits
Keep taking the derivative until you get out of the indeterminate situation

L'Hôpital's rule for $\frac{\infty}{\infty}$ and $\frac{0}{0}$ indeterminate forms

$$\lim_{n \rightarrow \infty} \frac{9n^2 + 4n + 5}{3n^2}$$

$$\lim_{n \rightarrow \infty} \frac{18n + 4}{6n}$$

$$\lim_{n \rightarrow \infty} \frac{18}{6} = 3$$

$$f(n) = O(n^3) \quad \text{YES}$$

$$f(n) = \Omega(n^3) \quad \text{YES}$$

$$f(n) = \Theta(n^3) \quad \text{YES}$$

$$f(n) = O(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq \infty$$

$$f(n) = \Omega(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq 0$$

$$f(n) = \Theta(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq 0, \infty$$

$$f(n) = o(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0$$

$$f(n) = \omega(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \infty$$


$$f(n) \sim g(n) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 1$$

$$f(n) = o(n^3) \quad \text{NO}$$


$$f(n) = \omega(n^3) \quad \text{NO}$$




Visual inspection (O)




Visual inspection (O)




Visual inspection (O)




Visual inspection(O)




Visual inspection (O)




Visual inspection (O)




Visual inspection Ω




Visual inspection Ω




Visual inspection Ω




Visual inspection Ω



Visual inspection Ω



Visual inspection Ω



Visual Inspection

$$f(n) = 3n^3 + 2n^2 + 5n + 7$$

$$1 < \log(n) < \sqrt{n} < n < n\log(n) < n^2 < \dots < n^3 < \dots < 2^n < e^n < \dots < n^n$$



Visual Inspection

$$f(n) = 3n^3 + 2n^2 + 5n + 7$$

$$1 < \log(n) < \sqrt{n} < n < n\log(n) < n^2 < \dots < n^3 < \dots < 2^n < e^n < \dots < n^n$$

O

Visual Inspection

$$f(n) = 3n^3 + 2n^2 + 5n + 7$$

$$\underbrace{1 < \log(n) < \sqrt{n} < n < n \log(n) < n^2 < \dots < n^3}_{\Omega} < \dots < 2^n < e^n < \dots < n^n \underbrace{\phantom{1 < \log(n) < \sqrt{n} < n < n \log(n) < n^2 < \dots < n^3}}_O$$



Visual Inspection

$$f(n) = 3n^3 + 2n^2 + 5n + 7$$

$$\underbrace{1 < \log(n) < \sqrt{n} < n < n \log(n) < n^2 < \dots < n^3}_{\Omega} < \dots < 2^n < e^n < \dots < n^n \underbrace{\hspace{10em}}_O$$

While it is mathematically correct to say that $f(n) = O(2^n)$ or even $f(n) = O(n^n)$, it does not capture the growth rate in a useful way



Visual Inspection

$$f(n) = 3n^3 + 2n^2 + 5n + 7$$

$$1 < \log(n) < \sqrt{n} < n < n\log(n) < n^2 < \dots < n^3 < \dots < 2^n < e^n < \dots < n^n$$

The diagram shows a sequence of growth rates: $1 < \log(n) < \sqrt{n} < n < n\log(n) < n^2 < \dots < n^3 < \dots < 2^n < e^n < \dots < n^n$. A bracket under the first four terms ($1, \log(n), \sqrt{n}, n$) is labeled with the Greek letter Ω . A bracket under the last four terms ($n^2, \dots, 2^n, e^n, n^n$) is labeled with the letter O . An arrow points from the text below towards the O bracket.

While it is mathematically correct to say that $f(n) = O(2^n)$ or even $f(n) = O(n^n)$, it does not capture the growth rate in a useful way

In general, the most useful asymptotic notation is one that accurately captures the dominant term or terms of the function, e.g. the smallest upper bound or the largest lower bound.

Example

$$f(n) = \sqrt{6n^3 + 7n^2 + 5n + 5}$$

$$f(n) = \mathbf{O}(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq \infty$$

$$f(n) = \mathbf{\Omega}(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq 0$$

$$f(n) = \mathbf{\Theta}(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq 0, \infty$$

$$f(n) = \mathbf{o}(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0$$

$$f(n) = \omega(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \infty$$

$$f(n) \sim g(n) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 1$$



Example

$$f(n) = n^3 \log_2 n$$

$$g(n) = 3n \log_8 n$$

$$f(n) = \mathbf{O}(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq \infty$$

$$f(n) = \mathbf{\Omega}(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq 0$$

$$f(n) = \mathbf{\Theta}(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq 0, \infty$$

$$f(n) = \mathbf{o}(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0$$

$$f(n) = \omega(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \infty$$

$$f(n) \sim g(n) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 1$$



Example

$$f(n) = 8^n$$

$$g(n) = 4^n$$

$$f(n) = \mathbf{O}(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq \infty$$

$$f(n) = \mathbf{\Omega}(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq 0$$

$$f(n) = \mathbf{\Theta}(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq 0, \infty$$

$$f(n) = \mathbf{o}(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0$$

$$f(n) = \omega(g(n)) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \infty$$

$$f(n) \sim g(n) \quad \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 1$$

