

Algorithm & Data Structure Analysis

Lecture 1: Growth of Function

Appetizer

Sorting: Why should I care?




Which one do you prefer?




Measuring efficiency

- Let n be the input size
- Runtime
- Memory usage
- (usually interested in large n)


Logarithmic runtime




Linear runtime




Quadratic runtime




Cubic runtime



Exponential runtime



Growth of function



Asymptotic notations

Asymptotic notations

$$O(g(n)) = \{f(n) : \exists c > 0, \exists n_0 \in \mathbb{N}_+, \forall n \geq n_0, f(n) \leq c \cdot g(n)\}$$

Asymptotic notations

$$O(g(n)) = \{f(n) : \exists c > 0, \exists n_0 \in \mathbb{N}_+, \forall n \geq n_0, f(n) \leq c \cdot g(n)\}$$

$$\Omega(g(n)) = \{f(n) : \exists c > 0, \exists n_0 \in \mathbb{N}_+, \forall n \geq n_0, f(n) \geq c \cdot g(n)\}$$

Asymptotic notations

$$O(g(n)) = \{f(n) : \exists c > 0, \exists n_0 \in \mathbb{N}_+, \forall n \geq n_0, f(n) \leq c \cdot g(n)\}$$

$$\Omega(g(n)) = \{f(n) : \exists c > 0, \exists n_0 \in \mathbb{N}_+, \forall n \geq n_0, f(n) \geq c \cdot g(n)\}$$

$$f(n) = \Theta(g(n)) \text{ if and only if } f(n) = O(g(n)) \text{ and } f(n) = \Omega(g(n))$$

Asymptotic notations

$$O(g(n)) = \{f(n) : \exists c > 0, \exists n_0 \in \mathbb{N}_+, \forall n \geq n_0, f(n) \leq c \cdot g(n)\}$$

$$\Omega(g(n)) = \{f(n) : \exists c > 0, \exists n_0 \in \mathbb{N}_+, \forall n \geq n_0, f(n) \geq c \cdot g(n)\}$$

$f(n) = \Theta(g(n))$ if and only if $f(n) = O(g(n))$ and $f(n) = \Omega(g(n))$

$$o(g(n)) = \{f(n) : \forall c > 0, \exists n_0 \in \mathbb{N}_+, \forall n \geq n_0, f(n) < c \cdot g(n)\}$$

Asymptotic notations

$$O(g(n)) = \{f(n) : \exists c > 0, \exists n_0 \in \mathbb{N}_+, \forall n \geq n_0, f(n) \leq c \cdot g(n)\}$$

$$\Omega(g(n)) = \{f(n) : \exists c > 0, \exists n_0 \in \mathbb{N}_+, \forall n \geq n_0, f(n) \geq c \cdot g(n)\}$$

$$f(n) = \Theta(g(n)) \text{ if and only if } f(n) = O(g(n)) \text{ and } f(n) = \Omega(g(n))$$

$$o(g(n)) = \{f(n) : \forall c > 0, \exists n_0 \in \mathbb{N}_+, \forall n \geq n_0, f(n) < c \cdot g(n)\}$$

$$\omega(g(n)) = \{f(n) : \forall c > 0, \exists n_0 \in \mathbb{N}_+, \forall n \geq n_0, f(n) > c \cdot g(n)\}$$

Examples

$$5n$$

Examples

$$5n \quad : \quad O(n), \Omega(n), \Theta(n), o(n \log n), \omega(\sqrt{n})$$

Examples

$$5n \quad : \quad O(n), \Omega(n), \Theta(n), o(n \log n), \omega(\sqrt{n})$$

$$n^2 - n \log n$$

Examples

$$5n \quad : \quad O(n), \Omega(n), \Theta(n), o(n \log n), \omega(\sqrt{n})$$

$$n^2 - n \log n : O(n^2), \Omega(n^2), \Theta(n^2), o(n^3), \omega(n \log n)$$

Examples

$$5n \quad : \quad O(n), \Omega(n), \Theta(n), o(n \log n), \omega(\sqrt{n})$$

$$n^2 - n \log n : O(n^2), \Omega(n^2), \Theta(n^2), o(n^3), \omega(n \log n)$$

$$100n$$

Examples

$$5n \quad : \quad O(n), \Omega(n), \Theta(n), o(n \log n), \omega(\sqrt{n})$$

$$n^2 - n \log n : O(n^2), \Omega(n^2), \Theta(n^2), o(n^3), \omega(n \log n)$$

$$100n \quad : \quad O(n^2), \Omega(\sqrt{n}), \Theta(n), o(n \log n), \omega(\sqrt{n})$$

True of false

$$5n \log n \in O(n \log n)$$

True or false

$5n \log n \in O(n \log n)$ True

True of false

$5n \log n \in O(n \log n)$ True

$5n \log n \in O(n^2)$

True of false

$5n \log n \in O(n \log n)$ True

$5n \log n \in O(n^2)$ True

True of false

$5n \log n \in O(n \log n)$ True

$5n \log n \in O(n^2)$ True

$5n \log n \in \Omega(n^2)$

True of false

$5n \log n \in O(n \log n)$ True

$5n \log n \in O(n^2)$ True

$5n \log n \in \Omega(n^2)$ False

True of false

$5n \log n \in O(n \log n)$ True

$5n \log n \in O(n^2)$ True

$5n \log n \in \Omega(n^2)$ False

$5n \log n \in o(n^2)$

True of false

$5n \log n \in O(n \log n)$ True

$5n \log n \in O(n^2)$ True

$5n \log n \in \Omega(n^2)$ False

$5n \log n \in o(n^2)$ True

True of false

$5n \log n \in O(n \log n)$ True

$5n \log n \in O(n^2)$ True

$5n \log n \in \Omega(n^2)$ False

$5n \log n \in o(n^2)$ True

$5n \log n + n^2 \in O(n \log n)$

True of false

$5n \log n \in O(n \log n)$ True

$5n \log n \in O(n^2)$ True

$5n \log n \in \Omega(n^2)$ False

$5n \log n \in o(n^2)$ True

$5n \log n + n^2 \in O(n \log n)$ False

True of false

$5n \log n \in O(n \log n)$ True

$5n \log n \in O(n^2)$ True

$5n \log n \in \Omega(n^2)$ False

$5n \log n \in o(n^2)$ True

$5n \log n + n^2 \in O(n \log n)$ False

$5n \log n + n^2 \in O(n^2)$

True of false

$5n \log n \in O(n \log n)$ True

$5n \log n \in O(n^2)$ True

$5n \log n \in \Omega(n^2)$ False

$5n \log n \in o(n^2)$ True

$5n \log n + n^2 \in O(n \log n)$ False

$5n \log n + n^2 \in O(n^2)$ True

Summary

- Efficiency of data structures and algorithms
- Growth function given input size
- Asymptotic behavior and complexity classes
- Reading: Introduction to Algorithms
 - ▶ Chapter 3.1: Growth of Functions

Next lecture

- Integer arithmetics
- Reading: Algorithms and Data Structures
 - ▶ Chapter 1.1: Addition
 - ▶ Chapter 1.2: Multiplication: The School Method