

# PCA Numerical Example

---

Let's understand steps of PCA with an example of two dimensional data.

We start with the following set of 5 points:

$$\mathbf{X} = \begin{bmatrix} 4 & 6 & -1 & -2 & -3 \\ 5 & 9 & -7 & -5 & -9 \end{bmatrix}$$

This is a more compact way of writing (using vectors & matrices); otherwise you will have many summations to deal with.

# PCA Numerical Example

---

Calculate mean:

$$\bar{X} = \begin{bmatrix} 0.8 \\ -1.4 \end{bmatrix}$$

# PCA Numerical Example

---

Center the points by performing  $(X - \bar{X})$

$$\begin{bmatrix} 3.2 & 5.2 & -1.8 & -2.8 & -3.8 \\ 6.4 & 10.4 & -5.6 & -3.6 & -7.6 \end{bmatrix}$$

# PCA Numerical Example

---

The covariance matrix A is:  $1/n * (X - \bar{X})(X - \bar{X})^T$   
where n is the number of points (=5).

$$A = \begin{bmatrix} 12.56 & 24.72 \\ 24.72 & 50.24 \end{bmatrix}$$

# PCA Numerical Example

---

Finding the eigenvalues:

$$\begin{bmatrix} 12.56 & 24.72 \\ 24.72 & 50.24 \end{bmatrix} - \lambda I$$

I is the identity matrix.  $I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

# PCA Numerical Example

---

Finding the eigenvalues:

$$\begin{bmatrix} 12.56 - \lambda & 24.72 \\ 24.72 & 50.24 - \lambda \end{bmatrix}$$

# PCA Numerical Example

---

Finding the eigenvalues:

$$\text{Determinant of } \begin{bmatrix} 12.56 - \lambda & 24.72 \\ 24.72 & 50.24 - \lambda \end{bmatrix} = 0$$

# PCA Numerical Example

---

Finding the eigenvalues:

$$(12.56 - \lambda) * (50.24 - \lambda) - 24.72^2 = 0$$



# PCA Numerical Example

---

Finding the eigenvalues:

$$(\lambda^2) - 62.8\lambda + 19.936 = 0$$

# PCA Numerical Example

---

Finding the eigenvalues:

Now, we find the roots of this quadratic equation as:

$$\frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

# PCA Numerical Example

---

Finding the eigenvalues:

This gives the roots as:

$$\lambda_1 = 62.4809$$

$$\lambda_2 = 0.3190$$

# PCA Numerical Example

---

Calculate the Eigenvector for the 1<sup>st</sup> Eigenvalue:

$$(A - \lambda_1 I) * X = 0$$

where  $X = \begin{bmatrix} a_1 \\ b_1 \end{bmatrix}$

# PCA Numerical Example

---

Calculate the Eigenvector for the 1<sup>st</sup> Eigenvalue:

$$(A - \lambda_1 I) = \begin{bmatrix} 12.56 & 24.72 \\ 24.72 & 50.24 \end{bmatrix} - \begin{bmatrix} 62.48 & 0 \\ 0 & 62.48 \end{bmatrix}$$

$$(A - \lambda_1 I) = \begin{bmatrix} -49.92 & 24.72 \\ 24.72 & -12.24 \end{bmatrix}$$

# PCA Numerical Example

---

Calculating the Eigenvector for the 1<sup>st</sup> Eigenvalue:

$$\begin{bmatrix} -49.92 & 24.72 \\ 24.72 & -12.24 \end{bmatrix} * \begin{bmatrix} a_1 \\ b_1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$a_1^2 + b_1^2 = 1$$

From these equations, we can get the 1<sup>st</sup> eigenvector.

# PCA Numerical Example

Calculating the Eigenvector for the 1<sup>st</sup> Eigenvalue:

$$\begin{bmatrix} -49.92 & 24.72 \\ 24.72 & -12.24 \end{bmatrix} * \begin{bmatrix} a_1 \\ b_1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \Rightarrow 24.72 a_1 = 12.24 b_1 \Rightarrow a_1 = 0.4951 b_1$$

$$a_1^2 + b_1^2 = 1 \Rightarrow 0.2452 b_1^2 + b_1^2 = 1 \Rightarrow 1.2452 b_1^2 = 1 \Rightarrow b_1^2 = 0.8031 \\ \Rightarrow b_1 = + \text{ or } - 0.8962 \Rightarrow a_1 = 0.4951 \times +/\text{- } 0.8962 = 0.4437$$

1<sup>st</sup> Eigen vector  $\begin{pmatrix} 0.4437 \\ 0.8962 \end{pmatrix}$

# PCA Numerical Example

---

Calculate the Eigenvector for the 2<sup>nd</sup> Eigenvalue:

$$(A - \lambda_2 I) * X = 0$$

where  $X = \begin{bmatrix} a_2 \\ b_2 \end{bmatrix}$



# PCA Numerical Example

---

Calculate the Eigenvector for the 2<sup>nd</sup> Eigenvalue:

$$(A - \lambda_2 I) = \begin{bmatrix} 12.56 & 24.72 \\ 24.72 & 50.24 \end{bmatrix} - \begin{bmatrix} 0.03 & 0 \\ 0 & 0.03 \end{bmatrix}$$

$$(A - \lambda_2 I) = \begin{bmatrix} 12.53 & 24.72 \\ 24.72 & 50.21 \end{bmatrix}$$

# PCA Numerical Example

---

Calculate the Eigenvector for the 2<sup>nd</sup> Eigenvalue:

$$\begin{bmatrix} 12.53 & 24.72 \\ 24.72 & 50.21 \end{bmatrix} * \begin{bmatrix} a_2 \\ b_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$a_2^2 + b_2^2 = 1$$

From these equations, we can get the 2<sup>nd</sup> eigenvector.

# PCA Numerical Example

---

Calculate the Eigenvector for the 2<sup>nd</sup> Eigenvalue:

$$\begin{bmatrix} 12.53 & 24.72 \\ 24.72 & 50.21 \end{bmatrix} * \begin{bmatrix} a_2 \\ b_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \Rightarrow 12.53 a_2 = -24.72 b_2 \Rightarrow a_2 = -1.97 b_2$$

$$a_2^2 + b_2^2 = 1 \Rightarrow 3.89 b_2^2 + b_2^2 = 1 \Rightarrow 4.89 b_2^2 = 1 \Rightarrow b_2^2 = 0.2045 \\ \Rightarrow b_2 = + \text{ or } - 0.4522 \Rightarrow a_2 = -1.97 \times +/ - 0.4522 = 0.8909$$

From these equations, we can get the 2<sup>nd</sup> eigenvector.

$$\text{2<sup>nd</sup> Eigen vector} \begin{pmatrix} 0.8909 \\ -0.4522 \end{pmatrix}$$

# PCA Numerical Notes

---

The variance of projections in the line of the principal components is equal to the Eigenvalues of the principal components.

Sum of all the Eigenvalues is equal to the total variance of all the variance.

1<sup>st</sup> eigen value is bigger than 2<sup>nd</sup>, 2<sup>nd</sup> is bigger than 3<sup>rd</sup> and so on.