


Improved bounds for the sunflower lemma

Kewen Wu
Peking University

Joint work with

Ryan Alweiss
Princeton



Shachar Lovett
UCSD



Jiapeng Zhang
Harvard



Definitions

Definition (w -set system and r -sunflower)

A w -set system is a family of sets of size at most w .

An r -sunflower is r sets S_1, \dots, S_r where

- **Kernel:** $Y = S_1 \cap \dots \cap S_r$;
- **Petals:** $S_1 \setminus Y, \dots, S_r \setminus Y$ are pairwise disjoint.

Definitions

Definition (w -set system and r -sunflower)

A w -set system is a family of sets of size at most w .

An r -sunflower is r sets S_1, \dots, S_r where

- **Kernel:** $Y = S_1 \cap \dots \cap S_r$;
- **Petals:** $S_1 \setminus Y, \dots, S_r \setminus Y$ are pairwise disjoint.

Example

$\{\{1, 2\}, \{1, 3, 4, 6\}, \{1, 5\}, \{2, 3\}\}$ is a 4-set system of size 4.

It has a 3-sunflower $\{\{1, 2\}, \{1, 3, 4, 6\}, \{1, 5\}\}$ with kernel $\{1\}$ and petals $\{2\}, \{3, 4, 6\}, \{5\}$.

Main result

Theorem (Erdős-Rado sunflower)

Any w -set system of size s has an r -sunflower.

Main result

Theorem (Erdős-Rado sunflower)

Any w -set system of size s has an r -sunflower.

Let's focus on $r = 3$.

- Erdős and Rado 1960: $s = w! \cdot 2^w \approx w^w$.
- Kostochka 2000: $s \approx (w \log \log \log w / \log \log w)^w$.
- Fukuyama 2018: $s \approx w^{0.75w}$.
- Now: $s \approx (\log w)^w$ and this is tight for our approach.

Actual bound and further refinement

Theorem (Improved sunflower lemma)

For some constant C , any w -set system of size s has an r -sunflower, where

$$s = (Cr^2 \cdot (\log w \log \log w + (\log r)^2))^w.$$

Actual bound and further refinement

Theorem (Improved sunflower lemma)

For some constant C , any w -set system of size s has an r -sunflower, where

$$s = (Cr^2 \cdot (\log w \log \log w + (\log r)^2))^w.$$

Recently, Anup Rao improved it to

$$s = (Cr(\log w + \log r))^w.$$

Applications – Theoretical computer science

- Circuit lower bounds
- Data structure lower bounds
- Matrix multiplication
- Pseudorandomness
- Cryptography
- Property testing
- Fixed parameter complexity
- Communication complexity
- ...

Applications – Combinatorics

- Erdős-Szemerédi sunflower lemma
- Intersecting set systems
- Packing Kneser graphs
- Alon-Jaeger-Tarsi nowhere-zero conjecture
- Thresholds in random graphs
- ...

Section 3

Proof overview

Make it robust

Assume $\mathcal{F} = \{S_1, \dots, S_m\}$ is a w -set system.

Define a width- w DNF $f_{\mathcal{F}}$ as $f_{\mathcal{F}} = \bigvee_{i=1}^m \bigwedge_{j \in S_i} x_j$.

Example

If $\mathcal{F} = \{\{1, 2\}, \{1, 3, 4, 6\}, \{1, 5\}, \{2, 3\}\}$, then

$$f_{\mathcal{F}} = (x_1 \wedge x_2) \vee (x_1 \wedge x_3 \wedge x_4 \wedge x_6) \vee (x_1 \wedge x_5) \vee (x_2 \wedge x_3).$$

Make it robust

Assume $\mathcal{F} = \{S_1, \dots, S_m\}$ is a w -set system.

Define a width- w DNF $f_{\mathcal{F}}$ as $f_{\mathcal{F}} = \bigvee_{i=1}^m \bigwedge_{j \in S_i} x_j$.

Example

If $\mathcal{F} = \{\{1, 2\}, \{1, 3, 4, 6\}, \{1, 5\}, \{2, 3\}\}$, then

$$f_{\mathcal{F}} = (x_1 \wedge x_2) \vee (x_1 \wedge x_3 \wedge x_4 \wedge x_6) \vee (x_1 \wedge x_5) \vee (x_2 \wedge x_3).$$

Definition (Satisfying system)

\mathcal{F} is satisfying if $\Pr[f_{\mathcal{F}}(x) = 0] < 1/3$ with $\Pr[x_i = 1] = 1/3$,
i.e., $\Pr[\forall i \in [m], S_i \not\subseteq S] < 1/3$ with $\Pr[x_i \in S] = 1/3$.

Satisfyingness implies sunflower

Assume \mathcal{F} is a set system on ground set $\{x_1, \dots, x_n\}$.

Lemma

If \mathcal{F} is satisfying, then it has 3 pairwise disjoint sets.

In particular, 3 pairwise disjoint sets is a 3-sunflower.

Satisfyingness implies sunflower

Assume \mathcal{F} is a set system on ground set $\{x_1, \dots, x_n\}$.

Lemma

If \mathcal{F} is satisfying, then it has 3 pairwise disjoint sets.

In particular, 3 pairwise disjoint sets is a 3-sunflower.

Proof.

Color x_1, \dots, x_n to red, green, blue uniformly and independently.

Satisfyingness implies sunflower

Assume \mathcal{F} is a set system on ground set $\{x_1, \dots, x_n\}$.

Lemma

If \mathcal{F} is satisfying, then it has 3 pairwise disjoint sets.

In particular, 3 pairwise disjoint sets is a 3-sunflower.

Proof.

Color x_1, \dots, x_n to red, green, blue uniformly and independently.
By definition, \mathcal{F} contains a purely red (green/blue) set w.p $> 2/3$.

Satisfyingness implies sunflower

Assume \mathcal{F} is a set system on ground set $\{x_1, \dots, x_n\}$.

Lemma

If \mathcal{F} is satisfying, then it has 3 pairwise disjoint sets.

In particular, 3 pairwise disjoint sets is a 3-sunflower.

Proof.

Color x_1, \dots, x_n to red, green, blue uniformly and independently. By definition, \mathcal{F} contains a purely red (green/blue) set w.p $> 2/3$. By union bound, \mathcal{F} contains one purely red set, one purely green set, and one purely blue set w.p > 0 . □

Structure vs pseudorandomness

Assume $\mathcal{F} = \{S_1, \dots, S_m\}$, $m > \kappa^w$ is a w -set system. Define link $\mathcal{F}_Y = \{S_i \setminus Y \mid Y \subset S_i\}$, which is a $(w - |Y|)$ -set system.

Example

If $\mathcal{F} = \{\{1, 2\}, \{1, 3, 4\}, \{1, 5\}, \{2, 3\}\}$, then $\mathcal{F}_{\{2\}} = \{\{1\}, \{3\}\}$.

Structure vs pseudorandomness

Assume $\mathcal{F} = \{S_1, \dots, S_m\}$, $m > \kappa^w$ is a w -set system. Define link $\mathcal{F}_Y = \{S_i \setminus Y \mid Y \subset S_i\}$, which is a $(w - |Y|)$ -set system.

Example

If $\mathcal{F} = \{\{1, 2\}, \{1, 3, 4\}, \{1, 5\}, \{2, 3\}\}$, then $\mathcal{F}_{\{2\}} = \{\{1\}, \{3\}\}$.

If there exists Y such that $|\mathcal{F}_Y| \geq m/\kappa^{|Y|} > \kappa^{w-|Y|}$, then we can apply induction and find 3-sunflower in \mathcal{F}_Y .

Structure vs pseudorandomness

Assume $\mathcal{F} = \{S_1, \dots, S_m\}$, $m > \kappa^w$ is a w -set system. Define link $\mathcal{F}_Y = \{S_i \setminus Y \mid Y \subset S_i\}$, which is a $(w - |Y|)$ -set system.

Example

If $\mathcal{F} = \{\{1, 2\}, \{1, 3, 4\}, \{1, 5\}, \{2, 3\}\}$, then $\mathcal{F}_{\{2\}} = \{\{1\}, \{3\}\}$.

If there exists Y such that $|\mathcal{F}_Y| \geq m/\kappa^{|Y|} > \kappa^{w-|Y|}$, then we can apply induction and find 3-sunflower in \mathcal{F}_Y .

So induction starts at such \mathcal{F} , that $|\mathcal{F}_Y| < m/\kappa^{|Y|}$ holds for any Y .

Lemma

Let $\kappa \geq (\log w)^{O(1)}$. If $|\mathcal{F}_Y| < m/\kappa^{|Y|}$ holds for any Y , then \mathcal{F} is satisfying, which means \mathcal{F} has 3 pairwise disjoint sets.

Randomness preserves pseudorandomness

Let $\mathcal{F} = \{S_1, \dots, S_m\}$ be a w -(multi-)set system.

Randomness preserves pseudorandomness

Let $\mathcal{F} = \{S_1, \dots, S_m\}$ be a w -(multi-)set system.

Assume $|\mathcal{F}_Y| < m/\kappa^{|Y|}$ holds for any Y . $\Leftarrow \mathcal{F}$ is pseudorandom

Randomness preserves pseudorandomness

Let $\mathcal{F} = \{S_1, \dots, S_m\}$ be a w -(multi-)set system.

Assume $|\mathcal{F}_Y| < m/\kappa^{|Y|}$ holds for any Y . $\Leftarrow \mathcal{F}$ is pseudorandom

Take $\approx 1/\sqrt{\kappa}$ -fraction of the ground set as W ,

and construct a $w/2$ -(multi-)set system \mathcal{F}' from each S_i :

Randomness preserves pseudorandomness

Let $\mathcal{F} = \{S_1, \dots, S_m\}$ be a w -(multi-)set system.

Assume $|\mathcal{F}_Y| < m/\kappa^{|Y|}$ holds for any Y . $\Leftarrow \mathcal{F}$ is pseudorandom

Take $\approx 1/\sqrt{\kappa}$ -fraction of the ground set as W ,

and construct a $w/2$ -(multi-)set system \mathcal{F}' from each S_i :

- Good: If there exists $|S_j \setminus W| \leq w/2$ and $S_j \setminus W \subset S_i \setminus W$, then put $S_j \setminus W$ into \mathcal{F}' ; (j may equal i)

Randomness preserves pseudorandomness

Let $\mathcal{F} = \{S_1, \dots, S_m\}$ be a w -(multi-)set system.

Assume $|\mathcal{F}_Y| < m/\kappa^{|Y|}$ holds for any Y . $\Leftarrow \mathcal{F}$ is pseudorandom

Take $\approx 1/\sqrt{\kappa}$ -fraction of the ground set as W ,

and construct a $w/2$ -(multi-)set system \mathcal{F}' from each S_i :

- Good: If there exists $|S_j \setminus W| \leq w/2$ and $S_j \setminus W \subset S_i \setminus W$, then put $S_j \setminus W$ into \mathcal{F}' ; (j may equal i)

E.g., $S_j \setminus W = \{1\}$, $S_i \setminus W = \{1, 2, 3, 4, 5\}$.

Randomness preserves pseudorandomness

Let $\mathcal{F} = \{S_1, \dots, S_m\}$ be a w -(multi-)set system.

Assume $|\mathcal{F}_Y| < m/\kappa^{|Y|}$ holds for any Y . $\Leftarrow \mathcal{F}$ is pseudorandom

Take $\approx 1/\sqrt{\kappa}$ -fraction of the ground set as W ,

and construct a $w/2$ -(multi-)set system \mathcal{F}' from each S_i :

- Good: If there exists $|S_j \setminus W| \leq w/2$ and $S_j \setminus W \subset S_i \setminus W$, then put $S_j \setminus W$ into \mathcal{F}' ; (j may equal i)

E.g., $S_j \setminus W = \{1\}$, $S_i \setminus W = \{1, 2, 3, 4, 5\}$.

- Bad: otherwise, we do nothing for S_i .

Example

If $\mathcal{F} = \{\{1, 2\}, \{1, 3\}, \{2, 3, 4\}, \{4, 5, 6, 7\}\}$ and $w = 4$, $W = \{1\}$, then $\mathcal{F}' = \{\{1, 2\}, \{1, 3\}, \{2, 3, 4\}, \{4, 5, 6, 7\}\}$.

One reduction step

Then $|\mathcal{F}'_Y| \leq |\mathcal{F}_Y|$ and $|\mathcal{F}'| \approx |\mathcal{F}|$. $\Leftarrow \mathcal{F}'$ is also pseudorandom

One reduction step

Then $|\mathcal{F}'_Y| \leq |\mathcal{F}_Y|$ and $|\mathcal{F}'| \approx |\mathcal{F}|$. $\Leftarrow \mathcal{F}'$ is also pseudorandom
Prove by encoding **bad** $(W, i) \rightarrow (W' = W \cup S_i, \text{aux}_1, k, \text{aux}_2)$,
where S_i ranks $k < |\mathcal{F}|/\kappa^{w/2}$ in $\mathcal{F}_{S_j \cap S_i}$ for the first $j \leq i$ that
 $S_j \setminus W \subset S_i \setminus W$.

One reduction step

Then $|\mathcal{F}'_Y| \leq |\mathcal{F}_Y|$ and $|\mathcal{F}'| \approx |\mathcal{F}|$. $\Leftarrow \mathcal{F}'$ is also pseudorandom
Prove by encoding **bad** $(W, i) \rightarrow (W' = W \cup S_i, \text{aux}_1, k, \text{aux}_2)$,
where S_i ranks $k < |\mathcal{F}|/\kappa^{w/2}$ in $\mathcal{F}_{S_j \cap S_i}$ for the first $j \leq i$ that
 $S_j \setminus W \subset S_i \setminus W$.

Example

$\mathcal{F}' = \{\{1, 2\}, \{2, 3, 4\}, \{1, 4, 5, 6\}, \{4, 5, 6, 7\}\}, W = \{1\}, i = 4$.
Encode/decode **bad** pair (W, i) :

One reduction step

Then $|\mathcal{F}'_Y| \leq |\mathcal{F}_Y|$ and $|\mathcal{F}'| \approx |\mathcal{F}|$. $\Leftarrow \mathcal{F}'$ is also pseudorandom
Prove by encoding **bad** $(W, i) \rightarrow (W' = W \cup S_i, \text{aux}_1, k, \text{aux}_2)$,
where S_i ranks $k < |\mathcal{F}|/\kappa^{w/2}$ in $\mathcal{F}_{S_j \cap S_i}$ for the first $j \leq i$ that
 $S_j \setminus W \subset S_i \setminus W$.

Example

$\mathcal{F}' = \{\{1, 2\}, \{2, 3, 4\}, \{1, 4, 5, 6\}, \{4, 5, 6, 7\}\}, W = \{1\}, i = 4$.
Encode/decode **bad** pair (W, i) :

- $W' = W \cup S_i = \{1, 4, 5, 6, 7\}$

One reduction step

Then $|\mathcal{F}'_Y| \leq |\mathcal{F}_Y|$ and $|\mathcal{F}'| \approx |\mathcal{F}|$. $\Leftarrow \mathcal{F}'$ is also pseudorandom
Prove by encoding **bad** $(W, i) \rightarrow (W' = W \cup S_i, \text{aux}_1, k, \text{aux}_2)$,
where S_i ranks $k < |\mathcal{F}|/\kappa^{w/2}$ in $\mathcal{F}_{S_j \cap S_i}$ for the first $j \leq i$ that
 $S_j \setminus W \subset S_i \setminus W$.

Example

$\mathcal{F}' = \{\{1, 2\}, \{2, 3, 4\}, \{1, 4, 5, 6\}, \{4, 5, 6, 7\}\}, W = \{1\}, i = 4$.
Encode/decode **bad** pair (W, i) :

- $W' = W \cup S_i = \{1, 4, 5, 6, 7\}$ we find $j = 3$ with $S_j \subset W'$

One reduction step

Then $|\mathcal{F}'_Y| \leq |\mathcal{F}_Y|$ and $|\mathcal{F}'| \approx |\mathcal{F}|$. $\Leftarrow \mathcal{F}'$ is also pseudorandom
 Prove by encoding **bad** $(W, i) \rightarrow (W' = W \cup S_i, \text{aux}_1, k, \text{aux}_2)$,
 where S_i ranks $k < |\mathcal{F}|/\kappa^{w/2}$ in $\mathcal{F}_{S_j \cap S_i}$ for the first $j \leq i$ that
 $S_j \setminus W \subset S_i \setminus W$.

Example

$\mathcal{F}' = \{\{1, 2\}, \{2, 3, 4\}, \{1, 4, 5, 6\}, \{4, 5, 6, 7\}\}, W = \{1\}, i = 4$.

Encode/decode **bad** pair (W, i) :

- $W' = W \cup S_i = \{1, 4, 5, 6, 7\}$ we find $j = 3$ with $S_j \subset W'$
- $\text{aux}_1 = \text{\$}\text{\$}\text{\$}\text{\$}$ **with at least $w/2$ $\text{\$}$ s**

One reduction step

Then $|\mathcal{F}'_Y| \leq |\mathcal{F}_Y|$ and $|\mathcal{F}'| \approx |\mathcal{F}|$. $\Leftarrow \mathcal{F}'$ is also pseudorandom
 Prove by encoding **bad** $(W, i) \rightarrow (W' = W \cup S_i, \text{aux}_1, k, \text{aux}_2)$,
 where S_i ranks $k < |\mathcal{F}|/\kappa^{w/2}$ in $\mathcal{F}_{S_j \cap S_i}$ for the first $j \leq i$ that
 $S_j \setminus W \subset S_i \setminus W$.

Example

$\mathcal{F}' = \{\{1, 2\}, \{2, 3, 4\}, \{1, 4, 5, 6\}, \{4, 5, 6, 7\}\}, W = \{1\}, i = 4$.

Encode/decode **bad** pair (W, i) :

- $W' = W \cup S_i = \{1, 4, 5, 6, 7\}$ we find $j = 3$ with $S_j \subset W'$
- $\text{aux}_1 = \$\$ \$\$$ **with at least $w/2$ \$s** we know $S_j \cap S_i = \{4, 5, 6\}$

One reduction step

Then $|\mathcal{F}'_Y| \leq |\mathcal{F}_Y|$ and $|\mathcal{F}'| \approx |\mathcal{F}|$. $\Leftarrow \mathcal{F}'$ is also pseudorandom
 Prove by encoding **bad** $(W, i) \rightarrow (W' = W \cup S_i, \text{aux}_1, k, \text{aux}_2)$,
 where S_i ranks $k < |\mathcal{F}|/\kappa^{w/2}$ in $\mathcal{F}_{S_j \cap S_i}$ for the first $j \leq i$ that
 $S_j \setminus W \subset S_i \setminus W$.

Example

$\mathcal{F}' = \{\{1, 2\}, \{2, 3, 4\}, \{1, 4, 5, 6\}, \{4, 5, 6, 7\}\}, W = \{1\}, i = 4$.

Encode/decode **bad** pair (W, i) :

- $W' = W \cup S_i = \{1, 4, 5, 6, 7\}$ we find $j = 3$ with $S_j \subset W'$
- $\text{aux}_1 = \$\$ \$\$$ **with at least $w/2$ \$s** we know $S_j \cap S_i = \{4, 5, 6\}$
- $k = 2$

One reduction step

Then $|\mathcal{F}'_Y| \leq |\mathcal{F}_Y|$ and $|\mathcal{F}'| \approx |\mathcal{F}|$. $\Leftarrow \mathcal{F}'$ is also pseudorandom
 Prove by encoding **bad** $(W, i) \rightarrow (W' = W \cup S_i, \text{aux}_1, k, \text{aux}_2)$,
 where S_i ranks $k < |\mathcal{F}|/\kappa^{w/2}$ in $\mathcal{F}_{S_j \cap S_i}$ for the first $j \leq i$ that
 $S_j \setminus W \subset S_i \setminus W$.

Example

$\mathcal{F}' = \{\{1, 2\}, \{2, 3, 4\}, \{1, 4, 5, 6\}, \{4, 5, 6, 7\}\}, W = \{1\}, i = 4$.

Encode/decode **bad** pair (W, i) :

- $W' = W \cup S_i = \{1, 4, 5, 6, 7\}$ we find $j = 3$ with $S_j \subset W'$
- $\text{aux}_1 = \$\$ \$\$$ **with at least $w/2$ $\$$ s** we know $S_j \cap S_i = \{4, 5, 6\}$
- $k = 2$ S_i ranks 2 in $\mathcal{F}_{\{4, 5, 6\}}$, we recover $i = 4$

One reduction step

Then $|\mathcal{F}'_Y| \leq |\mathcal{F}_Y|$ and $|\mathcal{F}'| \approx |\mathcal{F}|$. $\Leftarrow \mathcal{F}'$ is also pseudorandom
 Prove by encoding **bad** $(W, i) \rightarrow (W' = W \cup S_i, \text{aux}_1, k, \text{aux}_2)$,
 where S_i ranks $k < |\mathcal{F}|/\kappa^{w/2}$ in $\mathcal{F}_{S_j \cap S_i}$ for the first $j \leq i$ that
 $S_j \setminus W \subset S_i \setminus W$.

Example

$\mathcal{F}' = \{\{1, 2\}, \{2, 3, 4\}, \{1, 4, 5, 6\}, \{4, 5, 6, 7\}\}, W = \{1\}, i = 4$.

Encode/decode **bad** pair (W, i) :

- $W' = W \cup S_i = \{1, 4, 5, 6, 7\}$ we find $j = 3$ with $S_j \subset W'$
- $\text{aux}_1 = \$$$$ with at least $w/2$ **$s** we know $S_j \cap S_i = \{4, 5, 6\}$$
- $k = 2$ S_i ranks 2 in $\mathcal{F}_{\{4, 5, 6\}}$, we recover $i = 4$
- $\text{aux}_2 = \$$$$$

One reduction step

Then $|\mathcal{F}'_Y| \leq |\mathcal{F}_Y|$ and $|\mathcal{F}'| \approx |\mathcal{F}|$. $\Leftarrow \mathcal{F}'$ is also pseudorandom
 Prove by encoding **bad** $(W, i) \rightarrow (W' = W \cup S_i, \text{aux}_1, k, \text{aux}_2)$,
 where S_i ranks $k < |\mathcal{F}|/\kappa^{w/2}$ in $\mathcal{F}_{S_j \cap S_i}$ for the first $j \leq i$ that
 $S_j \setminus W \subset S_i \setminus W$.

Example

$\mathcal{F}' = \{\{1, 2\}, \{2, 3, 4\}, \{1, 4, 5, 6\}, \{4, 5, 6, 7\}\}, W = \{1\}, i = 4$.

Encode/decode **bad** pair (W, i) :

- $W' = W \cup S_i = \{1, 4, 5, 6, 7\}$ we find $j = 3$ with $S_j \subset W'$
- $\text{aux}_1 = \$$$$ with at least $w/2$ $s we know $S_j \cap S_i = \{4, 5, 6\}$$
- $k = 2$ S_i ranks 2 in $\mathcal{F}_{\{4, 5, 6\}}$, we recover $i = 4$
- $\text{aux}_2 = \$$$$$ we recover $W = W' \setminus \{4, 5, 6, 7\}$

Reductions

Let $\mathcal{F} = \{S_1, \dots, S_m\}$ be a w -(multi-)set system on $\{x_1, \dots, x_n\}$.
Assume $|\mathcal{F}_Y| < m/\kappa^{|Y|}$ holds for any Y , and $\kappa = (\log w)^{O(1)}$.

Reductions

Let $\mathcal{F} = \{S_1, \dots, S_m\}$ be a w -(multi-)set system on $\{x_1, \dots, x_n\}$.
Assume $|\mathcal{F}_Y| < m/\kappa^{|Y|}$ holds for any Y , and $\kappa = (\log w)^{O(1)}$.

It suffices to prove

- \mathcal{F} is satisfying
 \iff w.h.p S contains some set of \mathcal{F} , and $\Pr[x_i \in S] = 1/3$.

Reductions

Let $\mathcal{F} = \{S_1, \dots, S_m\}$ be a w -(multi-)set system on $\{x_1, \dots, x_n\}$.
Assume $|\mathcal{F}_Y| < m/\kappa^{|Y|}$ holds for any Y , and $\kappa = (\log w)^{O(1)}$.

It suffices to prove

- \mathcal{F} is satisfying
 \iff w.h.p S contains some set of \mathcal{F} , and $\Pr[x_i \in S] = 1/3$.

Split S to several steps,

- $\Pr[x_i \in S] = 1/3$
 \approx take $1/3$ -fraction of the ground set as S
 \approx view S as $W_1, W_2, \dots, W_{\log w}$, each of $\approx 1/\sqrt{\kappa}$ -fraction

Reductions

Let $\mathcal{F} = \{S_1, \dots, S_m\}$ be a w -(multi-)set system on $\{x_1, \dots, x_n\}$.
Assume $|\mathcal{F}_Y| < m/\kappa^{|Y|}$ holds for any Y , and $\kappa = (\log w)^{O(1)}$.

It suffices to prove

- \mathcal{F} is satisfying
 \iff w.h.p S contains some set of \mathcal{F} , and $\Pr[x_i \in S] = 1/3$.

Split S to several steps,

- $\Pr[x_i \in S] = 1/3$
 \approx take $1/3$ -fraction of the ground set as S
 \approx view S as $W_1, W_2, \dots, W_{\log w}$, each of $\approx 1/\sqrt{\kappa}$ -fraction

Then we iteratively apply reductions,

$$\mathcal{F} \xrightarrow{W_1} \mathcal{F}' \xrightarrow{W_2} \mathcal{F}'' \xrightarrow{W_3} \dots \xrightarrow{W_{\log w}} \mathcal{F}^{\text{last}}.$$

Final step

Recall $S = W_1 \cup \dots \cup W_{\log w}$ and

$\underbrace{\mathcal{F}}_{\text{width-}w}$

Final step

Recall $S = W_1 \cup \dots \cup W_{\log w}$ and

$$\underbrace{\mathcal{F}}_{\text{width-}w} \xrightarrow{W_1} \underbrace{\mathcal{F}'}_{\text{width-}w/2}$$

Final step

Recall $S = W_1 \cup \dots \cup W_{\log w}$ and

$$\underbrace{\mathcal{F}}_{\text{width-}w} \xrightarrow{W_1} \underbrace{\mathcal{F}'}_{\text{width-}w/2} \xrightarrow{W_2} \underbrace{\mathcal{F}''}_{\text{width-}w/4}$$

Final step

Recall $S = W_1 \cup \dots \cup W_{\log w}$ and

$$\underbrace{\mathcal{F}}_{\text{width-}w} \xrightarrow{W_1} \underbrace{\mathcal{F}'}_{\text{width-}w/2} \xrightarrow{W_2} \underbrace{\mathcal{F}''}_{\text{width-}w/4} \xrightarrow{W_3} \dots \xrightarrow{W_{\log w}} \underbrace{\mathcal{F}^{\text{last}}}_{\text{width-}0}.$$

Final step

Recall $S = W_1 \cup \dots \cup W_{\log w}$ and

$$\underbrace{\mathcal{F}}_{\text{width-}w} \xrightarrow{W_1} \underbrace{\mathcal{F}'}_{\text{width-}w/2} \xrightarrow{W_2} \underbrace{\mathcal{F}''}_{\text{width-}w/4} \xrightarrow{W_3} \dots \xrightarrow{W_{\log w}} \underbrace{\mathcal{F}^{\text{last}}}_{\text{width-}0}.$$

- either we stop at W_i when some set is contained in $\bigcup_{j < i} W_j$,
 $\Rightarrow S$ contains some set of \mathcal{F}

Final step

Recall $S = W_1 \cup \dots \cup W_{\log w}$ and

$$\underbrace{\mathcal{F}}_{\text{width-}w} \xrightarrow{W_1} \underbrace{\mathcal{F}'}_{\text{width-}w/2} \xrightarrow{W_2} \underbrace{\mathcal{F}''}_{\text{width-}w/4} \xrightarrow{W_3} \dots \xrightarrow{W_{\log w}} \underbrace{\mathcal{F}^{\text{last}}}_{\text{width-}0}.$$

- either we stop at W_i when some set is contained in $\bigcup_{j < i} W_j$,
 $\Rightarrow S$ contains some set of \mathcal{F}
- or, $\mathcal{F}^{\text{last}}$ is a width-0 (multi-)set system of size $\approx m > \kappa^w$,
 and $|\mathcal{F}_Y^{\text{last}}| \lesssim |\mathcal{F}^{\text{last}}| / \kappa^{|Y|}$ still holds for any Y .
 \Rightarrow Impossible

Final step

Recall $S = W_1 \cup \dots \cup W_{\log w}$ and

$$\underbrace{\mathcal{F}}_{\text{width-}w} \xrightarrow{W_1} \underbrace{\mathcal{F}'}_{\text{width-}w/2} \xrightarrow{W_2} \underbrace{\mathcal{F}''}_{\text{width-}w/4} \xrightarrow{W_3} \dots \xrightarrow{W_{\log w}} \underbrace{\mathcal{F}^{\text{last}}}_{\text{width-}0}.$$

- either we stop at W_i when some set is contained in $\bigcup_{j < i} W_j$,
 $\Rightarrow S$ contains some set of \mathcal{F}
- or, $\mathcal{F}^{\text{last}}$ is a width-0 (multi-)set system of size $\approx m > \kappa^w$,
 and $|\mathcal{F}_Y^{\text{last}}| \lesssim |\mathcal{F}^{\text{last}}| / \kappa^{|Y|}$ still holds for any Y .
 \Rightarrow Impossible

Thus, (informally) we proved such \mathcal{F} is satisfying, which means \mathcal{F} has 3-sunflower (3 pairwise disjoint sets).

Section 4

Open problems

Erdős-Rado sunflower

Problem (Erdős-Rado sunflower conjecture)

Any w -set system of size $O_r(1)^w$ has r -sunflower.

- Our approach cannot go beyond $(\log w)^{(1-o(1))w}$.
We need new ideas.
- Lift the sunflower size?
 $r = 3 \implies r = 4$.
- Is $(\log w)^{(1-o(1))w}$ actually tight? Counterexamples?

Erdős-Szemerédi sunflower

Assume $\mathcal{F} = \{S_1, \dots, S_m\}$ and $S_i \subset \{1, 2, \dots, n\}$.

Problem (Erdős-Szemerédi sunflower conjecture)

There exists function $\varepsilon = \varepsilon(r) > 0$, such that, if $m > 2^{n(1-\varepsilon)}$, then \mathcal{F} has r -sunflower.

- Now:
 - general r : $\varepsilon = O_r(1/\log n)$ from ER sunflower.
 - $r = 3$: Naslund proved it using polynomial method.
- ER sunflower conjecture \implies ES sunflower conjecture.

Section 5

Thanks