




Supplementary Fig. S10 | Receptor and transporter availability in the negative-profile regions, across 39 PET maps. Each row is one PET map from the neuromaps collection. The unit of observation is one Shen-268 region: the 32 regions plotted as the negative profile are the nodes of its 29 edges, a superset of the 19 nodes of the 12-edge NP factor, and are compared with the remaining 236 regions. x, Hedges' g for the difference in mean regional value (negative profile minus other regions); a filled marker marks a map that survives correction, an open one a map that does not; the map name is printed in black or grey to match. The test is a permutation test on the difference in means: the region labels are reassigned at random 10,000 times with the group sizes held at 32 and 236, and the two-sided P value is $(\#|null| \geq |observed| + 1) / (10,000 + 1)$. P values are corrected across all 39 maps with the Benjamini-Hochberg FDR, and 11 maps reach $q < 0.05$, all of them in the direction of higher availability in the negative-profile regions: 5-HT1a way savli ($g = 0.52$, $q = 0.030$), 5-HT2a alt savli ($g = 0.54$, $q = 0.030$), 5-HT2a cimbi beliveau ($g = 0.54$, $q = 0.030$), 5-HT2a mdl talbot ($g = 0.55$, $q = 0.030$), M1 lsn naganawa ($g = 0.57$, $q = 0.030$), mGluR5 abp rosaneto ($g = 0.54$, $q = 0.030$), mGluR5 abp dubois ($g = 0.58$, $q = 0.030$), mGluR5 abp smart ($g = 0.61$, $q = 0.030$), 5-HT1a cumi beliveau ($g = 0.50$, $q = 0.039$), CB1 FMPEPd2 laurikainen ($g = 0.49$, $q = 0.039$), 5-HT6 gsk radhakrishnan ($g = 0.49$, $q = 0.041$). Every map is shown, so the selection is visible: the effect sizes of the maps that do not survive run from -0.30 to +0.41. The region set carries the negative-profile colour rather than the NP-factor blue of the main figures because it is defined on the 29-edge profile; repeating the test on the 19 NP-factor nodes alone leaves 6 maps below $q = 0.05$.