

Maxpro 20 mg

Patient Information Leaflet — Esomeprazole 20 mg. Read this leaflet carefully before you start taking this medicine.

Brand name	Maxpro 20 mg
Active ingredient	Esomeprazole 20 mg
Form	Enteric-coated (delayed-release) tablet
Manufacturer	Renata PLC, Bangladesh
Pack & price	Unit price ₳7.00; strip of 14: ₳98.00; pack of 10 × 14: ₳980.00

1. What Maxpro is and what it is used for

Maxpro contains esomeprazole, a proton pump inhibitor that reduces the amount of acid produced in the stomach. It is used for:

- chronic heartburn and the symptoms of gastro-oesophageal reflux disease (GERD)
- healing and long-term maintenance of erosive oesophagitis
- duodenal and gastric ulcers
- eradication of *Helicobacter pylori* (together with amoxicillin and clarithromycin) in duodenal ulcer disease
- Zollinger-Ellison Syndrome and acid-related dyspepsia

2. Before you take Maxpro

Do not take if you are hypersensitive to esomeprazole or any component of the formulation.

- A good symptomatic response does not rule out the presence of a gastric malignancy; tell your doctor if symptoms persist.
- Do not chew or crush the tablet/pellets.

3. How to take Maxpro

Swallow the tablet whole with water, **one hour before a meal** — do not chew or crush it. If swallowing is difficult, the pellets may be mixed with cool apple sauce and taken immediately.

Condition	Usual dose
Symptomatic GERD	20 mg once daily for 4 weeks
Healing of erosive oesophagitis	20 or 40 mg once daily for 4–8 weeks
Duodenal ulcer	20 mg once daily for 2–4 weeks
<i>H. pylori</i> eradication	40 mg once daily plus antibiotics for 10 days
Zollinger-Ellison Syndrome	20–80 mg once daily

Antacids may be used at the same time if needed.

4. Possible side effects

Common: headache, diarrhoea, nausea, flatulence, abdominal pain, constipation, and dry mouth. The rate of side effects is similar for long-term (up to 12 months) and short-term use.

5. Use in pregnancy and breast-feeding

There are no adequate controlled studies in pregnant women, although animal studies showed no teratogenic effects. Excretion in breast milk has not been studied, so breast-feeding should be discontinued if treatment is considered essential. Ask your doctor for advice.

6. How to store Maxpro

Store at a temperature not exceeding 30 °C in a dry place. Protect from light and moisture. **Keep out of the reach of children.**

Source: product information from MedEx (medex.com.bd). This leaflet is reproduced for an illustrative technical assessment exercise only and is not medical advice.