

Rolac 10 mg

Patient Information Leaflet — Ketorolac Tromethamine 10 mg. Read this leaflet carefully before you start taking this medicine.

Brand name	Rolac 10 mg
Active ingredient	Ketorolac Tromethamine 10 mg
Form	Tablet
Manufacturer	Renata PLC, Bangladesh
Pack & price	Unit price ₳12.00; strip of 14: ₳168.00; pack of 4 × 14: ₳672.00

1. What Rolac is and what it is used for

Rolac contains ketorolac, a non-steroidal anti-inflammatory drug (NSAID). It is used for the **short-term management of moderate to severe acute post-operative pain**.

2. Before you take Rolac

Do not take if you: are hypersensitive to ketorolac or other NSAIDs; are under 16 years of age; or intend to use it as a preventive painkiller before surgery.

Use with caution, and tell your doctor, if you:

- are over 65 years of age
- have an active or suspected peptic ulcer or gastrointestinal bleeding
- have asthma or liver dysfunction

3. How to take Rolac

Patient	Dose
Adults	10 mg every 4–6 hours

- **For short-term use only — up to 7 days.** Not recommended for chronic use.
- **Do not exceed 40 mg per day.**

4. Possible side effects

Nausea, vomiting, gastrointestinal bleeding, black stools (melaena), peptic ulcer, pancreatitis, anxiety, drowsiness, headache, fatigue, slow heartbeat (bradycardia), high blood pressure, palpitations, chest pain, and pulmonary oedema. Stop taking and seek medical help if you notice signs of stomach bleeding or a serious reaction.

5. Use in pregnancy and breast-feeding

US FDA Pregnancy Category C. Should be avoided during pregnancy and breast-feeding unless the potential benefits outweigh the possible risks to the baby. Ask your doctor for advice.

6. How to store Rolac

Keep in a dry place, away from light and heat. **Keep out of the reach of children.**

Source: product information from MedEx (medex.com.bd). This leaflet is reproduced for an illustrative technical assessment exercise only and is not medical advice.