

Rolip 10 mg

Patient Information Leaflet — Rosuvastatin Calcium 10 mg. Read this leaflet carefully before you start taking this medicine.

Brand name	Rolip 10 mg
Active ingredient	Rosuvastatin (as Rosuvastatin Calcium) 10 mg
Form	Tablet
Manufacturer	Renata PLC, Bangladesh
Pack & price	Unit price ₳24.00; strip of 10: ₳240.00; pack of 3 × 10: ₳720.00

1. What Rolip is and what it is used for

Rolip contains rosuvastatin, a statin that lowers cholesterol. It is used, alongside a suitable diet, to treat lipid disorders — including primary hyperlipidaemia, mixed dyslipidaemia, hypertriglyceridaemia, Type III hyperlipoproteinaemia, and familial hypercholesterolaemia. It also helps slow the progression of atherosclerosis and reduce the risk of heart attack, stroke, and arterial procedures in high-risk patients, and is used for heterozygous familial hypercholesterolaemia in children aged 10–17 when diet alone is not enough.

2. Before you take Rolip

Do not take if you: are hypersensitive to any component; have active liver disease (including unexplained, persistently raised liver enzymes); are pregnant or may become pregnant; or are breast-feeding.

Tell your doctor before taking if you have factors that increase the risk of muscle problems (myopathy) — being over 65, an under-treated underactive thyroid, or kidney impairment. Stop and contact your doctor if you develop unexplained muscle pain, tenderness, or weakness. Your doctor will check your liver with blood tests before treatment, at 12 weeks, and periodically thereafter.

3. How to take Rolip

Can be taken **with or without food, at any time of day.**

Patient	Dose
Usual range (adults)	5–40 mg once daily
Very high cholesterol / high risk	up to 40 mg once daily
Homozygous familial hypercholesterolaemia	start 20 mg once daily
Asian patients	consider starting at 5 mg once daily
Severe renal impairment (not on dialysis)	

Patient	Dose
	start at 5 mg, not exceeding 10 mg daily

If you also take cyclosporine, limit to 5 mg daily; with certain protease-inhibitor combinations, limit to 10 mg daily.

4. Possible side effects

Generally well tolerated. Common: headache, muscle pain (myalgia), constipation, weakness (asthenia), abdominal pain, and nausea. Serious muscle effects (myopathy/rhabdomyolysis) are rare — see the precautions above.

5. Use in pregnancy and breast-feeding

Give to women who could become pregnant only when conception is highly unlikely and the hazards have been explained; stop immediately if pregnancy occurs. It is contraindicated in nursing mothers.

6. How to store Rolip

Keep below 30 °C, protected from light and moisture. **Keep out of the reach of children.**

Source: product information from MedEx (medex.com.bd). This leaflet is reproduced for an illustrative technical assessment exercise only and is not medical advice.