



**BALTORGANIC**

LIQUID HUMIC  
CONCENTRATE

**BALTORGANIC**

based on lowland peat

2026



# ABOUT THE COMPANY



The **BALTORGANIC** group of companies implements projects related to the production of original complex organic and organo-mineral fertilizers based on animal and plant raw materials.

Production facilities are located in the Leningrad Region and the Republic of Mari El.



## Product range



Organic fertilizers  
in granules and pellets



Organo-mineral fertilizers



Liquid humic fertilizers

# STRESS DEPLETES YOUR HARVEST



Modern agriculture faces a number of factors that reduce the yield potential of crops.

## Main causes of harvest loss:



Soil degradation and depletion



Decline in natural fertility



Climatic stresses  
droughts and return frosts



Impaired nutrient uptake by plants

Under such conditions, plants require additional support for full growth, development and yield formation.



# SOLUTION — **BALTORGANIC** HUMIC CONCENTRATE

Liquid concentrates based on ecologically clean lowland peat.  
4 products for any agronomic challenge.

## Base foundation



HA/FA  
Humic and fulvic acids.



100% organic  
Free from coal humate impurities and synthetics.



### **Concentrate 30+%** **Ultra-base**

Maximum concentration  
of humic acids for powerful  
stimulation and soil restoration



### **Concentrate 16-18%** **Base**

Optimal concentration  
for regular support  
and plant stress relief



### **Concentrate 16-18%** **+ NPK**

Enriched with a macronutrient  
complex (N, P, K, Mg, S)  
for intensive nutrition during  
the growing season



### **Concentrate 16-18%** **+ poultry manure**

Enhanced with natural organics  
(Ca, Fe + microelements)  
to support weakened crops





# SCIENTIFIC APPROACH: ACTION OF HUMIC SUBSTANCES

Molecules of humic substances and fulvic acids can penetrate the plant cell already at the seed germination stage, transmitting growth energy to the plant.

This promotes:



Accelerated root system development



Activation of plant respiration



More intensive growth  
at early development stages



Plants receive additional support during the most vulnerable period — the initial growth stage.

# SUPPORT FROM SEED TO FRUIT



Humic substance molecules continue to actively work throughout the entire plant life cycle.



Active action from seedling emergence to harvest




Promotes restoration of soil structure and microflora



Acts as a soil ameliorant



Retains moisture in the root zone





# MAXIMUM PROFIT, MINIMUM COST

Application of **BALTORGANIC** humic concentrate improves the economic efficiency of crop cultivation.

## Yield increase:



For grain  
crops

up to

**10–30%**



For industrial  
crops

up to

**40–50%**

Result — increased output  
of high-quality marketable products.



Pesticide cost  
reduction

by **30–70%**



Mineral fertilizer  
consumption reduction

by **15–20%**

# ECOLOGY AND PRODUCT SAFETY



Sharp **reduction**  
in **nitrate** levels in fruits



**Increase** in sugar,  
protein and vitamin content



Effective binding  
and neutralization of heavy metals



**Improved crop**  
**shelf life during storage**



# THREE STEPS TO A RICH HARVEST



## STAGE 1

Pre-sowing seed  
and tuber treatment



increases germination energy  
and ensures a powerful start  
to plant development.

## STAGE 2

Soil treatment  
before planting



improves soil structure  
and activates  
microbiological processes.

## STAGE 3

Foliar spraying  
and root-zone treatment  
during the growing season



supports active plant growth  
and yield formation.

# PREPARATION AND COMPATIBILITY



The product easily integrates into existing crop cultivation technologies.



Compatible with **UAN, mineral fertilizers and crop protection products**

Can be used in tank mixes

Reduces chemical stress on plants when applying agrochemicals

Application rate for pre-sowing soil treatment:

**10–35 l/ha** | depending on soil type

For restoration of depleted and degraded lands:

**30–40 l/ha**

# WHEAT | RYE | BARLEY | OATS



## 1 SEED TREATMENT

Soaking seeds before planting together with a seed dresser.



**Working solution:**  
200–300 ml  
per 10 l of water  
for 100 kg of seeds.

## 2 VEGETATION PERIOD TREATMENT (SPRAYING)

3 applications:

1st — tillering stage

2nd — end of stem elongation stage

3rd — grain filling stage



**Working solution:**  
2–3 l per 1 ha

## 3 ROOT-ZONE SOIL TREATMENT


2 applications:

1st — emergence of 1–2 leaves

2nd — grain filling stage



**Consumption:**  
18–20 l per 1 ha



# RICE



## 1 SEED TREATMENT

Soaking seeds together with a seed dresser.



**Working solution:**  
300–400 ml  
per 10 l of water  
for 100 kg of seeds.

## 2 VEGETATION PERIOD TREATMENT (SPRAYING)

2 applications:

**1st** —  
tillering to early stem elongation stage

**2nd** — every 10–15 days



**Working solution:**  
2–3 l  
in 150–200 l of water  
per 1 ha

## 3 ROOT-ZONE SOIL TREATMENT


2 applications:

**1st** — tillering stage

**2nd** — shoot formation stage



**Consumption:**  
20–30 l per 1 ha



# SOYBEAN | RAPESEED | LENTIL | BEAN



## 1 SEED TREATMENT

Soaking seeds together with a seed dresser.



**Working solution:**  
200–300 ml  
per 10 l of water  
for 100 kg of seeds

## 2 VEGETATION PERIOD TREATMENT (SPRAYING)

3 applications:

**1st** — 1–2 leaf pairs emergence

**2nd** — every 10–15 days

**3rd** — grain filling stage



**Working solution:**  
2–3 l/1 ha

## 3 ROOT-ZONE SOIL TREATMENT


2 applications:

**1st** — during seedling emergence

**2nd** — pod formation stage



**Consumption:**  
20–25 l per 1 ha





# PEAS

## 1 SEED TREATMENT

Soaking seeds together with a seed dresser.



**Working solution:**  
250–350 ml  
per 10 l of water  
for 100 kg of seeds

## 2 VEGETATION PERIOD TREATMENT (SPRAYING)

3 applications:

**1st** — seed-to-leaf stage

**2nd** — stem elongation to 2–3 true leaves

**3rd** — budding stage



**Working solution:**  
2–3 l/1 ha

## 3 ROOT-ZONE SOIL TREATMENT


2 applications:

**1st** — during seedling emergence

**2nd** — pod formation stage



**Consumption:**  
20–25 l per 1 ha





# SUNFLOWER

## 1 SEED TREATMENT

Soaking seeds together with a seed dresser.



**Working solution:**  
200–300 ml  
per 10 l of water  
for 100 kg of seeds

## 2 VEGETATION PERIOD TREATMENT (SPRAYING)

3 applications:

1st — seedling emergence

2nd — 3–4 pairs of true leaves

3rd — after 10–15 days



**Working solution:**  
2–3 l/1 ha

## 3 ROOT-ZONE SOIL TREATMENT

2 applications:


1st — 2–4 leaf pairs emergence

2nd — head formation

3rd — grain filling stage



**Consumption:**  
20–25 l per 1 ha



# COTTON



## 1 SEED TREATMENT

Soaking seeds together with a seed dresser.



**Working solution:**  
400–500 ml  
per 10 l of water  
for 1,000 kg of seeds

## 2 VEGETATION PERIOD TREATMENT (SPRAYING)

3 applications:

**1st** — 2–3 leaves emergence

**2nd** — bud formation

**3rd** — after 12–15 days



**Working solution:**  
2–3 l in 200–300 l  
per 1 ha

## 3 ROOT-ZONE SOIL TREATMENT






2 applications:

**1st** — 3–5 leaves emergence

**2nd** — after 12–15 days



**Consumption:**  
20–30 l per 1 ha

|                                         | 7-12<br>Seed<br>germination                                                         | 14-15<br>Appearance<br>of 2–3 leaves                                                | 55-65<br>Bud formation                                                               | 70-75<br>Flowering                                                                    | 120-145<br>Boll opening                                                               | 150-190<br>Full ripeness                                                              |
|-----------------------------------------|-------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------|
| Seed<br>treatment<br>before<br>planting |  |  |  |  |  |  |
|                                         |                                                                                     | 1st application                                                                     | 2nd application                                                                      | 3rd application                                                                       |                                                                                       |                                                                                       |



# SUGAR BEET

## 1 SEED TREATMENT

Soaking seeds together with a seed dresser.



**Working solution:**  
300–400 ml  
per 10 l of water  
for 100 kg of seeds

## 2 VEGETATION PERIOD TREATMENT (SPRAYING)

3 applications:

**1st** — formation of 2–3 pairs of true leaves

**2nd** — 4 leaf pairs to row closure

**3rd** — after 12–15 days



**Working solution:**  
2–3 l/1 ha

## 3 ROOT-ZONE SOIL TREATMENT


2 applications:

**1st** — during 2nd irrigation

**2nd** — during 3rd irrigation



**Consumption:**  
20–30 l per 1 ha



# CORN



## 1 SEED TREATMENT

Soaking seeds together with a seed dresser.



**Working solution:**  
200–300 ml  
per 10 l of water  
for 100 kg of seeds

## 2 VEGETATION PERIOD TREATMENT (SPRAYING)

2 applications:

**1st** — emergence to 3–5 leaves

**2nd** — tasseling stage



**Working solution:**  
2–3 l/1 ha

## 3 ROOT-ZONE SOIL TREATMENT


2 applications:

**1st** — 3–5 leaves emergence

**2nd** — shoot formation stage



**Consumption:**  
25–35 l per 1 ha





# TOMATOES AND CUCUMBERS

## 1 SEED TREATMENT

Soaking seeds together with a seed dresser.



**Working solution:**  
200–300 ml  
per 10 l of water  
for 100 kg of seeds

## 2 VEGETATION PERIOD TREATMENT (SPRAYING)

3 applications:

**1st** — 2–4 leaves emergence

**2nd** — at 15-day intervals



**Working solution:**  
2–3 l per 1 ha

## 3 ROOT-ZONE SOIL TREATMENT

3 applications:


**1st** — 3–5 leaves emergence

**2nd** — flowering stage

**3rd** — fruit formation



**Consumption:**  
20–30 l per 1 ha

|                                                                                                                                          | Appearance<br>of 2–4 leaves                                                                                   | Appearance<br>of 5–7 leaves                                                         | Ovary formation                                                                      | Flowering                                                                                                       | Fruit formation                                                                                                 | Fruit ripening                                                                        | Harvest                                                                               |
|------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------|
| <b>Seed<br/>treatment<br/>before<br/>planting</b><br> | <b>1st application</b><br> |  |  | <b>2nd application</b><br> | <br><b>3rd application</b> |  |  |

# POTATO



## 1 TUBER TREATMENT

Soaking tubers before planting.



**Working solution:**  
400–500 ml  
per 10 l of water  
for 100 kg of tubers

## 2 VEGETATION PERIOD TREATMENT (SPRAYING)

2 applications:

**1st** — 5–7 leaves emergence

**2nd** — budding stage



**Working solution:**  
2–3 l per 1 ha

## 3 ROOT-ZONE SOIL TREATMENT

2 applications:

**1st** — during 2nd irrigation

**2nd** — during 3rd irrigation



**Consumption:**  
20–30 l per 1 ha

|                                | Sowing                                                                              | Seedling emergence                                                                  | Appearance of 5–7 leaves                                                             | Budding phase                                                                         | Flowering                                                                             | Ripening                                                                              | Wilting                                                                               |
|--------------------------------|-------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------|
| Seed treatment before planting |  |  |  |  |  |  |  |

# ONION



## 1 BULB TREATMENT

Soaking bulbs before planting for 3 hours.



**Working solution:**  
300 ml  
per 12 l of water  
for 100 kg of bulbs

## 2 VEGETATION PERIOD TREATMENT (SPRAYING)

3 applications:

**1st** — 2–3 leaves emergence

**2 subsequent** at 10–15 day intervals



**Working solution:**  
2–3 l per 1 ha

## 3 ROOT-ZONE SOIL TREATMENT


3 applications:

**1st** — 2–5 leaves emergence

**2 subsequent** at 10–15 day intervals



**Consumption:**  
20–30 l per 1 ha





# ФРУКТЫ И ЯГОДЫ

## 1 CUTTING AND SEEDLING TREATMENT

Treatment together with fertilizers and biopreparations.  
Soaking cuttings for 24 hours.



**Consumption:**  
18–20 l  
per 100 cuttings

## 2 VEGETATION PERIOD TREATMENT (SPRAYING)

4 applications:

1st — 5–7 days after flowering

2nd — onset of physiological fruit drop

3rd — flower bud formation period

4th — intensive fruit growth period



**Working solution:**  
2–3 l in 100 l of water  
per 1 ha

## 3 ROOT-ZONE SOIL TREATMENT


5 applications:

First irrigation — during leaf formation

Last — during intensive fruit growth



**Consumption:**  
20–40 l  
per 100 l of water

|                                | Leaf formation                                                                      | Shoot formation                                                                     | Beginning of flowering                                                               | Flowering                                                                             | Petal drop                                                                            | Appearance of 2nd fruit                                                               | Leaf develop                                                                          | Beginning of ripening                                                                 |
|--------------------------------|-------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------|
| Seed treatment before planting |  |  |  |  |  |  |  |  |

1st application

2nd application

3rd application

4th application



# APPLICATION IN MODERN GROWING SYSTEMS

**BALTORGANIC** liquid humic concentrate is effectively used not only in traditional farming but also in modern plant growing technologies.

Use of humic concentrate allows:



Improved root system development



Increased plant stress resistance



Accelerated growth and yield formation



Improved efficiency of nutrient solutions

The product is suitable for various systems:

## Protected cultivation

Greenhouse complexes, nurseries



## Hydroponics

Cultivation on nutrient solutions



## Aeroponics

Intensive soilless technologies



# CONTACT US



**BALTORGANIC**

## **Sales Department**

8 Dobrolyubova Ave.,  
St. Petersburg

## **Production facilities**

Pervomaiskoye,  
Leningrad Region

Ishimovo,  
Republic of Mari El

Kingisepp,  
Leningrad Region



 +7 (921) 910-03-00

 [export@baltorganic.com](mailto:export@baltorganic.com)

 [baltorganic.com](http://baltorganic.com)

