

Integrative Structure Validation Report ?

January 05, 2026 - 12:15 PM PST

The following software was used in the production of this report:

IHMValidation Version 3.0

Python-IHM Version 2.7

MolProbity Version 4.5.2

PDB ID	8ZZF pdb_00008zzf
PDB-Dev ID	PDBDEV_00000015
Structure Title	Structure of human mitochondrial iron sulfur cluster core complex (NIAUF)2
Structure Authors	Cai, K.; Frederick, R.O.; Dashti, H.; Markley, J.L.
Deposited on	2018-01-24

This is a PDB-IHM Structure Validation Report.

We welcome your comments at helpdesk@pdb-ihm.org

A user guide is available at https://pdb-ihm.org/validation_help.html with specific help available everywhere you see the ? symbol.

List of references used to build this report is available [here](#).

1. Overview ?

1.1. Summary ?


This entry consists of 1 model(s). A total of 5 dataset(s) were used to build this entry.

Name	Type	Count
Crosslinking-MS data	Experimental data	2
NMR data	Experimental data	1
Experimental model	Starting model	2


1.2. Overall quality ?

This validation report contains model quality assessments for all structures, data quality and fit to model assessments for SAS and crosslinking-MS datasets. Data quality and fit to model assessments for other datasets and model uncertainty are under development. Number of plots is limited to 256.

Model Quality: Excluded Volume Analysis ?



Fit to Data Used for Modeling ?



2. Model Details ?

2.1. Ensemble information ?

This entry consists of 0 distinct ensemble(s).

2.2. Representation ?

This entry has 1 representation(s).

ID	Model(s)	Entity ID	Molecule name	Chain(s) [auth]	Total residues	Rigid segments	Flexible segments	Model coverage/ Starting model coverage (%)	Scale
1	1	1	NFS1	A	406	-	3-401	98.28 / 100.00	Atomic
		2	ISD11	B	91	-	5-85	89.01 / 100.00	Atomic
		3	Acp	C	77	-	4-74	92.21 / 100.00	Atomic
		4	ISCU	D	150	-	6-133	85.33 / 100.00	Atomic
		1	NFS1	E	406	-	3-403	98.77 / 100.00	Atomic
		2	ISD11	F	91	-	3-85	91.21 / 100.00	Atomic

ID	Model(s)	Entity ID	Molecule name	Chain(s) [auth]	Total residues	Rigid segments	Flexible segments	Model coverage/ Starting model coverage (%)	Scale
		3	Acp	G	77	-	3-72	90.91 / 100.00	Atomic
		4	ISCU	H	150	-	10-133	82.67 / 100.00	Atomic
		5	FXN	I	119	-	1-119	100.00 / 100.00	Atomic
				J					
		7	S-[2-({N-[(2R)-2-hydroxy-3,3-dimethyl-4-(phosphonooxy)butanoyl]-beta-alanyl]amino)ethyl] dodecanethioate	L [C]	Non-polymeric	-	-	Not available / Not available	Atomic
				O [G]					
		8	ZINC ION	M [D]	Non-polymeric	-	-	Not available / Not available	Atomic
				P [H]					

2.3. Datasets used for modeling ?

There are 5 unique datasets used to build the models in this entry.

ID	Dataset type	Database name	Data access code
1	Experimental model	PDB	pdb_00005wlv
2	Experimental model	PDB	pdb_00001ekg
3	NMR data	BMRB	27171
4	Crosslinking-MS data	PRIDE	PXD006938
5	Crosslinking-MS data	PRIDE	PXD006928

2.4. Methodology and software ?

This entry is a result of 1 distinct protocol(s).

Step number	Protocol ID	Method name	Method type	Method description	Number of computed models	Multi state modeling	Multi scale modeling
1	1	Not available	Not available	Not available	Not available	False	False

There is 1 software package reported in this entry.

ID	Software name	Software version	Software classification	Software location
1	HADDOCK	2.20	molecular docking	http://haddock.science.uu.nl/services/HADDOCK/

3. Data quality ?

3.2. Crosslinking-MS

At the moment, data validation is only available for crosslinking-MS data deposited as a fully [compliant](#) dataset in the [PRIDE Crosslinking](#) database. Correspondence between crosslinking-MS and entry entities is established using [pyHMMER](#). Only residue pairs that passed the reported threshold are used for the analysis. The values in the report have to be interpreted in the context of the experiment (i.e. only a minor fraction of in-situ or in-vivo dataset can be used for modeling).

Crosslinking-MS dataset is not available in the [PRIDE Crosslinking](#) database.

3.4. NMR ?

Validation for this section is under development.

4. Model quality ?

For models with atomic structures, MolProbity analysis is performed. For models with coarse-grained or multi-scale structures, excluded volume analysis is performed.

4.1a. Excluded Volume Analysis ?

Excluded volume satisfaction for the models in the entry are listed below. The Analysed column shows the number of particle-particle or particle-atom pairs for which excluded volume was analysed.

Model ID	Analysed	Number of violations	Excluded Volume Satisfaction (%)
1	73732296	43684	99.94

5. Fit to Data Used for Modeling Assessment ?

5.2. Crosslinking-MS ?

5.2.1. Restraint types ?

This table summarizes information about crosslinker(s) used for data generation, and how crosslinking information was translated into actual modeling restraints. Restraints assigned "by-residue" are interpreted as between CA atoms. Restraints between coarse-grained beads are indicated as "coarse-grained". [Restraint group](#) represents a set of crosslinking restraints applied collectively in the modeling.


There are 16 crosslinking restraints combined in 8 restraint groups.

Linker	Residue 1	Atom 1	Residue 2	Atom 2	Restraint type	Distance, Å	Count
sulfo-SMCC	CYS	CA	LYS	CA	upper bound	22.00	16

Distograms of individual restraints

Distograms (i.e., histogram plots of distances) provide an overview of distributions of distances between residues for which chemical crosslinks were identified. The shift of the distogram relative to the threshold value may indicate a poor model. Restraints with identical thresholds are grouped into one plot. Only the best distance per restraint per model

group/ensemble is plotted. Inter- and intramolecular (including self-links) restraints are also grouped into one plot. Distance for a restraint between coarse-grained beads is calculated as a minimal distance between shells; if beads intersect, the distance will be reported as 0.0. A bead with the highest available resolution for a given residue is used for the assessment.




5.2.2. Satisfaction of restraints ?

Satisfaction of restraints is calculated on a [restraint group](#) (a set of crosslinking restraints applied collectively in the modeling) level. Satisfaction of a restraint group depends on satisfaction of individual restraints in the group and the conditionality (all/any). A restraint group is considered satisfied, if the condition was met in at least one model of the model group/ensemble. The number of measured restraints can be smaller than the total number of restraint groups if crosslinks involve non-modeled residues. Only deposited models are used for validation right now.

State group	State	Model group	# of Deposited models/Total	Restraint group type	Satisfied (%)	Violated (%)	Count (Total=8)
1	1	1	1/1	All	75.00	25.00	8
				Heteromeric links/Intermolecular	75.00	25.00	8

Per-model satisfaction rates in ensembles

Every point represents one model in a model group/ensemble. Where possible, boxplots with quartile marks are also plotted.



5.4. NMR ?

Validation for this section is under development.

6. Fit to Data Used for Validation Assessment ?

Validation for this section is under development.

Acknowledgments

The development of integrative model validation metrics, implementation of a model validation pipeline, and creation of a validation report for integrative structures are funded by NSF awards to the [PDB-IHM team](#) (DBI-1756248, DBI-2112966, DBI-2112967, DBI-2112968, and DBI-1756250) and awards from NSF, NIH, and DOE to the [RCSB PDB](#) (DBI-2321666, R01GM157729, and DE-SC0019749). The PDB-IHM team and members of the [Sali lab](#) contributed model validation metrics and software packages.

Dr. Jill Trehwella, Dr. Dina Schneidman, and members of the [SASBDB](#) repository are acknowledged for their advice and support in implementing SAS validation methods. Team members from the labs of Dr. Juri Rappsilber, Dr. Alexander Leitner, Dr. Andrea Graziadei, and members of [PRIDE](#) database are acknowledged for their advice and support in implementing crosslinking-MS validation methods. We are grateful to Dr. Shruthi Viswanath for discussions about uncertainty assessment of integrative structural models.

Members of the [wwPDB Integrative/Hybrid Methods Task Force](#) provided recommendations and community support for the project.