Competitive Landscape: non-small cell lung cancer (US, 2023)

Report date: 2023-10-02 | Prepared by: Omar Faruk

Market Overview

In the US non-small cell lung cancer market, 4 brands compete for first-line+ patients as of Q4 2023. The segment is shaped by guideline updates, reimbursement decisions and real-world evidence.

The number of branded options in non-small cell lung cancer has grown since 2020, increasing pressure on standard of care positioning.

Positioning: Keytrava

In the US non-small cell lung cancer market, Keytrava maintains a 27% market share as of Q3 2023, with the strongest uptake among oncologists.

Sentiment index for Keytrava in US: 0.68 (scale 0 to 1, 2023). Clinicians view the brand strongly positively; payer prior-authorisation requirements slow initiation.

Positioning: Opdivra

In the US non-small cell lung cancer market, Opdivra maintains a 37% market share as of Q1 2023, with the strongest uptake among oncologists.

Sentiment index for Opdivra in US: 0.30 (scale 0 to 1, 2023). Clinicians view the brand negatively; the dosing schedule complicates adherence.

Positioning: Tagrisa

In the US non-small cell lung cancer market, Tagrisa maintains a 9% market share as of Q1 2023, with the strongest uptake among oncologists.

Sentiment index for Tagrisa in US: 0.77 (scale 0 to 1, 2023). Clinicians view the brand strongly positively; the dosing schedule complicates adherence.

Positioning: Lumakro

In the US non-small cell lung cancer market, Lumakro maintains a 11% market share as of Q4 2023, with the strongest uptake among oncologists.

Sentiment index for Lumakro in US: 0.54 (scale 0 to 1, 2023). Clinicians view the brand moderately positively; cost and formulary access remain barriers.

Competitive Dynamics

Lumakro has been gaining ground since its 2021 approval for non-small cell lung cancer, pressuring Keytrava in first-line use.

Price competition, biosimilar entry and head-to-head evidence are expected to reshape formulary position over the next 24 months.

Outlook

We expect continued consolidation in non-small cell lung cancer. Differentiation will depend on tolerability, dosing convenience and evidence of overall survival or clinical benefit.

Synthetic document generated for the Watson prototype. All brand names and figures are fictional.