

# !! Sequence & Series !!

A.P. (Arithmetic Progression)

G.P. (Geometric Progression)

H.P. (Harmonic Progression)

A.G.P. (Arithmetic Geometric progression)

Special Sequences - 4 types

Sequence - ~~Seq~~ Succession of Terms which may be ~~all~~ algebraic, real & complex numbers written according to a definite rule.

Eg. 2, 3, 5, 7, 11 (Prime nos.)

4, 8, 12, 16, ... (Multiples of 4)

0, 0, 0, 0, ... (Multiples of 0)

→ Minimum no. of required terms of a sequence is 3.

Progression - Special case of sequence in which we can express the  $n^{\text{th}}$  term mathematically.

Eg. 2, 4, 6, 8, 10, ...  $a_n = 2n$

1, 2, 3, 4, 5, ...  $a_n = n$

Series - If we put sign of addition or subtraction between the terms of sequence then it is called series.

Eg.  $2 + 3 + 5 + 7 + 11 + \dots$

$0 + 7 + 26 + \dots$

$1 - 3 + 9 - 27 + \dots$

Q find out the 2<sup>nd</sup> & 4<sup>th</sup> terms of the sequence.

$$T_n / a_n = \left\{ \frac{-1}{n} \right\}^n$$

$$a_1 = (-1)^1 \Rightarrow a_1 = \left\{ \frac{-1}{1} \right\}^2 \quad a_4 = \left\{ \frac{-1}{4} \right\}^4$$
$$a_1 = -1 \quad a_4 = \frac{1}{256}$$
$$a_2 = \frac{1}{4}$$

Q write sequence where  $n^{th}$  terms are

①  $2^n$

OTTOE  
ARACTAIC  
MECHANICAL

②  $\log_a (nx)$

① 2, 4, 8, 16, 32, ...

②  $\log_a x, \log_a 2 + \log_a x, \log_a 3 + \log_a x, \dots$

~~sigma & pi notation~~

$$\sum_{i=0}^{10} x_i = x_1 + x_2 + x_3 + \dots + x_9 + x_{10}$$

$$\prod_{i=1}^{10} x_i = x_1 \cdot x_2 \cdot x_3 \cdot x_4 \cdot \dots \cdot x_9 \cdot x_{10} \quad (\text{Multiplication})$$

$$\sum_{i=1}^5 (ax_i + b) = (ax_1 + b) + (ax_2 + b) + (ax_3 + b) + (ax_4 + b) + (ax_5 + b)$$

$$\prod_{i=1}^5 (ax_i + b) = (ax_1 + b)(ax_2 + b)(ax_3 + b)(ax_4 + b)(ax_5 + b)$$

## Arithmetic Progression (A.P.)

→ Sequence Having two consecutive terms cross the constant difference.

$$\rightarrow a, a+nd, a+2d, a+3d \dots \overset{a+nd}{\downarrow}$$

$T_1 \quad T_2 \quad T_3 \quad T_4$

$T_{mn} = a + (m-1)d$

$a = \text{first term} \quad d = \text{common difference}$

Common Difference (C.D) can be +ve, -ve or zero

$$CD > 0$$

Increasing AP

$$CD < 0$$

Decreasing AP

$$CD = 0$$

Constant AP

Q1. If 6<sup>th</sup> term & 11<sup>th</sup> term of AP are 17 & 32 respectively  
find the 20<sup>th</sup> term.

$$\text{Q2. In an AP } T_2 + T_5 - T_3 = 10$$

$$T_2 + T_9 = 17$$

find  $T_1$  & CD.

Q3. If P<sup>th</sup>, Q<sup>th</sup> & R<sup>th</sup> Terms of an AP are a, b, c respectively  
then find  $a(a-a) + b(R-P) + c(P-a)$

Q4. If 11 times the 11<sup>th</sup> term of an AP is 9 times the  
9<sup>th</sup> term then find 20<sup>th</sup> term.

$$\text{Q1. } -17 = -a + 5d$$

$$32 = a + 10d$$

$$15 = 5d$$

$$d = 3$$

$$\begin{cases} a = 17 - 15 \\ a = 2 \end{cases}$$

$$T_{20} = 2 + 19 \times 3$$

$$= 59$$

Q2.

$$20 + 6d = 120$$

$$20 + 9d = 117$$

$$-3d = 3$$

$$\boxed{d = -1}$$

$$a + 3(-1) = 120$$

$$a - 3 = 120$$

$$\boxed{a = 123}$$

OTTOBLS

Q4.

$$11(a + 10d) = 9(0 + 8d)$$

$$11a + 110d = 9a + 72d$$

$$20 + 38d = 0$$

$$a + 19d = 0$$

$$+19d = 0$$

$$\boxed{T_{1020} = a + 19d = 0}$$

$$\begin{aligned}T_{20} &= a + 19d \\&= -17d + 19d \\&= 2d\end{aligned}$$

$$aQ = AQ + PdQ - dQ \quad aR = AR + PdR - dR$$

Q3.

$$a = A + (h-1)d$$

$$b = A + (e-1)d$$

$$c = A + (z-1)d$$

$$\begin{aligned}AQ + PdQ - dQ - A/R - PdR + dR + AR - AP + QdR - dR \\- QdP + dP - AQ + dQ + PdR - dR\end{aligned}$$

$$\boxed{= 0}$$

## Some Facts

→  $k^{\text{th}}$  term from the end of an AP

$$T_k = l + (k-1)(-d)$$

$$n, (n-1), (n-2) \dots 3, 2, 1 \Rightarrow -d$$

↓

$l$

→ If  $m^{\text{th}}$  term is  $n$  &  $n^{\text{th}}$  term is  $m$  then  $(m+n)^{\text{th}}$  term = 0

→ Sum of  $k^{\text{th}}$  term from beginning &  $k^{\text{th}}$  term from end is always constant

OTTOBLS  
ARACTA  
MUSICAL INSTRUMENTS

OLD

20-6-2024

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88

89

90

91

92

93

94

95

96

97

98

99

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

203

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

228

229

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

## Summation of an A.P.

$$S_n = a + (a+d) + (a+2d) + (a+3d) + \dots + (a+(n-2)d) + (a+(n-1)d)$$

$$+ S_n = a + (n-1)d + (a+d) + (a+2d) + (a+3d) + \dots + (a+(n-2)d) + (a+(n-1)d)$$

$$2S_n = 2a + (n-1)d + 2a + (n-1)d + 2a + (n-1)d + \dots + 2a + (n-1)d$$

$$= n(2a + (n-1)d)$$

$$\boxed{S_n = \frac{n}{2}(2a + (n-1)d)}$$

$$S_m = \frac{m}{2} [a + a + (m-1)d]$$

$$\boxed{S_m = \frac{m}{2}(a + T_m)}$$

$$S_m = \frac{m}{2} (\text{first term} + \text{last term})$$

Q If  $m$  in an A.P.  $T_m = \frac{1}{m}$  &  $T_{m-n} = \frac{1}{m-n}$  show that  $S_{mn} = \frac{1}{2}(mn+1)$

~~$T_n = a + (n-1)d$~~

~~$T_m = a + (m-1)d$~~

~~$S_{mn} = \frac{mn}{2}(2a + (mn-1)d)$~~

~~$a + (n-1)d = \frac{1}{m}$~~

~~$a + (m-1)d = \frac{1}{m-n}$~~

~~$a + (m-1)d = \frac{1}{m}$~~

~~$(n-1)d - (m-1)d = \frac{1}{m} - \frac{1}{m-n}$~~

~~$d(n-1) - d(m-1) = \frac{n-m}{mn}$~~

~~$d(n-m) = \frac{n-m}{mn}$~~

$$\boxed{d = \frac{1}{mn}}$$

~~$a + (m-1)d = \frac{1}{m}$~~

~~$a + (m-n)d = \frac{1}{m-n}$~~

~~$a + (mn-n)d = 1$~~

~~$a = \frac{1}{m} - nd + d$~~

~~$= \frac{1}{m} - \frac{1}{m} + \frac{1}{mn}$~~

~~$a = \frac{1}{mn}$~~

$$a = \frac{1}{mn}$$

$$S_m = \frac{m}{2} (2a + (m-1)d)$$

$$= \frac{m}{2} \left( 2 \times \frac{1}{m} + \frac{m \times 1}{m} - \frac{1}{m} \right)$$

$$= \frac{m}{2} \left( \frac{2}{m} - \frac{1}{m} + 1 \right)$$

$$= \frac{m}{2} \left( \frac{1+m}{m} \right)$$

$$\boxed{= \frac{1+m}{2}}$$

OTTOSLS

Q The sum of  $m$  terms of 2APs are in the ratio of  $7m+1 : 4m+27$ .  
Find the ratio of all 11 terms.

~~$\frac{7(1) + 1}{4(1) + 27}$~~

~~$S_{11} = 78 = \frac{n}{2} (2a + (n-1)d)$~~

~~$a_1 = 7$~~

~~$S_{11} = 71 = 67 + a_{10}$~~

~~$a_{10} = 4$~~

~~$\frac{a_{11}}{a_{10}} = \frac{7}{4}$~~

~~$\frac{a_1 + 10d}{a_1 + 9d}$~~

$$\frac{7m+1}{4m+27} = \frac{a + \frac{(m-1)d}{2}}{a + \frac{(m-1)d}{2}}$$

$$\text{for } \frac{m-1}{2} = 10$$

$$m = 21$$

$$\frac{S_m}{S_{11}} = \frac{\frac{m}{2} (2a + (m-1)d)}{\frac{11}{2} (2a + (m-1)d)}$$

$$= \frac{m}{11} a + \frac{(m-1)d}{2}$$

$$= \frac{a + \frac{(m-1)d}{2}}{a + \frac{(m-1)d}{2}}$$

$$\frac{148}{111} = \sqrt{\frac{4}{3}} = \frac{T_{11}}{T_{11}}$$

Note  $\rightarrow$  If the sum of an A.P ~~is~~  $S_n$  is given

$T_n = S_n - S_{(n-1)}$   
 $\Rightarrow$  In an A.P If  $S_n$  is a quadratic of  $n$  ~~then~~  $T_n$  is linear  
in  $n$  Then the series will be an A.P.

$$T_n = A_n + B$$

$$S_n = An^2 + Bn + C$$

AP with common difference  $A$

$\hookrightarrow$  AP with common difference  $= 2A$

& Given  $S_n = 2n + 3n^2$ . A new AP is formed with some first term & double common difference. find sum of new AP.

$$\begin{aligned} d &= 2A \\ d &= 2 \times 3 \\ d &= 6 \\ a &= 2(1) + 3(1)^2 \\ a &= 2 + 3 \\ a &= 5 \\ S_n &= \frac{n}{2} [2a + (n-1)d] \\ &= \frac{n}{2} (10 + (n-1)12) \\ &= \frac{n}{2} (10 + 12n - 12) \\ &= \frac{n}{2} (12n - 2) \\ &= 6n^2 - n \end{aligned}$$

## Geometric Progression

→ Collection of non zero terms in which each consecutive term bears the some constant ratio.

$$a, ar, ar^2, ar^3, ar^4, ar^5, \dots ar^{n-1}$$

$$T_n = ar^{(n-1)}$$

or → First Term

$r \rightarrow$  common ratio  
 $r \neq 0$

$r \in (0, 1) \downarrow$  decrease

$r \in (1, \infty) \uparrow$  increase

$r \in (-1, 0) \downarrow$  ~~decrease in odd~~

$r < 0 \quad r \in (-\infty, -1) \uparrow$

Note - Common Ratio can be  $\frac{1}{2}$  or ~~-2~~

Q  $\frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \dots$   
find  $20^{\text{th}}$  term &  $n^{\text{th}}$  term.

$$\begin{aligned} r &= \frac{\frac{1}{4}}{\frac{1}{2}} \\ &= \frac{1}{4} \times \frac{2}{1} \\ &= \frac{1}{2} \end{aligned}$$

$$\begin{aligned} r &= \frac{\frac{1}{8}}{\frac{1}{4}} \\ &= \frac{1}{8} \times \frac{4}{1} \\ &= \frac{1}{2} \end{aligned}$$

$$a = \frac{1}{2}$$

$$\begin{aligned} T_{n^{\text{th}}} &= \frac{1}{2} \times \frac{1}{2}^{n-1} \\ &= \frac{1}{2^n} \\ &= 2^{-n} \end{aligned}$$

$$T_{20} = 2^{-20}$$

Q Find the ratio of 6<sup>th</sup> term & 9<sup>th</sup> term of the series.

$$\frac{S_4}{4}, \frac{S_5}{5}, \frac{S_6}{16} \dots$$

$$r = \frac{\frac{S_5}{5}}{\frac{S_4}{4}} = \frac{5}{4}$$

$$= \frac{5}{8} \times \frac{4}{5} = \frac{1}{2}$$

$$\frac{T_6}{T_9} = \frac{ar^5}{ar^8} = \frac{a}{r^3} = \left(\frac{1}{2}\right)^3 = \frac{1}{8}$$

$\boxed{= 8}$

Q In GP,  $T_3 = 2$

$$T_6 = \frac{1}{4}$$

$$ar^2 = 2$$

$$ar^5 = -\frac{1}{4}$$

$$\frac{ar^2}{ar^5} = -2 \times 8$$
$$\frac{ar^2}{ar^5} = -8 \times \frac{1}{r^3} = -8$$
$$\frac{1}{r^3} = -8$$
$$\frac{1}{r} = -2$$
$$\boxed{r = -\frac{1}{2}}$$

$$T_{10} = ar^9$$

$$= a \times \left(-\frac{1}{2}\right)^9$$

$$= +8 \times \frac{-1}{512}$$

$$= \frac{-1}{64}$$

## Summation GP

$$S_m = a + ar + ar^2 + ar^3 + \dots + ar^{m-1}$$

$$S_m = a + ar + ar^2 + ar^3 + \dots + ar^{m-2} + ar^{m-1}$$

$$\therefore S_m = ar + ar^2 + ar^3 + ar^4 + \dots + ar^{n-1} + ar^n$$

$$(1-r)S_m = a - ar^n$$

$$(1-r)S_m = a(1-r^n)$$

$$\boxed{S_m = \frac{a(1-r^n)}{1-r}} \quad (r \neq 1)$$

$$S_\infty = \frac{a(1-r^\infty)}{1-r}$$

$n \rightarrow \infty$

for  $r \in (-1, 1)$

$$S_\infty = \frac{a(1-r)}{1-r} = \frac{a}{1-r}$$

$$\boxed{S_\infty = \frac{a}{1-r}}$$

$$r \in (-1, 1)^\infty = \emptyset$$

$$r \in (-\infty, -1) \cup (1, \infty) = \infty$$

i)  $1, 5, 25, 125, \dots$

ii) sum of 30 terms.

iii) sum w/ infinite terms.

$$S_\infty = \infty \text{ Not Defined}$$

$$r = \frac{5}{1} = 5$$

$$S_{30} = \frac{1(1-5^{30})}{1-5}$$

$$= \frac{1-5^{30}}{-4}$$

$$= \frac{5^{30}-1}{4}$$

$$= \frac{5^{30}}{4} - \frac{1}{4}$$

$$Q \quad 1, \frac{1}{\sqrt{3}}, \frac{1}{3}, \frac{1}{3\sqrt{3}}$$

$$r = \frac{\frac{1}{\sqrt{3}}}{1}$$

$$= \frac{1}{\sqrt{3}}$$

- i) sum of 10 terms  
 ii) sum of infinite terms

$$S_{10} = 1 \left( 1 - \left( \frac{1}{\sqrt{3}} \right)^{10} \right)$$

$$= \frac{\left( \frac{1}{\sqrt{3}} \right)^{10} - 1}{\frac{\sqrt{3}-1}{\sqrt{3}}}$$

~~$$= \frac{242}{9}$$~~

$$= \frac{242\sqrt{3}}{\sqrt{3}-1}$$

$$= \frac{242\sqrt{3}(\sqrt{3}+1)}{2}$$

$$= 121(3+\sqrt{3})$$

$$= 363 + 121\sqrt{3}$$

$$S_\infty = \frac{a}{1-r}$$

$$= \frac{1}{\frac{\sqrt{3}-1}{\sqrt{3}}}$$

$$= \frac{\sqrt{3}}{\sqrt{3}-1}$$

$$= \frac{\sqrt{3}}{\sqrt{3}-1} \times \frac{\sqrt{3}+1}{\sqrt{3}+1}$$

$$= \frac{3+\sqrt{3}}{2}$$

Q (i) If  $a \neq 1, 0$  and  $(a + a^2 + a^3 + \dots) = S(a + a^3 + a^5 + \dots)$ .  
 find  $a$

$$(a + a^3 + a^5 + \dots) + (a^2 + a^4 + a^6 + \dots) = S(a + a^3 + a^5 + \dots)$$

$$\begin{aligned} S_{10} &= S \\ \frac{a^2}{1-a^2} &= S \\ a^2 &= S - Sa^2 \\ 6a^2 &= S \\ a^2 &= \frac{S}{6} \end{aligned}$$

$$a = \frac{\sqrt{S}}{\sqrt{6}}$$

$$\frac{a}{1-a} = S \left( \frac{a}{1-a^2} \right)$$

$$a(1-a)(1+a) = Sa(1-a)$$

$$a(1-a)(1+a-S) = 0$$

$$1 + a - S = 0$$

$$\text{but } a \in (-1, 1) \text{ so } a \in \emptyset$$

$$Q \quad (S^x + S^{(x+1)} + S^{(x+2)} + \dots) = 150 \left( \frac{1}{S} + \frac{1}{S^3} + \frac{1}{S^5} + \dots \right)$$

find  ~~$x$~~  ' $x$ ',

$$(S^x + S^{(x+1)} + S^{(x+2)} \dots) = 150 \left( S^{-1} + S^{-3} + S^{-5} + S^{-7} \dots \right)$$

$$a = S^x$$

$$r = S$$

$$a = S^{-1}$$

$$r = S^{-2}$$

$$\frac{S^x}{1-S} = 150 \left( \frac{1}{S^1} \right) \quad \left| \quad \frac{S^x}{1-S} = 150 \left( \frac{1}{S^1} \times \frac{25}{24} S \right)$$

$$\frac{S^x}{-4} = 150 \times \left( \frac{S}{24} \right)$$

$$S^x = -600 \times \frac{S}{24}$$

$$S^x = -300 \times \frac{S}{12}$$

$$S^x = -250 \times \frac{S}{3}$$

$$S^x = -250$$

$$\boxed{x \in \emptyset}$$

Q

$$10x^2 - 11x - 2 = 0$$

$$10(t + at + a^2t + a^3t + \dots) (1 + b + b^2 + b^3 + \dots)$$

$$a = 1$$

$$rb = a$$

$$a = 1$$

$$r = b$$

$$10 \left( \frac{1}{1-a} \right) \left( \frac{1}{1-b} \right)$$

$$\frac{10}{1-b-a+ab}$$

$$\frac{10}{1-(a+b)+ab}$$

~~$\frac{10}{1-b-a+ab}$~~

$$\frac{10}{1-\frac{1}{11}-\frac{2}{11}}$$

$$\frac{10 \times 11}{11-3} = \frac{110}{8} = \boxed{22}$$

product

$$2^{\frac{1}{4}} \cdot 4^{\frac{1}{16}} \cdot 8^{\frac{1}{64}} \cdot 16^{\frac{1}{128}} \dots \infty$$

$$2^{\frac{1}{4}} \cdot 2^{\frac{1}{8}} \cdot 2^{\frac{1}{16}}$$

$$r = \frac{2^{\frac{1}{16}}}{2^{\frac{1}{4}}} \\ = 2^{\frac{1}{16} - \frac{1}{4}}$$

$$r = 2^{-\frac{1}{4}}$$

$$2^{\frac{1}{4} + \frac{1}{8} + \frac{1}{16} \dots}$$

$$r = \frac{1}{8} \times \frac{4}{1}$$

$$= \frac{1}{2}$$

$$S_{\infty} = \frac{\frac{1}{4}}{1 - \frac{1}{2}}$$

$$= \frac{1}{4}$$

$$= \frac{2-1}{2}$$

$$= \frac{1}{4} \times \frac{2}{1}$$

$$= \frac{1}{2}$$

$$2^{\frac{1}{2}} = \boxed{\frac{1}{2}}$$

$$= \sqrt{2}$$

$$Q (\ln \alpha^2) + (\ln \alpha^2)^2 + (\ln \alpha^2)^3 \dots \infty = (\ln \alpha) + (\ln \alpha)^2 + (\ln \alpha)^3$$

$$\begin{aligned} x &= \ln \alpha^2 \\ \alpha &= e^{\ln \alpha^2} \end{aligned}$$

$$\begin{aligned} f_2 &= \ln \alpha \\ a &= e^{\ln \alpha} \end{aligned}$$

$$S_\infty = \frac{\ln \alpha^2}{1 - \ln \alpha^2}$$

$$S_\infty = \frac{\ln \alpha}{1 - \ln \alpha}$$

$$= \frac{2 \ln \alpha}{1 - 2 \ln \alpha}$$

$$= \frac{2 \log_e \alpha}{1 - 2 \log_e \alpha}$$

$$= \frac{2 \log_e \alpha}{e^{-2 \log_e \alpha}}$$

$$= \frac{2 \log_e \alpha}{\frac{\log_e \frac{\alpha}{e}}{\log_e \alpha^2}}$$

$$= \frac{\log_e \alpha^2}{\frac{\log_e \frac{\alpha}{e}}{\log_e \alpha^2}}$$

$$= \frac{\log_e \alpha^2}{\log_e \alpha^2}$$

$$\frac{\ln \alpha^2}{1 - \ln \alpha^2} = \frac{\ln \alpha}{1 - \ln \alpha}$$

$$\begin{aligned} 2 \ln \alpha &= \ln \alpha \\ 1 - 2 \ln \alpha &= 1 - \ln \alpha \end{aligned}$$

$$\frac{2x}{1 - 2x} = \frac{x}{1 - x}$$

$$2x - 2x^2 = x - 2x^2$$

$$x = 0$$

$$\ln \alpha = 0$$

$$a = 1$$

x reject

$$\ln \alpha = +\frac{1}{4}$$

$$\alpha = e^{+\frac{1}{4}}$$

$$Q \text{ If } a = \sum_{n=0}^{\infty} x^n, b = \sum_{n=0}^{\infty} y^n, c = \sum_{n=0}^{\infty} (xy)^n$$

$x, y \in (-1, 1)$

$$\text{find } \left( \frac{a-1}{a} \right) \left( \frac{b-1}{b} \right) - \frac{(c-1)}{c}$$

$$x = -1 \quad y = -1$$

$$\text{Ans: } 1, -1 + 1 - 1 - \dots \infty$$

$$a = 0$$

$$x = 1$$

$$a = \infty$$

$$y = -1$$

$$a = 0$$

$$a = \infty$$

$$\begin{array}{l} C = 0 \\ C = \infty \end{array}$$

$$a = \frac{1}{1-x}$$

$$b = \frac{1}{1-y}$$

$$c = \frac{1}{1-x}y$$


$$\left( \frac{a-1}{a} \right) \left( \frac{b-1}{b} \right) = \left( \frac{c-1}{c} \right)$$

$$\frac{\frac{1}{1-x}-1}{\frac{1}{1-x}} = \frac{\frac{1}{1-y}-1}{\frac{1}{1-y}} = \frac{\frac{1}{1-x}y-1}{\frac{1}{1-x}y}$$

$$(x)(y) = (xy)$$

$$xy = xy$$

$$\boxed{x=0}$$



$$a) 7 + 7 \cdot 10 + 7 \cdot 10^2 + \dots + 7 \cdot 10^n$$

$$7 + 7 \cdot 10 + 7 \cdot 10^2 + 7 \cdot 10^3 + \dots + 7 \cdot 10^n$$

~~$7 \cdot 10^0 + 7 \cdot 10^1 + 7 \cdot 10^2$~~

↓

$$7 \cdot 10^0 + 7 \cdot 10^1 + 7 \cdot 10^2$$

$$\text{S} \rightarrow a = 7$$

$$r = 10$$

$$S_n = \frac{7(1 - 10^n)}{1 - 10}$$

$$\begin{aligned} & \text{OTTOBLS} \\ & \text{ARCTIC AIR} \\ & \text{MECHANISCHE KÜHLEN} \end{aligned}$$

$$\begin{aligned} & \frac{7 - 7 \cdot 10^n}{-9} \\ & = \frac{7 \cdot 10^n - 7}{9} \end{aligned}$$

$$\frac{7 \cdot 10^0 - 7}{9} + \frac{7 \cdot 10^1 - 7}{9} + \frac{7 \cdot 10^2 - 7}{9} + \dots + 7 \cdot 10^n$$

$$\frac{7}{9} [10^0 - 1 + (10^1 - 1) + (10^2 - 1) + \dots + 10^n]$$

$$\frac{7}{9} [10^0 + 10^1 + 10^2 + \dots + (1 + \dots + 1) \cdot n]$$

$$\frac{7}{9} \left[ \frac{10(10^n - 1)}{9} - n \right]$$

$$\boxed{\frac{7}{9} \left( \frac{10(10^n - 1)}{9} - n \right)}$$

Q In a sequence of 4 nos., 1<sup>st</sup> 3 nos. are in G.P & last 3 are in A.P with CD=6 If 1<sup>st</sup> & last terms of the sequence are equal then find last term.

$$ar + ar^2 + ar^3 + a$$

$$ar^2 - ar = 6 \quad a - ar^2 = 6$$

$$ar(r-1) = 6 \quad a(1-r^2) = 6$$

$$\frac{6}{1-r^2} \times r^2 = \frac{6}{1-r^2} r = 6 \quad a = \frac{6}{1-r^2}$$

$$\frac{6r^2}{1-r^2} - \frac{6r}{1-r^2} = 6$$

$$6r^2 - 6r = 6 - 6r^2$$

$$12r^2 - 6r - 6 = 0$$

$$2r^2 - r - 1 = 0$$

$$r = \frac{1 \pm \sqrt{1+8}}{2}$$

$$= \frac{1 \pm 3}{2}$$

$$= 2, -1$$

$$a = \frac{6}{1-4}$$

$$a = \frac{6}{-3}$$

$$= -2$$

$$a = \frac{6}{1-1} \times$$

$$a+6, a-6, a, a+6$$

GP

$$\frac{a-6}{a+6} = \frac{a}{a+6}$$

$$a^2 + 36 - 12a = a^2 + 6a$$

$$36 = 18a$$

$$a=2$$


$$8, -4, 2, 8$$

OTTO BLS  
ARACTA  
MECHANISCHE  
FABRIK

$$8$$

Properties of GP

- ① Product of  $k^{\text{th}}$  term from beginning &  $k^{\text{th}}$  term from end is always constant.



- ② Assuming terms in GP

$$3: \frac{a}{r}, a, ar$$

$$4: \frac{a}{r^3}, \frac{a}{r}, ar, ar^3$$

$$5: \frac{a}{r^5}, \frac{a}{r^3}, \frac{a}{r}, ar^2, ar^5$$

$$6: \frac{a}{r^5}, \frac{a}{r^3}, \frac{a}{r}, ar, ar^3, ar^5$$

③ If each term of GP is raised to some power, multiplied or divided then the resulting series is also a GP.

$$2, 4, 8, 16, 32 \dots$$

$$2^{\frac{1}{8}}, 4^{\frac{1}{8}}, 8^{\frac{1}{8}}, 16^{\frac{1}{8}}, 32^{\frac{1}{8}} \dots$$

$$\cancel{2} \frac{4^{\frac{1}{8}}}{2^{\frac{1}{8}}} = 2^{\frac{1}{8}} \quad \frac{8^{\frac{1}{8}}}{4^{\frac{1}{8}}} = 2^{\frac{1}{8}}$$

④ If Three Terms  $A, B, C$  are in GP Then  $b^2 = ac$

$$a, b, c$$

$$\frac{b}{a} = \frac{c}{b}$$

$$b^2 = ac$$

are +ve real nos. So They are in

⑤ For  $a_1, a_2, a_3, \dots, a_n$  GP then

$\log a_1, \log a_2, \log a_3, \log a_4, \dots, \log a_n$  are in AP

e.g.  $2, 4, 8 \dots$  are in GP

$\log 2, \log 4, \log 8$  are in AP

## Harmonic Progression (H.P.)

→ A non zero sequence is said to be in H.P. if reciprocal of its terms are in A.P.

e.g.  $a_1, a_2, a_3, a_4, \dots$  H.P

$\frac{1}{a_1}, \frac{1}{a_2}, \frac{1}{a_3}, \frac{1}{a_4}, \dots$  A.P

→ Standard form.

$\frac{1}{a}, \frac{1}{a+d}, \dots, \frac{1}{a+(n-1)d}$  ... H.P

Note: ① No term of H.P. can be zero  
② Reciprocal of every H.P. is A.P. but reciprocal of every A.P. can or cannot be H.P.

③ If  $a, b, c$  are in H.P. then

$a, b, c \rightarrow \text{H.P}$

$\frac{1}{a}, \frac{1}{b}, \frac{1}{c} \rightarrow \text{A.P}$

$$2 \times \frac{1}{b} = \frac{1}{a} + \frac{1}{c}$$

$$\frac{2}{b} = \frac{a+c}{abc}$$

$$\boxed{\frac{b}{a+c} = \frac{2ac}{a+c}}$$

(4) No. general formula for finding the sum of H.P.

Q Determine the 4<sup>th</sup> & 8<sup>th</sup> term of H.P. 6, 9, 3 ...

$$\frac{1}{6}, \frac{1}{4}, \frac{1}{3} \dots A.P$$

$$\frac{1}{4} - \frac{1}{6} = C.P$$

$$d = \frac{6-4}{24}$$

$$= \frac{2}{24} \\ = \frac{1}{12}$$

$$a_4 = a + 3d \\ = \frac{1}{6} + 3 \times \frac{1}{12} \\ = \frac{1}{6} + \frac{1}{4} \\ = \frac{4+6}{24} \\ = \frac{10}{24} (A.P)$$

$$a_4 = \frac{24}{10} (H.P)$$

$$a_8 = a + 7d \\ = \frac{1}{6} + 7 \times \frac{1}{12} \\ = \frac{12+42}{60} \\ = \frac{54}{60} (A.P) \\ = \frac{9}{10} (H.P)$$

Q Find 16<sup>th</sup> term of H.P. if 6<sup>th</sup> & 11<sup>th</sup> term of H.P. are 10, 18 respectively.

$$\frac{1}{10} = a + 5d$$

$$\frac{1}{18} = a + 10d$$

$$a + 5 \times \frac{-2}{225} = \frac{1}{10} \\ a = \frac{1}{10} + \frac{2}{45}$$

$$\frac{1}{18} - \frac{1}{10} = 5d$$

$$\frac{-8}{180} = 5d$$

$$\frac{-8}{900} = d$$

$$d = -\frac{2}{225}$$

$$a = \frac{450513}{45090}$$

$$a_{16} = \frac{47}{450} - 15 \times \frac{2}{225} \\ a_{16} = \frac{47-60}{450} \\ a_{16} = -\frac{13}{450}$$

$$a_{16} = \frac{13}{90} - 15 \times \frac{2}{225}$$

$$a_{16} = 90$$

Q If  $a, b, c, d, e$  are nos,  $a, b$  in GP.


$b, c, d$  in GP.

$c, d, e$  in HP.

a) Then  $a, c, e$  in GP

b)  $c = \frac{(2b-a)^2}{a}$

$$a-d, a, a+d, \frac{(a+d)^2}{a}, \frac{(a+d)^2}{a-d}$$



c)  $\frac{c}{a} = \frac{a+d}{a-d}$

$$\frac{e}{c} = \frac{(a+d)^2}{a-d} \times \frac{1}{a+d}$$

$$\frac{e}{c} = \frac{a+d}{d-d}$$

$\frac{c}{a} = \frac{e}{c}$  so  $a, c, e$  are in GP.

$$6) \frac{(2b-a)^2}{a}$$

$$\cancel{2a+d}$$

$$\frac{(2a-a+d)^2}{a-d}$$

$$= \frac{(a+d)^2}{a-d}$$

$$e = \frac{(a+d)^2}{a-d}$$

Hence LHS = RHS.

Method II

$$Qf = \frac{a+c}{2}$$

$$d = \frac{2ce}{c+e}$$

$$c^2 = fd$$

$$c^2 = \left(\frac{a+c}{2}\right)\left(\frac{2ce}{c+e}\right)$$

$$= \frac{2ace + 2c^2e}{2c + 2e}$$

$$= \frac{ace + c^2e}{c+e}$$

$$c^2(c+e) = (a+c)ce$$

$$c^2e + c^3e = ace + c^2e$$

~~$c^2e + c^3e$~~

~~$a = c^2$~~

$$c^2 = ae$$

Thus,  $a, c, d, e$  are in GP

$$6) \quad c = \frac{(2b-a)^2}{a}$$

$$\therefore c = (2b-a)^2$$

$$c^2 = (2b-a)^2$$

$$c = 2b-a = LHS$$

$$b = \frac{a+c}{2}$$

$$2b = a+c$$

$$2b - a = c = LHS = RHS$$

Arithmetic  
Mean

∴ Proved.

Mean :-

① Arithmetic Mean (AM) -

$$2, 4, 6 \dots (AP)$$

$$\frac{6+2}{2} = \frac{8}{2} = 4 \text{ (mean)}$$

$$2, 4 \text{ के बीच का mean?}$$

$$2, 3, 4$$

$$2, 2.5, 3$$

$a, b, c$  are in AP

$$b = \frac{a+c}{2}$$

Find means b/w  $a$  &  $b$ ?

$a, AM_1, AM_2, AM_3, AM_4, \dots, AM_n, b$  (AP)

$$A \quad b = a + (n+2-1)d$$

$$b = a + (n+1)d$$

$$b - a = (n+1)d$$

$$d = \frac{b-a}{n+1}$$

$$AM_1 = a + d = a + \frac{(b-a)}{n+1}$$

$$AM_2 = a + 2\left(\frac{b-a}{n+1}\right)$$

$$AM_n = a + (n-1)\left(\frac{b-a}{n+1}\right)$$

Sum of all AMs

$$AM_1 + AM_2 + AM_3 + \dots + AM_n$$

$$na + \left(\frac{b-a}{n+1}\right)(1+2+3+4+\dots+n)$$

$$na + \left(\frac{(b-a)}{n+1}\right) \times \frac{n(n+1)}{2}$$

$$na + \frac{n(b-a)}{2}$$

$$2na + nb = nb$$

$$\frac{nb + na}{2}$$

$$\frac{n(a+b)}{2} = S_n \text{ of AMs}$$

Q. Inscriit 20 AMs between 4 & 67

- ① find  $A.M.s$
- ② find sum of  $A.M.s$

e.g. 4,  $A.M_1$ ,  $A.M_2$ , ..., 67

$$n = 20$$

$$\textcircled{2} \quad 20 \left( \frac{67 + 4}{2} \right)$$

$$20 \cdot 10 \times 71$$

$$\boxed{S_n = 710}$$

$$\textcircled{1} \quad A.M_1 = 4 + n \left( \frac{63}{m+1} \right)$$

$$A.M_1 = 4 + \cancel{\frac{63}{21}}$$

$$A.M_2 = 4 + \cancel{\frac{2 \times 63}{21}}$$

$$= 4 + 4.2$$
  
$$= 4.6$$

$$\textcircled{1} \quad A.M = 4 + n \left( \frac{63}{21} \right)$$

$$= 4 + 3n$$

$$A.M_1 = 7$$

$$A.M_2 = 10$$

$$A.M_3 = 13$$

$$A.M_4 = 16$$

$$A.M_5 = 19$$

$$A.M_6 = 22$$

⋮

Q find  $s_0 A M_s$  when  $I \& 99$

$$S_m = \frac{99+1}{2} \times s_0$$

$$= s_0 \times s_0$$

$$\boxed{= 2500}$$

& If  $p A M_s$  required between  $P_5$  &  $P_1$ ,  $\frac{A M_3}{A M_{P-1}} = \frac{2}{s}$   
find ' $p'$

$$A M_3 = 5 + \frac{3}{s} \left( \frac{36}{P+1} \right)$$

$$= \frac{5P + 5 + 108}{P+1}$$

$$= \frac{5P + 113}{P+1}$$

$$A M_{P-1} = \frac{5P+5 + (P-1)36}{P+1}$$

$$= 5P + 5 + 36P - 36$$

$$= \frac{41P - 31}{P+1}$$

$$\frac{(5P + 113)}{(P+1)} \times \frac{(P+1)}{(41P - 31)} = \frac{2}{s}$$

$$\frac{25P + 565}{627} = \frac{82P - 62}{579}$$

$$\boxed{P = 11}$$

H.W. 25-06-24

DYS-3

RoC - 20-23

~~DYS = 5 (Q3-4)~~

Geometric mean. :-

$$a, \underbrace{Gm_1, Gm_2, Gm_3, Gm_4, \dots, b}_{n} \quad (\text{Total } n Gm_i + 2(a, b))$$

$$b = ar^{(n+1)+1}$$

$$b = ar^{n+1}$$

$$\left[ \left( \frac{b}{a} \right)^{\frac{1}{n+1}} = r \right]$$

$$Gm_1 = ar^2$$

$$Gm_2 = ar^2$$

$$\boxed{Gm_n = ar^n}$$

Product of Gms

$$AM_1 \times AM_2 \times AM_3 \times AM_4 \times AM_5 \dots GM_n$$

$$Gm_1 \times Gm_2 \times Gm_3 \times Gm_4 \times Gm_5 \dots GM_n$$

$$a \left( \frac{b}{a} \right)^{\frac{1}{n+1}} \times a \left( \frac{b}{a} \right)^{\frac{2}{n+1}} \times a \left( \frac{b}{a} \right)^{\frac{3}{n+1}} \dots \left( \frac{b}{a} \right)^{\frac{n}{n+1}}$$

$$a^n \left( \frac{b}{a} \right)^{\frac{1}{n+1}(1+2+3\dots n)}$$

$$a^n \left( \frac{b}{a} \right)^{\frac{1}{n+1}(\frac{n}{2} \times (n+1))}$$

$$a^n a \left( \frac{b}{a} \right)^{\frac{n}{2}}$$

$$\left( a \cdot \frac{\sqrt{b}}{\sqrt{a}} \right)^n$$

$$\boxed{\left( \sqrt{ab} \right)^n}$$

Q Insert 4 GPs & the SD of 160. find product of GP's.

$$A = \left(\frac{160}{5}\right)^{\frac{1}{5}}$$

$$= (32)^{\frac{1}{5}}$$

$$\underline{A = 2}$$

GPs  $\rightarrow$  10, 20, 40, ..., 80

$$\text{Product} = (\sqrt{5 \times 160})^4$$

$$\begin{aligned} &= (800)^2 \\ &= 640000 \\ &\boxed{= 640000} \end{aligned}$$

Q find GPs of the series  $3 + 3^1 + 3^2 + 3^3 + \dots + 3^n$

$$\begin{aligned} A.P. &= \sqrt{3 \times 3^1 \times 3^2 \times 3^3 \dots \times 3^n} \\ &= \sqrt{3 \times 3^{\frac{n(n+1)}{2}}} \\ &= 3^{\frac{n(n+1)}{2}} \times \frac{1}{2} \\ &= 3^{\frac{n(n+1)}{2}} \\ &= 3^{1+2+3+4+5+\dots+n} \\ &\boxed{3^{\frac{n(n+1)}{2}}} \end{aligned}$$

$$\left[ 3^{\frac{n(n+1)}{2}} \right]^{\frac{1}{n}}$$

$$\boxed{3^{\frac{(1+n)}{2}}}$$

$$\begin{aligned} &\sqrt{ab} \\ &\sqrt[3]{abc} \\ &\sqrt[4]{abcd} \end{aligned}$$

a) find product of 3 G.M.s between 2 & 8.

$$\text{Product} = (\sqrt{ab})^{15}$$
$$= (2 \times 8)^{15}$$
$$= 16^{15}$$

Q)  $AM(a, b) = 15$   
 $GM(a, b) = 9$ . find  $a, b$ .

$$\frac{a+b}{2} = 15$$

$$\sqrt{ab} = 9$$

$$ab = 81$$

$$a+b = 30$$

$$b = 30-a$$

$$(30-a) a = 81$$

$$30a - a^2 = 81$$

$$a^2 - 30a + 81 = 0$$

$$a = \frac{30 \pm \sqrt{900 - 4 \cdot 324}}{2}$$

$$a = \frac{30 \pm 24}{2}$$

$$a = 15 \pm 12$$

$$\therefore a = 27, 3$$

$$\Rightarrow [3, 27]$$

Harmonic Mean

$a, HM_1, HM_2, HM_3, \dots, HM_{n-1}, b$

$$\frac{1}{a}, \frac{1}{HM_1}, \frac{1}{HM_2}, \dots, \frac{1}{HM_n}, \frac{1}{b}, \dots, A.P.$$

$$\frac{1}{b} = \frac{1}{a} + (n+2-1)d$$

$$\frac{1}{b} - \frac{1}{a} = (n+1)d$$

$$d = \frac{a-b}{ab(n+2)}$$

3656  
371  
186  
540  
676

$$AM_1 = \frac{1}{a+d}$$

$$\frac{a}{d} = \frac{1}{a+d}$$

$$HM_1 = \frac{1}{\frac{a}{d} + 1}$$

$$HM_2 = \frac{1}{a+2d}$$

$$HM_n = \frac{1}{a+nd}$$

Q Insert 5 HM betw  $\frac{1}{3}$  &  $\frac{1}{21}$ .

$$d = \frac{\frac{1}{3} - \frac{1}{21}}{\frac{1}{3} \times \frac{1}{21} (5+1)}$$

$$\begin{aligned} &= \frac{\frac{18}{63} - \frac{1}{63}}{\frac{1}{63} \times 6} \\ &= \frac{\frac{17}{63}}{\frac{1}{63} \times 6} \\ &= \frac{17}{6} \\ &= 3 \end{aligned}$$

$$\begin{aligned} HM_1 &= \frac{1}{\frac{1}{3} + \frac{18}{7}} \\ &= \frac{1}{\frac{61}{21}} \\ &= \frac{21}{61} \end{aligned}$$

$$\begin{aligned} HM_2 &= \frac{1}{\frac{1}{3} + \frac{30}{7}} \\ &= \frac{1}{\frac{31}{7}} \\ &= \frac{7}{31} \end{aligned}$$

$$HM_1 = \frac{1}{a+c}$$

$$HM_2 = \frac{1}{\frac{1}{3} + \frac{29}{30}}$$

$$\cancel{\frac{1}{17}} = \frac{3}{18}$$

$$= \frac{1}{6}$$

$$= \frac{3}{30}$$

$$= \frac{1}{10}$$

Q. HM of roots of eq  $(5+\sqrt{2})x^2 - (4+\sqrt{3})x + (8+2\sqrt{3}) = 0$

$$x = \frac{(4+\sqrt{3}) \pm \sqrt{16+3+8\sqrt{3}-160-40\sqrt{3}-32\sqrt{2}-8\sqrt{6}}}{2(5+\sqrt{2})}$$

$$HM = \frac{2\alpha\beta}{\alpha+\beta}$$

$$= 2 \frac{(8+2\sqrt{3})}{5+\sqrt{2}}$$

$$\frac{4+\sqrt{3}}{5+\sqrt{2}}$$

$$= 2 \frac{16+4\sqrt{3}}{16+4\sqrt{3}}$$

$$= \frac{4(4+\sqrt{3})(4-\sqrt{3})}{(4-\sqrt{3})(4+\sqrt{3})}$$

$$= 4$$

H.W. 27-6-24

DYS-S (2-13)

Q Insert 2 H.M. b/w  $2.5$  &  $0.4$

$$\frac{2}{5}, \frac{2}{3}$$

$$d = \frac{\frac{2}{5} + \frac{2}{3}}{3}$$

$$= \frac{4+25}{30}$$

$$= \frac{29}{30}$$

$$HM_1 = \frac{1}{\frac{2}{5} + \frac{29}{30}}$$

$$= \frac{30}{41}$$

$$HM_2 = \frac{1}{\frac{2}{3} + \frac{0.58}{30}}$$

$$= \frac{30}{70}$$

$$= \frac{3}{7}$$

AM, GM, HM (Inequality & Relation)

Properties:-

$a, b, c, d \in \text{Positive Real nos.}$

(1)

$a, b$

$$AM = \frac{a+b}{2}$$

$$GM = \sqrt{ab}$$

$$HM = \frac{2}{\frac{1}{a} + \frac{1}{b}}$$

$a, b, c$

$$AM = \frac{a+b+c}{3}$$

$$GM = \sqrt[3]{abc}$$

$$HM = \frac{3}{\frac{1}{a} + \frac{1}{b} + \frac{1}{c}}$$

$$AM = \frac{a+b+c+d}{4}$$

$$GM = \sqrt[4]{abcd}$$

$$HM = \frac{4}{\frac{1}{a} + \frac{1}{b} + \frac{1}{c} + \frac{1}{d}}$$

$\alpha_1, \alpha_2, \alpha_3, \dots, \alpha_n$

$$AM = \frac{\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4 + \dots + \alpha_n}{n}$$

$$GM = \sqrt[n]{\alpha_1 \times \alpha_2 \times \alpha_3 \times \alpha_4 \times \dots \times \alpha_n}$$

$$HM = \frac{1}{\frac{1}{\alpha_1} + \frac{1}{\alpha_2} + \dots + \frac{1}{\alpha_n}}$$

(2)

$$AM \geq GM \geq HM$$

$$AM = GM = HM \text{ when } \alpha_1 = \alpha_2 = \alpha_3 = \dots = \alpha_n$$

Proof

$$AM \geq GM$$

$$\frac{a+b}{2} \geq \sqrt{ab}$$

$$(\sqrt{a})^2 + (\sqrt{b})^2 - 2\sqrt{ab} \geq 0$$

$$(\sqrt{a} - \sqrt{b})^2 \geq 0$$

always true

$$GM \geq HM$$

$$\sqrt{ab} \geq \frac{2ab}{a+b}$$

$$\sqrt{ab}(a+b) - 2ab \geq 0$$

$$\sqrt{ab}(a+b - 2\sqrt{ab}) \geq 0$$

$$(\sqrt{a} - \sqrt{b})^2 \geq 0$$

Always true

(3) For any ~~two~~<sup>three</sup> numbers there AM, GM & HM are in A.P.

$$GM^2 = AM \times HM$$

(4)  $x^2 - 2AMx + GM^2 = 0$

$$x^2 - (a+b)x + ab = 0$$

$$AM = \frac{a+b}{2} \Rightarrow a+b = 2(AM)$$

$$GM = \sqrt{ab} \Rightarrow GM^2 = ab$$

$$x^2 = ab$$

$$x^2 - 2AMx + GM^2$$

(5) If  $a$  and  $b$  are two numbers and  $\frac{a^n + b^n}{a^n + b^n}$

$$n=0 \Rightarrow \frac{a+b}{2} = AM$$

$$n=-1 \Rightarrow \sqrt{ab} = GM$$

$$n=-2 \Rightarrow \frac{2ab}{a+b} = HM$$

(6) If ~~am~~ are  $a, b, c$  are in A.P. as well as in G.P.  
~~and~~ then  $a=b=c \neq 0$

(7) i)  $\frac{a-b}{b-c} = \frac{a}{c} = 1$ ;  $a, b, c$  are in G.P.  $\Rightarrow a, b, c$  are in G.P.

ii)  $\frac{a-b}{b-c} = \frac{a}{b}$  or  $\frac{a-b}{b-c} = \frac{b}{c} \Rightarrow ac - bc = b^2 - bc$

iii)  $\frac{a-b}{b-c} = \frac{c}{b}$ ;  $a, b, c$  are in H.P.

## Inequalities

Q1. If  $x \& y \in R^+$  Then  $\frac{xy}{y} + \frac{y}{x}$  min

$$GM = \sqrt{\frac{x}{y} \times \frac{y}{x}} = \sqrt{1} = 1$$

$$AM = \frac{\frac{x}{y} + \frac{y}{x}}{2}$$

$$AM \geq GM$$

$$\frac{xy}{y} + \frac{y}{x} \geq 2$$

~~$\frac{x^2+y^2}{2} \geq xy$~~

$$T_{\min} = 2$$

Q2.  $x \& y \in R^+$   $\left(\frac{1}{x} + \frac{3}{y}\right)$  min value  $xy^3 = 16$

$$\begin{aligned} & \cancel{\frac{1}{x} + \frac{3}{y}} = \frac{3x + y}{xy} \\ & \frac{1}{x}, \frac{3}{y} \quad \frac{3}{y} \\ & \frac{1}{y}, \frac{1}{y} \quad \frac{1}{y} \\ & \frac{1}{x} + \frac{1}{y} + \frac{1}{y} + \frac{1}{y} \geq 4\sqrt[4]{\frac{1}{x} \cdot 3} \end{aligned}$$

$$\frac{1}{x} + \frac{3}{y} \geq 2$$

$$\boxed{\min = 2}$$

$$\text{Q} \quad p + 2\theta + 4R = 9 \quad P, Q, R \in \mathbb{F}^+$$

find  $(P^2 \oplus^4 R^3)_{\text{max}}$

$$\begin{array}{c}
 \cancel{37} \quad 2 \sqrt{PQR} \\
 \cancel{27} \quad PQR \\
 \hline
 \cancel{8} \quad PQR = \cancel{27} \\
 \max
 \end{array}$$

```

graph LR
    P((P)) -- "1/2" --> Q1((Q))
    Q1 -- "1/2" --> R1((R))
    Q1 -- "1/2" --> R2((R))
    Q1 -- "1/2" --> R3((R))
    R1 -- "1/3" --> S1((S))
    R2 -- "1/3" --> S2((S))
    R3 -- "1/3" --> S3((S))
  
```

$$\frac{q}{q} \geq \sqrt{\frac{P^2 + R^3}{2R}}$$

$$I^q \geq P^2 Q^q R^{\frac{q}{2}}$$

$$P^2 \varnothing^4 R^3 \leq 27$$

$$\text{mod } c = 17$$

$$x^3 + x^2 + x + 1 \geq x^3 \text{ für } x > 0.$$

$\alpha$  if  $\alpha > 0$  &  $\beta = \alpha$

$$\begin{aligned} & \cancel{(1+0)(1+0^2)} \\ & \cancel{(1+0)(1+0^2)} \\ & \cancel{(1+0)(1+0^2)} \\ & \cancel{(1+0)(1+0^2)} \end{aligned}$$


A hand-drawn diagram showing two sets of intersecting curves. The top set consists of curves labeled 3,  $0^2 + 0 + 1$ , 3, 4, and  $0^6$ . The bottom set consists of curves labeled  $0^3 + 0^2 + 0 + 1$ , 4, and  $0^{3/2}$ . A label "BESCHREIBUNG" is written below the bottom set.

H.W. (28-6-24)

0-1 (01, 2-5, 7, 11, 13, 15, 17, 19)

$a^3 + a^2 + a + 1 \geq ka^3$ ,  $a \neq 0$

Not feasible



$$2x + 3y = 15. \quad (\text{first } \Rightarrow \text{ max})$$

$$\frac{x+x+3y}{3} = \sqrt[3]{3x^2y}$$

22 5x+5

$$5 \cdot \frac{15}{3} \geq \sqrt[3]{3x^2y}$$

$$125 \geq 3x^2y$$

$$\frac{125}{3} \rightarrow 0 \ x^2y \text{ TOOLS}$$

$$\boxed{\max(x^2y) = \frac{125}{3}}$$

$$\text{Q } a, b, c \in \text{Positive real } ab^2c^3 = 64 \quad \left(\frac{1}{a} + \frac{2}{b} + \frac{3}{c}\right) \text{ min.}$$

$$\frac{1}{a} + \frac{9}{b} + \frac{1}{b} + \frac{1}{c} + \frac{1}{c} + \frac{1}{c}$$

$$\frac{10}{6} = \sqrt[6]{\frac{1}{64}}$$

$$x = 6 \times 2^{-1}$$

$$x = \frac{6}{2}$$

$$x = 3$$

$$\text{Q } a, b, c, d > 0 \quad a + 2b + 3c + 4d = 50 \quad \left(\frac{a^2 b^4 c^3 d}{16}\right)^{\frac{1}{10}} \text{ max}$$

$$\frac{a}{2} + \frac{a}{2} + \frac{b}{2} + \frac{b}{2} + \frac{b}{2} + \frac{b}{2} + c + c + c + c + 4d \geq \sqrt[10]{\frac{a^2 b^4 c^3 d}{16}}$$

10

$$\boxed{1 - 5 =}$$

$$\text{Q } p, q, r \in \mathbb{R}^+ \quad pq = 1$$

$$\sqrt{(1+p+p^2)(1+q+q^2)(1+r+r^2)} \geq \min$$

$$1 \leq p(1+p)$$

$$\frac{1+p+p^2}{3} \geq (1+p+p^2)^{\frac{1}{3}} \geq p$$

$$\frac{1+q+q^2}{3} \geq q$$

$$\frac{1+r+r^2}{3} \geq r$$

multiply

$$(1+p+p^2)(1+q+q^2)(1+r+r^2) \geq pqrR$$

.27

$$\boxed{\min = 27}$$

Q If  $\alpha_1, \alpha_2, \alpha_3, \dots, \alpha_n \in \mathbb{R}^+$   
&  $\alpha_1 \times \alpha_2 \times \alpha_3 \dots \alpha_n = 1$

$$\frac{1+\alpha_1+\alpha_1^2}{3} \geq \alpha_1$$

$$\frac{1+\alpha_2+\alpha_2^2}{3} \geq \alpha_2$$

$$\frac{1+\alpha_n+\alpha_n^2}{3} \geq \alpha_n$$

$$\frac{(1+\alpha_1+\alpha_1^2)(1+\alpha_2+\alpha_2^2)\dots(1+\alpha_n+\alpha_n^2)}{3^n} \geq 1$$

$$\boxed{3^n = \min}$$

$$\text{Q } x > 0, \left( \frac{x^{10}}{1+x+x^2+\dots+x^{20}} \right)_{\text{max}}$$

$$\begin{aligned} & \cancel{x^{10}} \\ & \cancel{\frac{x^{20}-1}{x-1}} \\ & x^{10}(x-1) \\ & \frac{1+\dots}{21} \geq x^{\frac{20}{21}} \end{aligned}$$

$$\begin{aligned} & \frac{1+x^{20}}{21} \Rightarrow \text{OTTOSLS} \\ & \frac{1+x^{20}}{x^{10}} \Rightarrow 21 \\ & \frac{x^{10}}{1+x^{20}} \leq \frac{1}{21} \\ & \boxed{\text{max} = \frac{1}{21}} \end{aligned}$$

### Arith - Geometric Series (A.G.P)

→ Series formed by multiplying the corresponding terms of AP and GP.

$$a b, (a+d)b r, (a+2d)b r^2, \dots, (a+(n-1)d)b r^{n-1}$$

$a$  → first term of AP

$d$  → CP of AP

$b$  → 1st term of GP

$r$  → C.R of GP

$$T_n = (a + (n-1)d) b^{n-1}$$

Sum of A.G.P :-

Process :-

- ① Find the C.R.
- ② Multiply the C.R. with given series to form a new series.
- ③ Subtract the given series & the new series.
- ④ After subtraction we either get AP or GP hence Apply sum.
- ⑤ In rare case we have to repeat the subtraction process two or three times if we don't get the A.P. or G.P. after first subtraction.

~~Q~~ If  $|x| < 1$ , find the sum of

~~a.~~  $\frac{1}{1+x} + \frac{1}{1+2x} + \dots$  find sum

$$1 + 2x + 3x^2 + 4x^3 + \dots \infty$$

$$S = 1 + 2x + 3x^2 + 4x^3 + \dots \infty$$

$$CR = x$$

$$\text{Ans} Sx = x + 2x^2 + 3x^3 + 4x^4 + \dots \infty$$

$$S - Sx = (x-1) + 2x(x-1) + 3x^2(x-1) + \dots \infty$$

$$S = 1 + 2x + 3x^2 + \dots$$

$$S(1-x) = 1 + x + x^2 + x^3 + \dots \infty$$

$$S(1-x) = \frac{1}{1-x}$$

$$S = \frac{1}{(1-x)^2}$$

$$Q \quad S = 2 + \frac{4}{3} + \frac{6}{3^2} + \frac{8}{3^3} + \frac{10}{3^4} + \dots$$

$$\frac{S}{3} = \frac{2}{3} + \frac{4}{3^2} + \frac{6}{3^3} + \frac{8}{3^4} + \dots$$

$$S - \frac{S}{3} = 2 + \frac{2}{3} + \frac{2}{3^2} + \frac{2}{3^3} + \frac{2}{3^4} + \dots$$

$$S\left(1 - \frac{1}{3}\right) = \frac{2}{1 - \frac{1}{3}}$$

$$S\left(\frac{2}{3}\right) = \frac{2 \times 3}{1 - \frac{1}{3}}$$

$$S = \frac{2}{2} \times \frac{3}{3}$$

$$\boxed{S = \frac{9}{2}}$$

$$Q \quad 1 + 2 \times 2 + 3 \times 2^2 + 4 \times 2^3 + \dots + 100 \times 2^{99}, \text{ sum.}$$

$$S = 1 + 2 \times 2^1 + 3 \times 2^2 + 4 \times 2^3 + \dots + 100 \times 2^{99}$$

$$2S = 2^1 + 2 \times 2^2 + 3 \times 2^3 + 4 \times 2^4 + \dots + 200 \times 2^{100}$$

$$S - 2S = 1 + 2^1$$

$$S = 1 + 2^1 + 2^2 + 2^3 + \dots + 2^{99} = 100 \times 2^{99}$$

$$\cancel{-S = 1}$$

$$\cancel{-S = 1 - 2^1}$$

$$\boxed{\cancel{S = 1 - 2^1}}$$

$$-S = \frac{(1 - 2^{100})}{2(1 - 2)} - 100 \times 2^{100}$$

$$S = 1 - 2^{100} + 100 \times 2^{100}$$

$$\cancel{S = 1 - 101 \times 2^{100}}$$

$$\boxed{S = 99 \times 2^{100} + 1}$$

$$\cancel{\textcircled{1}} \quad \cancel{\frac{4}{3^2} + \frac{4}{3^3} + \dots}$$

$$\textcircled{2} \quad \frac{4}{7} - \frac{5}{7^2} + \frac{4}{7^3} - \frac{5}{7^4} + \frac{4}{7^5} - \frac{5}{7^6} \dots \infty$$

$$\frac{(28-5)}{7^2} + \frac{(28-5)}{7^4} + \frac{(28-5)}{7^6} \dots \infty$$

$$d = (28-5)/7^2$$

$$r = 1/7^2$$

$$S_{\infty} = \frac{d}{1-r}$$

$$= \frac{(28-5)}{7^2} \cdot \frac{1}{1-\frac{1}{7^2}}$$

$$= \frac{23}{48} \cdot \frac{49}{48}$$

$$\boxed{= \frac{23}{48}}$$

$$\textcircled{3} \quad S = 1 + 3x + 6x^2 + 10x^3 + \dots \infty$$

$$S_{2x} = x + 3x^2 + 6x^3 + 10x^4 + \dots \infty$$

$$S - S_{2x} = 1 + 2x + 3x^2 + 4x^3 \dots \infty$$

$$S(1-x) = 1 + 2x + 3x^2 + 4x^3 \dots \infty$$

$$Sx(1-x) = x + 2x^2 + 3x^3 \dots \infty$$

$$S(1-x)^2 = 1 + x + x^2 + x^3 \dots \infty$$

$$S(1-x)^2 = \frac{1}{1-x}$$

$$S = \frac{1}{(1-x)^3}$$

Q Sum of 10 terms

$$\left(x + \frac{1}{x}\right)^2 + \left(x^2 + \frac{1}{x^2}\right)^2 + \left(x^3 + \frac{1}{x^3}\right)^2 \dots$$

$$\left(x + \frac{1}{x}\right) + \left(x^2 + \frac{1}{x^2}\right)$$

$$\left(x^2 + \frac{1}{x^2} + 2\right) + \left(x^4 + \frac{1}{x^4} + 2\right) + \left(x^6 + \frac{1}{x^6} + 2\right) \dots$$

$$x^2 + x^4 + x^6 + x^8 \dots + \frac{x^{20}}{x^2} + \frac{1}{x^4} + \frac{1}{x^6} \dots + 2 + 2^{20} \dots$$

$$D = x^2$$

$$d = x^2$$

$$S_{10} = \frac{x^2(x^{20}-1)}{x^2-1}$$

OTTOBLS

ANACTACT

$$S_{10} = \frac{x^2(\chi^{20}-1)}{\sqrt{x^{20}-1}} \quad S_{10} = 20$$

$$= \frac{x^2(\chi^{20}-1)x^2}{1-x^2}$$

$$= \frac{(1-\frac{1}{x^{20}})}{x^2-1}$$

$$\frac{x^2(x^{20}-1) + 1 - \frac{1}{x^{20}}}{x^2-1} + 20$$

$$\frac{x^{40} - x^2 + 1 - \frac{1}{x^{20}} + 20x^2 - 20}{x^2-1}$$

$$\frac{x^{60} - x^{22} + x^{20} - 1 + 20x^{22} - 20x^{20}}{x^{20}(x^2-1)}$$

$$\frac{x^{60} + 19x^{22} - 19x^{20} - 1}{x^{20}(x^2-1)}$$

$$\cancel{\frac{x^{40} + 19x^2 - 19}{x^2-1}}$$

# Special Sequence

Type-1 :- Using Summation ( $\Sigma$ )

Type-2 :- Method of Difference (MOD)

Type-3 :- Splitting in denominators

Type-4 :- Splitting in Numerators

Q. W- 1-7-24

DYS-6 (Full)

Type-1 :- Results :-

$$\textcircled{1} \quad 1 + 2 + 3 + 4 + 5 + \dots + n = \frac{n(1+n)}{2}$$

$$\textcircled{2} \quad 1 + 2^2 + 3^2 + 4^2 + 5^2 + \dots + n^2 = \frac{n(n+1)(2n+1)}{6}$$

$$\textcircled{3} \quad 1 + 2^3 + 3^3 + 4^3 + 5^3 + \dots + n^3 = \left(\frac{n(n+1)}{2}\right)^2$$

$$\textcircled{4} \quad 1 + 3 + 5 + \dots + (2n-1) = n^2$$

$$\textcircled{5} \quad 2 + 4 + 6 + \dots + 2n = n(n+1)$$

Proof:-

$$\textcircled{2} \quad 1^2 + 2^2 + 3^2 + \dots + n^2 = \frac{n(n+1)(2n+1)}{6}$$

$$(x+1)^3 = x^3 + 1 + 3x + 3x^2$$

$$(x+1)^3 - x^3 = 1 + 3x + 3x^2$$

~~$$x^3 - 0^3 = 1 + 0 + 0$$~~

$$x=1 \quad 2^3 - 1^3 = 1 + 3(1) + 3(1)^2$$

$$x=2 \quad 3^3 - 2^3 = 1 + 3(2) + 3(2)^2$$

$$x=3 \quad 4^3 - 3^3 = 1 + 3(3) + 3(3)^2$$

$$x=n \quad (n+1)^3 - n^3 = 1 + 3(n) + 3(n)^2$$

ddd dle

$$(n+1)^3 - 1 = n + 3(1^2 + 2^2 + 3^2 + \dots + n^2) + 3 \frac{(1^2 + 2^2 + 3^2 + \dots + n^2)}{t}$$

$$\cancel{(n+1)^3 - 1} = \cancel{n} + \frac{3n(n+1)}{2} + 3t$$

$$(n+1)^3 - 1 - n - \frac{3n(n+1)}{2} = 3t$$

$$2(n+1)^3 - 2n - 3n(n+1) - 2 = 3t$$

$$\frac{2n^3 + 2n^2 + 6n + 6n^2 + 3n^3 - 3n^2 - 3n - 2}{2} = 3t$$

$$3t = \frac{2n^3 + 3n^2 + n}{2}$$

$$3t = \frac{n(2n^2 + 3n + 1)}{2}$$

$$3t = \frac{n(n+1)(2n+1)}{2}$$

$$t = \frac{n(n+1)(2n+1)}{6}$$

Method to find a sum :-

→ when  $n^{th}$  term is given. → Apply  $S_n = \sum T_n$

→ when  $n^{th}$  term is not given:- → Find  $n^{th}$  term → Apply  $S_n = \sum T_n$

Q find the sum of  $n$  terms upto  $n^{\text{th}}$  term is

①  $2n + 3n - 1$

2, 5, 8, 11, ... general

$$S_n = \sum (3n - 1)$$

$$= \sum 3n - \sum 1$$

$$= 3\sum n - \sum 1$$

$$= 3(1 + 2 + 3 + \dots + n) - (\underbrace{1 + 1 + 1}_{n \text{ times}})$$

$$= 3 \times \frac{n(n+1)}{2}$$

~~$$= 3n^2 + 3$$~~

$$= n \left( \frac{3(n+1)}{2} - 1 \right)$$

$$= n \left( \frac{3n+3-2}{2} \right)$$

$$= n \left( \frac{3n+1}{2} \right)$$

$$= \frac{n(3n+1)}{2}$$

②  $15 + n$

$$S_n = \sum 15 + \sum n$$

$$= 15n + \frac{n(n+1)}{2}$$

$$= \frac{30n + n^2 + n}{2}$$

$$= \boxed{\frac{n(n+3)}{2}}$$

③  $2n^2 + 3n$

$$S_n = 2\sum n^2 + 3\sum n$$

$$= \frac{2n(n+1)(2n+1)}{6} + \frac{3n(n+1)}{2}$$

$$= \frac{n+1}{2} \left( \frac{2n(2n+1)}{3} + 3n \right)$$

$$= \frac{n+1}{2} \left( \frac{4n^2+2n+9n}{3} \right)$$

$$= \frac{n+1}{6} (4n^2 + 11n)$$

$$= \boxed{\frac{n(n+1)(4n+11)}{6}}$$

$$\textcircled{4} \quad T_n = 3^n - 2^n$$

$$S_n = \sum (3^n - 2^n)$$

$$= \sum 3^n - \sum 2^n$$

$$= \cancel{2(3+9+27+\dots 3^n)} - \cancel{2(2+4+8+16+\dots 2^n)}$$

$$= (3 + 3^2 + 3^3 + 3^4 + \dots 3^n) - (2 + 2^2 + 2^3 + 2^4 + \dots 2^n)$$

$$= \frac{3(3^n - 1)}{2} - \frac{2(2^n - 1)}{1}$$

$$= \frac{3(3^n - 1)}{2} - 4(2^n - 1)$$

$$= \frac{3^{n+1} - 3 - 2^{n+2} + 4}{2} = \boxed{\frac{3^{n+1} - 2^{n+2} + 1}{2}}$$

$$\textcircled{5} \quad T_m = 3^k + k \cdot k^3$$

$$S_n = \sum 3^k + k \cdot k^3 = \sum k^3$$

$$= (3 + 3^2 + 3^3 + \dots 3^k) + (\cancel{1+8+27} (1^3 + 2^3 + 3^3 + \dots n^3))$$

$$= \left[ \frac{n(n+1)}{2} \right]^2 + \frac{3(3^n - 1)}{2}$$

$$= \frac{[n(n+1)]^2 + 6(3^n - 1)}{4}$$

14. W. 04-07-2024

DYS-7 (Q2)

Q Find the sum of following series.

①  $1 \times 2 + 2 \times 3 + 3 \times 4 + 4 \times 5 + \dots n \text{ terms}$

$$1, 2, 3, 4, \dots$$

$$T_m = m$$

$$2, 3, 4, \dots$$

$$T_m = m+1$$

$$T_m = n(n+1)$$

$$\begin{aligned} S_n &= \sum T_m \\ &= \sum n^2 + \sum n \\ &= \frac{n(n+1)(2n+1)}{6} + \frac{n(n+1)}{2} \end{aligned}$$

$$= \frac{n(n+1)(2n+1+3)}{6}$$

$$= \frac{n(n+1)(n+2)}{3}$$

②.

$$1 \cdot 2 \cdot 3 + 2 \cdot 3 \cdot 4 + 3 \cdot 4 \cdot 5 + \dots n$$

$$1, 2, 3, 4, \dots n$$

$$T_m = m$$

$$2, 3, 4, 5, \dots n$$

$$T_m = m+1$$

$$3, 4, 5, 6, \dots n$$

$$T_m = m+2$$

$$T_m = m(m+1)(m+2)$$

$$S_n = \sum (m^3 + m^2 + 2m)$$

$$S_n = \sum (m^3 + 3m^2 + 2m)$$

$$S_n = \sum m^3 + 3\sum m^2 + 2\sum m$$

$$\left. \begin{aligned} S_n &= \frac{n(n+1)^2}{24} + \frac{3n(n+1)(2n+1)}{62} + \frac{2n(n+1)}{2} \\ S_n &= \frac{n^2(n+1)^2}{4} + \frac{2n(n+1)(2n+1)}{4} + \frac{4n(n+1)}{4} \\ S_n &= n(n+1) \left[ n(n+1) + 2(2n+1) + 4(n+1) \right] \end{aligned} \right\} 4$$

$$Q3. 1 \cdot 3 \cdot 5 + 3 \cdot 5 \cdot 7 + 5 \cdot 7 \cdot 9 + \dots n$$

~~+ 3 + 5~~

1, 3, 5, ... n

$$\text{S.T. } T_n = 2n - 1$$

3, 5, 7, ... n

$$T_n = 2n + 1$$

5, 7, 9, ...

$$T_n = 2n + 3$$


$$T_n = \frac{(2n-1)(2n+1)(2n+3)}{6}$$

$$= (4n^2 - 1)(2n+3)$$

$$= 8n^3 + 12n^2 - 2n - 3$$

$$S_n = 8 \sum n^3 + 12 \sum n^2 - 2 \sum n - 3$$

$$Q) 1^2 + \frac{1^2 + 2^2}{2} + \frac{1^2 + 2^2 + 3^2}{3} + \frac{1^2 + 2^2 + 3^2 + 4^2}{4} + \dots n \text{ times}$$



$$\frac{1}{1}, \frac{1}{2}, \frac{1}{3}, \frac{1}{4} (\text{H.P.})$$

$$T_n = \frac{1}{n}$$

$$1^2, 1^2 + 2^2, 1^2 + 2^2 + 3^2, \dots$$

$$T_n = \frac{\sum n^2}{6}$$

$$T_n = \frac{n(n+1)(n+2+1)}{6}$$

$$T_n = \frac{(n+1)(2n+1)}{6}$$

$$= \frac{2n^2 + 3n + 1}{6}$$

$$S_n = \frac{1}{6} \left[ 2 \sum n^2 + 3 \sum n + n \right]$$

$$= \frac{\sum n^2}{3} + \frac{\sum n}{2} + \frac{n}{6}$$

$$⑤ 1 + \frac{1^2 + 2^2}{1+2}, 1^2 + \frac{2^2 + 3^2}{1+2+3} + \frac{1^2 + 2^2 + 3^2 + 4^2}{1+2+3+4} \dots$$

$$\frac{1}{1}, \frac{1}{1+2}, \frac{1}{1+2+3} \quad (H.P)$$

$$1, 1+2, 1+2+3 \quad (A.P)$$

$$T_m = \sum n$$

$$T_m = \frac{1}{\sum n} \quad (H.r)$$

$$1 + 1^2 + 2^2, 1^2 + 2^2 + 3^2$$

$$T_m = \sum n^2$$

$$T_m = \frac{\sum n^2}{\sum n}$$

$$= \frac{n(n+1)(2n+1)}{6 \cdot 3} \times \frac{2}{n(n+1)}$$


$$= \frac{2n+1}{3}$$

$$S_m = \frac{1}{3} [2\sum n + n]$$

$$= \frac{1}{3} [\sum n^2 + n]$$

$$= \frac{n^2 + 2n}{3}$$

## Type-3 :- Method of Difference



Q)  $3 + 7 + 13 + 21 + 31 + \dots$  n terms.

$\begin{matrix} 3 & 7 & 13 & 21 & 31 \\ 4 & 6 & 8 \\ 2 & 2 \end{matrix}$  → constant A.P.

$$T_n = an^2 + bn + c$$

$$n=1; 3 = a + b + c - \textcircled{1}$$

$$n=2; 7 = 4a + 2b + c - \textcircled{2}$$

$$n=3; 13 = 9a + 3b + c - \textcircled{3}$$

$$a = b = c = 1$$

$$T_n = n^2 + n + 1$$

$$\begin{aligned} S_n &= \sum T_n \\ &= \sum n^2 + \sum n + \sum 1 \\ &= \frac{n(n+1)(2n+1)}{6} + \frac{3n(n+1)}{6} + \frac{6n}{6} \end{aligned}$$

$$= \frac{2n^3 + 3n^2 + n + 3n^2 + 3n + 6n}{6}$$

$$= \frac{2n^3 + 6n^2 + 10n}{6}$$

$$= \boxed{\frac{n^3 + 3n^2 + 5n}{3}}$$

Q2.  $1 + \underbrace{q + 10 + \dots}_{\text{n terms}} + 22 + \dots$

~~13~~  $\underbrace{6}_{3} \quad 12 \rightarrow GP$

~~2762~~  $\rightarrow$  constant of

$T_n = ar^{n-1} + b$

~~$a = 6$~~

$q = ar + b \rightarrow q = 6r$

$10 = ar^2 + b \rightarrow 10 = ar^2 \rightarrow \frac{5}{2} = r$

~~$a = \frac{8}{5}$~~

$T_n = \frac{8}{5} \times \left(\frac{5}{2}\right)^{n-1} + 1$

$T_n = \frac{8}{5}$

$1 = 6 + a$

$4 = a(2) + b \rightarrow a = \frac{3}{2}$

$q = a2 + b$

~~$3 = a$~~

~~$b = -2$~~

~~$T_n = 3 \times \frac{2^n}{2} - 2$~~

~~$T_n = 3 \cdot 2^{n-1} - 2$~~

~~$S_n = 3 \sum 2^{n-1} - 2n$~~

~~$= 3(1 + 2 + 4 + \dots + n) - 2n$~~

~~$= 3(\frac{2^n - 1}{2}) - 2n$~~

~~$= 3 \cdot 2^n - 3 - 2n$~~

$$S_n = 3 \sum 2^{n-1} - 2n$$

$$= \frac{3}{2} (2 + 4 + 8 \cdot 2^n) - 2n$$

$$= \frac{3}{2} \times \frac{2(2^n - 1)}{2} - 2n$$

$$= 3(2^n - 1) - 2n$$

$$Q) 6 + 13 + 22 + 33 \dots n$$

$$\begin{array}{cccc} 6 & 13 & 22 & 33 \\ 7 & 13 & 21 & \\ 2 & 2 & & \end{array}$$

$$T_n = an^2 + bn + c$$

$$6 = a + b + c$$

$$13 = 4a + 2b + c$$

$$22 = 9a + 3b + c$$

$$9 = 5a + b \quad | \quad \boxed{b=4}$$

$$7 = 3a + b \quad | \quad \boxed{C=1}$$

$$2 = 2a$$

$$\boxed{a=1}$$

$$T_n = n^2 + 4n + 1$$

$$S_n = \sum n^2 + 4n + 1 \cdot n$$

$$= n \left( \frac{(n+1)(2n+1)}{6} + \frac{4n(n+1)}{2} + n \right)$$

$$Q) 9 + 16 + 25 + 36 + 49 + 64 \dots n \text{ terms}$$

$$\begin{array}{cccc} 9 & 16 & 25 & 36 & 49 \\ 7 & 13 & 21 & 29 & 37 \\ 16 & 12 & 24 & & \end{array}$$

→ 2nd Order G.P

$$T_n = an^{n-1} + bn + c$$

$$a = 2$$

$$9 = a + b + c$$

$$16 = 2a + 2b + c$$

$$25 = 4a + 3b + c$$

$$36 = 9a + 6b + c$$

$$7 = a + b$$

$$\boxed{b=0}$$

$$\boxed{b=1}$$

$$\boxed{c=2}$$

$$T_n = 6 \times 2^{n-1} + n + 2$$

$$S_n = 13 \sum (2 + 4 + 6 \dots n) + \frac{n(n+1)}{2} + 2n$$

$$S_n =$$

Type -3

Splitting in denominator

Splitting the  $n^{\text{th}}$  term (denominator)

$$Q \quad \frac{1}{1 \cdot 2} + \frac{1}{2 \cdot 3} + \frac{1}{3 \cdot 4} + \frac{1}{4 \cdot 5} \dots n$$

$$1 + 2 + 3 + 4 + 5 + \dots n$$

$$2, 3, 4, 5, 6$$

$$T_n = n+1$$

$$T_1 = n(n+1) (\cancel{n})$$

$$T_n = \frac{1}{n(n+1)} (\cancel{n})$$

$$T_n = \left( \frac{n+1-n}{n(n+1)} \right)$$

$$= \frac{n+1}{n(n+1)} - \frac{n}{n(n+1)}$$

$$= \frac{1}{n} - \frac{1}{n+1}$$

$$n = 1$$

$$\frac{1}{1} - \frac{1}{2}$$

$$n = 2$$

$$\frac{1}{2} - \frac{1}{3}$$

$$n = 3$$

$$\frac{1}{3} - \frac{1}{4}$$

$$n = n \quad \cancel{\frac{1}{n}} - \frac{1}{n+1}$$

$$S_n = 1 - \frac{1}{n+1}$$

$$= \frac{n}{n+1}$$

$$Q \quad \frac{1}{1 \cdot 4} + \frac{1}{2 \cdot 3} + \frac{1}{7 \cdot 10} + \dots n^{\text{terms}}$$

$$\begin{aligned} T_n &= n(n+3) \\ &= n(n+3) \\ T_n &= \frac{1}{n(n+3)} \\ &= \frac{(n+3)-n}{n(n+3)} \\ &= \frac{1}{n} - \frac{1}{n+3} \\ S_n &= \frac{1}{1} - \frac{1}{n+3} \\ &= \frac{n+3-1}{n+3} \\ &= \boxed{\frac{n+2}{n+3}} \end{aligned}$$

$$\begin{aligned} T_n &= (3n-2)(3n+1) \\ &= \frac{1}{3} \times \frac{3}{(3n-2)(3n+1)} \\ &= \frac{1}{3} \left[ \frac{(3n+1)(-3n-2)}{(3n-2)(3n+1)} \right] \\ &= \frac{1}{3} \left( \frac{3n+1 - 3n-2}{(3n-2)(3n+1)} \right) \end{aligned}$$

$$\boxed{\frac{1}{(3n-2)(3n+1)}}$$

$$S_n = \frac{9n+2}{3n(3n+1)}$$

H.W. 05-07-2024

$$\text{DXS } 7 \{1, 4, 5, 6\} \{3, 7\}$$

$$01 \{27, 28, 30\}$$

$$1^2 - 2^2 + 3^2 - 4^2 + 5^2 - 6^2 + 7^2 - 8^2 + \dots + 99^2 - 100^2.$$

$$\begin{aligned} Q \quad 1^2 - 2^2 + 3^2 - 4^2 + 5^2 - \dots 99^2 - (2^2 + 4^2 + 6^2 + \dots 100^2) \\ T_n = \frac{(2n-1)^2}{6} \quad T_n = \frac{(2n)^2}{4n^2} \\ = 4n^2 - 4n + 1 \end{aligned}$$

$$S_n = 4n \frac{(n+1)(2n+1)}{6} - \frac{4n(n+1)}{2} + n$$

$$S_n = \frac{4n(n+1)(2n+1)}{6}$$

$$S_n = \frac{200(51)(101)}{6} - \frac{200(51)}{2} + 50$$

$$S_n = \frac{200(51)(101)}{6}$$

$$S_n = 100 \times 17 \times 101 - 5100 + 50$$

$$S_n = 100 \times 101 \times 17$$

$$S_n = 171700 - 5100 + 50$$

$$S_n = 171700$$


$$\boxed{S_n = -5050}$$

Q Sum of  $\infty$  terms ..

$$\frac{1}{1 \cdot 4} + \frac{1}{4 \cdot 7} + \frac{1}{7 \cdot 10}$$

$$1 + 4 + 7 \dots \dots \dots$$
$$T_n = 1 + (n-1)3$$
$$= 1 + 3n - 3$$
$$= 3n - 2$$

$$4 + 7 + 10$$

$$T_n = 3n + 1$$


$$T_n = (3n+2)(3n+1)(A.P)$$

$$S_n = \frac{n}{3n+1}$$

$$S_n = \frac{\frac{n}{n}}{\frac{3n+1}{n}}$$
$$= \frac{n}{3+\frac{1}{n}}$$
$$\boxed{= \frac{1}{3}}$$

$$\frac{1}{\infty} = 0$$

Q)  $T_n = \frac{n}{1-n^2}$

$$= \frac{n}{(1+n)(1-n)}$$
$$= \frac{(1-n) - (1+n)}{-2(1+n)(1-n)}$$

$$= \frac{-1}{2(1+n)} + \frac{1}{2(1-n)}$$

$$= \frac{1}{2(1-n)} - \frac{1}{2(1+n)}$$

start,  $n=2$

$$S_n = \frac{-1}{4(2)} + \frac{1}{2(2-n)}$$

$$= \frac{2n-2+1}{4(2-n)}$$

~~$$= \frac{n+1}{4(2-n)}$$~~

$$\boxed{= \frac{2n}{2(1-n)}}$$

Q Find sum of the sequence upto infinite terms:-

$$\frac{1}{1+1^2+1^4} + \frac{2}{1+2^2+2^4} + \frac{3}{1+3^2+3^4}$$

$$(1+1^2+1^4), (1+2^2+2^4), (1+3^2+3^4)$$

$$T_n = \frac{n(n^3-1)}{n-1}$$

$$1, 2, 3$$

$$T_n = n$$

$$\begin{aligned} T_n &= n \times \frac{(n-1)}{n(n^3-1)} \\ &= \frac{n-1}{n^3-1} \\ &= \frac{n-1}{(n-1)(n^2+n+1+2n)} \\ T_n &= \frac{1}{n^2+2n+1} \end{aligned}$$

~~$$T_n = \frac{n}{1+n^2+n^4}$$~~

~~$$T_n = \frac{n}{n^2+n^3}$$~~

~~$$T_n = \frac{n}{(n^4+n^2+1)}$$~~

$$\begin{aligned} T_n &= \frac{n}{(n^2+n+1)(n^2-n+1)} \\ &= \frac{(n^2-n+1) - (n^2+n+1)}{2(n^2+n+1)(n^2-n+1)} \end{aligned}$$

$$T_n = \frac{1}{2(n^2+n+1)} - \frac{1}{2(n^2-n+1)}$$

~~$$n=1; \frac{1}{6} - \frac{1}{2}$$~~

~~$$n=2; \frac{1}{14} - \frac{1}{6}$$~~

$$S_n = \frac{1}{2} \left[ 1 - \frac{1}{n^2+n+1} \right]$$

$$S_n = \frac{1}{2} \left[ \frac{n^2+n+1-1}{n^2+n+1} \right]$$

$$= \frac{1}{2} \left[ \frac{n^2+n}{n^2+n+1} \right]$$

$$= \frac{1}{\frac{2}{n^2+n} + \frac{1}{n^2+n}}$$

$$= \frac{1}{2}$$

Type-1 Using Difference in numerator

Q  $1 \cdot 2 + 2 \cdot 3 + 3 \cdot 4 + 4 \cdot 5 + \dots n \text{ terms}$

$$T_n = n(n+1)$$

$$\begin{aligned} T_n &= \frac{1}{3} n(n+1) [(n+2) - (n-1)] \\ &= \frac{n(n+1)(n+2)}{3} - \cancel{\frac{(n-1)(n)(n+2)}{3}} \end{aligned}$$

$$n=1; \cancel{\frac{1}{3} \cdot 1 \cdot 2 \cdot 3 - \frac{1}{3} \cdot 0} = 0$$

$$n=2; \cancel{\frac{1}{3} \cdot 2 \cdot 3 \cdot 4 - \frac{1}{3} \cdot 1 \cdot 2 \cdot 3} = 10$$

$$n=n \quad \frac{n(n+1)(n+2)}{3} = 0$$

$$S_n = \frac{n(n+1)(n+2)}{3}$$

Q  $1 \cdot 2 \cdot 3 + 2 \cdot 3 \cdot 4 + 3 \cdot 4 \cdot 5 + \dots n \text{ terms.}$

$$T_n = n(n+1)(n+2)$$

$$T_n = \frac{n(n+1)(n+2)}{4} \cancel{[(n+3) - (n-1)]}$$

$$T_n = \frac{n(n+1)(n+2)(n+3)}{4} - \frac{(n-1)(n)(n+1)(n+2)}{4}$$

$$n=1; \cancel{\frac{1 \cdot 2 \cdot 3 \cdot 4}{4}} = 0$$

$$n=2; \frac{2 \cdot 3 \cdot 4 \cdot 5}{4} - \cancel{\frac{1 \cdot 2 \cdot 3 \cdot 4}{4}}$$

$$S_n = \frac{n(n+1)(n+2)(n+3)}{4}$$

$$Q) 1+3+5+\dots+2n-1 = n^2$$

$$T_n = \frac{n(n+1)(n+2)}{6}$$

$$T_n = \frac{(2n-1)(2n+1)(2n+3)}{6}$$

$$T_n = \frac{(2n-1)(2n+1)(2n+3)}{6} \quad \text{or } 8$$

$$T_n = \frac{8(2n-1)(2n+1)(2n+3)(2n+5)}{6 \cdot 8} = \frac{(2n+3)(2n-1)(2n+1)(2n+3)}{6 \cdot 8}$$

$$n=1; \quad 1+3+5+7+9 = 25$$

$$n=2; \quad 3+5+7+9 = 25$$

$$S_n = \frac{(2n-1)(2n+1)(2n+3)(2n+5)}{6}$$

Q) PYS 7  
06 ii)

$$\sum_{n=1}^{\infty} n(n+1)(n+2)(n+3)$$

$$T_n = n(n+1)(n+2)(n+3)$$

$$T_n = (n+1)(n)(n+1)(n+2)(n+3) - (n-1)(n)(n+1)(n+2)(n+3)$$

$$n=1;$$

$$S_n = 22 \left\{ \frac{n(n+1)(n+2)(n+3)(n+4)}{5} \right\}$$

$$Q \quad S_p = 1 + 2 + 3 + \dots + p \text{ terms}$$

$$\sum_{p=1}^n = \frac{1}{8} S_p$$

$$T_n = \frac{1}{\sum n}$$

$$= \frac{2}{n(n+1)}$$

$$T_n = \frac{2[(n+1) - n]}{n(n+1)}$$

$$T_n = \frac{2}{n} - \frac{2}{n+1}$$

$$n=1; \quad 2 - \frac{2}{2}$$

$$n=2; \quad \cancel{\frac{2}{2}} - \frac{2}{3}$$

$$n=3; \quad \cancel{\frac{2}{3}} - \frac{2}{4}$$

$$S_n = 2 - \frac{2}{n+1}$$

$$= 2n + 2 - 2$$

$$= \frac{2n}{n+1}$$

$$S_n = \frac{2n}{n+1}$$

$$Q \quad 1 + \frac{(1+2)^2}{1+3} + \frac{1(1+2+3)^2}{1+3+5} + \frac{1(1+2+3+4)^2}{1+3+5+7} + \dots \dots \dots \text{Ans}$$

O-1

$$Q \quad 16-30 = \{27, 29, 30\}$$

O-2

$$Q \quad 1-15 (16-2)$$

O-3

$$Q \quad 1-10$$

$$T_m = \frac{(\sum n)^2}{\cancel{\sum} (2m-1)}$$

$$T_m = \frac{n(n+1)}{2}^2$$

$$= \frac{n^2(n^2+1+2n)}{4n^2}$$

$$T_n = \frac{n^2+2n+1}{4}$$

~~Rei~~

$$S_m = \frac{1}{4} \left[ \frac{n(n+1)(2n+1)}{6} + \frac{2(n+1)n}{2} + n \right]$$

$$= \frac{1}{4} \left[ \frac{n(n+1)(2n+1) + 6n(n+1) + 6n}{6} \right]$$

$$= \frac{n}{24} [(n+1)(2n+1+6) + 6]$$

$$\boxed{S_n = \frac{n}{24} [(n+1)(2n+1+6) + 6]}$$

①.  $S_n = n(n+1)$  Find ①  $\frac{1}{T_m}$  ②  $\sum_{n=1}^{10} \frac{1}{T_m}$

$$S_n = \sum T_n = n(n+1)$$

$$T_n = 2^n$$

$$\boxed{\frac{1}{T_m} = \frac{1}{2^n}}$$

$$\frac{1}{2} + \frac{1}{4} + \frac{1}{8} \dots$$

~~$$\frac{1}{T_m} = \frac{1}{2^n}$$~~

$$S_m = \frac{1}{2} \left( 1 - \left( \frac{1}{2} \right)^{10} \right)$$

$$= 1 - \frac{1}{1024}$$

$$\boxed{= \frac{1023}{1024}}$$

$$Q2. \frac{S}{1^2 \cdot 4^2} + \frac{11}{4^2 \cdot 7^2} + \frac{17}{7^2 \cdot 10^2} \dots$$

$$T_m = S + (n-1)6 \dots \\ = S + 6n - 6$$

$$\boxed{= 6n-1}$$

$$T_m = 1^2 + 4^2 + 7^2 + \dots \\ = (1 + (n-1)3)^2 \\ = (1 + 3n - 3)^2 \\ = (3n - 2)^2$$

$$T_m = 4^2 + 7^2 + 10^2 \dots \\ = (3n+1)^2 \\ = 9n^2 + 1 + 6n \\ = 9n^2 + 6n + 1$$

$$T_m = \frac{6n-1}{(9n^2-6n+4)(9n^2+6n+1)}$$

$$S = \frac{(1+1)}{4^2 \cdot 1^2} + \frac{(7+1)}{7^2 \cdot 4^2} + \frac{(10+1)}{10^2 \cdot 7^2} \\ = \frac{1}{3} \left[ \frac{(4+1)(4-1)}{4^2 \cdot 1^2} + \frac{(7+1)(7-1)}{7^2 \cdot 4^2} + \frac{(10+1)(10-1)}{10^2 \cdot 7^2} \right] \\ = \frac{1}{3} \left[ \frac{4^2 - 1^2}{4^2 \cdot 1^2} + \frac{7^2 - 4^2}{7^2 \cdot 4^2} + \frac{10^2 - 7^2}{10^2 \cdot 7^2} \right] \\ = \frac{1}{3} \left[ \frac{1}{4^2} - \frac{1}{1^2} + \frac{1}{4^2} - \frac{1}{7^2} + \frac{1}{7^2} - \frac{1}{10^2} \right]$$

$$\boxed{= \frac{1}{3}}$$

$$\text{Q} (1^2 - a_1) + (2^2 - a_2) + (3^2 - a_3) + \dots + (n^2 - a_n) = \frac{1}{3} n (n^2 - 1)$$

$$a_1 = ?$$

$$\text{S}_n - \text{S}_{n-1}$$

$$0 + \underbrace{2 + 6 + 12}_{\begin{matrix} 2 & 4 \\ 2 & 2 \end{matrix}}$$

$$T_n = an^2 + bn + c$$

$$0 = a + b + c$$

$$2 = 4a + 2b + c$$

$$6 = 9a + 3b + c$$

$$4 = 5a + b$$

$$2 = 3a + b$$

$$2 = 2a$$

$$a = 1$$

$$b = -1$$

$$c = 0$$

$$T_n = n^2 - n$$

$$1 - a_1 = 0$$

$$a_1 = 1$$

$$4 - a_2 = 2$$

$$a_2 = 2$$

$$9 - a_3 = 6$$

$$a_3 = 3$$

$$n^2 - a_n = n^2 + n$$

$$a_n = n$$

$$a_7 = 7$$







