Exploring The Most Beautiful Mathematical Formula

$$e^{i\pi} = -1$$

Shreya Balaji

Berkeley Carroll School

May 1, 2020

Why is the Equation Beautiful

Euler's identity is beautiful because it combines five of the most important numbers in mathematics into one equation. These numbers are...

- 1 the basis for all other numbers
- 0 the concept of nothingness
- pi the ratio of a circle's circumference to its diameter
- e the base of natural logarithms
- i the "imaginary" square root of -1

History of Mathematicians Involved

A number of mathematicians laid the building blocks for understanding this equation. As you can see, it has taken painstaking work of many mathematicians for us to come to such an elegant equation. Below are are some of the mathematicians whose works contributed to this equation :

```
Jacob Bernoulli - Swiss mathematician Invented e
Gottfried Wilhelm Leibniz - Along with Newton invented Calculus
Issac Newton - Along with Leibniz invented Calculus
Carl Friedrich Gauss - One of greatest mathematician of all time - Imaginary
numbers
Leonhard Euler - Worked on imaginary numbers
Augustin-Louis Cauchy - Worked on imaginary numbers
Greek, Chinese and Indian Mathematicians - Contributed to \pi
James Gregory, Brook Taylor and Colin Maclaurin - Contributed to
understanding of expansion series
```

Taylor and Maclaurin Series

James Gregory and Brook Taylor invtented the idea of infinite series expansion of functions. Scottish mathematician Colin Maclaurin used the work extensively to study functions centered around zero. The basic idea of Maclaurin series is that any function can be expanded into a infinite series centered around zero as given below:

$$f(x) = f(0) + x \cdot \frac{f'(0)}{1!} + x^2 \cdot \frac{f''(0)}{2!} + x^3 \cdot \frac{f'''(0)}{3!} + \dots$$

In the next three slides, we will see the application of the same for three basic functions:

$$e(x) \dots \sin(x) \dots \cos(x)$$



Taylor Series for e(x)

f(x)	f'(x)	f''(x)	f'''(x)	f''''(x)
e^x	e^x	e^x	e^x	e^x
1	1	1	1	1

Applying the Maclaurin formula we see that the infinite series expansion for e^{x} is :

$$f(x) = f(0) + x \cdot \frac{f'(0)}{1!} + x^2 \cdot \frac{f''(0)}{2!} + x^3 \cdot \frac{f'''(0)}{3!} + \dots$$
$$e^x = 1 + x \cdot \frac{1}{1!} + x^2 \cdot \frac{1}{2!} + x^3 \cdot \frac{1}{3!} + \dots$$

By extension:

$$e^{ix} = 1 + i \cdot x \cdot \frac{1}{1!} - x^2 \cdot \frac{1}{2!} - i \cdot x^3 \cdot \frac{1}{3!} + x^4 \cdot \frac{1}{4!} \dots$$

Taylor Series for sin(x)

f(x)	f'(x)	f''(x)	f'''(x)	f''''(x)
$\sin(x)$	$\cos(x)$	$-\sin(x)$	$-\cos(x)$	$\sin(x)$
0	1	0	-1	0

Applying the Maclaurin formula we see that the infinite series expansion for e^{x} is :

$$f(x) = f(0) + x \cdot \frac{f'(0)}{1!} + x^2 \cdot \frac{f''(0)}{2!} + x^3 \cdot \frac{f'''(0)}{3!} + \dots$$
$$\sin(x) = 0 + x \cdot \frac{1}{1!} - x^3 \cdot \frac{1}{3!} + x^5 \cdot \frac{1}{5!} + \dots$$

Taylor Series for cos(x)

f(x)	f'(x)	f''(x)	f'''(x)	f''''(x)
$\cos(x)$	$-\sin(x)$	-cos(x)	$\sin(x)$	$\cos(x)$
1	0	-1	0	1

Applying the Maclaurin formula we see that the infinite series expansion for e^{x} is :

$$f(x) = f(0) + x \cdot \frac{f'(0)}{1!} + x^2 \cdot \frac{f''(0)}{2!} + x^3 \cdot \frac{f'''(0)}{3!} + \dots$$
$$\cos(x) = 1 - x^2 \cdot \frac{1}{2!} + x^4 \cdot \frac{1}{4!} + \dots$$

Bringing it all together

8/9

A Graphical Representation