



EVALUATED PHASE DIAGRAMS OF BINARY METAL-TELLURIUM SYSTEMS  
OF THE D-BLOCK TRANSITION ELEMENTS

*by*

G. Chattopadhyay and S. R. Bharadwaj  
Water Chemistry Division

1989

B.A.R.C. - 1449

B.A.R.C. - 1449

GOVERNMENT OF INDIA  
ATOMIC ENERGY COMMISSION

**EVALUATED PHASE DIAGRAMS OF BINARY METAL-TELLURIUM SYSTEMS  
OF THE D-BLOCK TRANSITION ELEMENTS**

by

G. Chatopadhyay and S.R. Bharadwaj  
Water Chemistry Division

BHABHA ATOMIC RESEARCH CENTRE  
BOMBAY, INDIA

1989

### 13. SYSTEM MO-TE

**Liquid Phase:** No liquid phase in this system upto 1 atm of Te vapor is known.

**Terminal Solid Solution:** The solid solubility of Te in Mo at 1373 K is less than 1-2 at.% (1). The molybdenum solvus is given by  $\ln X_{Te} = 6600/T$  between 700 and 1573 K (2). The solid solubility of Mo in Te must be negligible in view of the low solubility of Mo in liquid Te.

**Intermediate Phases:** There are two intermediate phases  $Mo_3Te_4$  (2,3) and  $MoTe_2$  (2-12). The ranges of homogeneity of  $Mo_3Te_4$  (3,12) and of  $MoTe_2$  (3,9,12) have been assessed by (13) as follows. The Mo-rich boundary of the  $Mo_3Te_4$  phase ranges from 56.5 at.% Te at low temperatures to 55.9 at.% Te at 1570 K. The Te-rich boundary ranges from 57.4 at.% Te at low temperatures to 58.3 at.% Te at 1450 K. The  $Mo_2Te_3$  phase of (4) and (7) probably belongs to the  $Mo_3Te_4$  phase. The Mo-rich boundary of  $MoTe_2$  is nearly independent of temperature at 65.5 at.% Te. The Te-rich boundary shifts from 65.5 at.% Te at 1450 K to 66.6 at.% Te at 1260 K (9,12).  $MoTe_2$  undergoes phase transition from hexagonal to monoclinic form, the transition temperature varying from 1153 K at the Mo-rich end to 1093 K at the Te-rich end.

**Crystallographic Data:** The crystallographic data along with the homogeneity ranges are given in Table 13 A according to the compilation by (13).

**Phase Diagram:** The phase diagram has been evaluated by (13). The isobaric phase diagram at 1 atm. based on the dissociation pressure data of (14) is given in Fig. 13. (13) surmised in analogy to Mo-S and Mo-Se systems (15), Mo-Te system should show Mo- $Mo_3Te_4$  and  $Mo_3Te_4$ - $MoTe_2$  eutectics around 1673 K with congruent melting points of 1773 to 1873 K at high pressures.

TABLE 13 A

Crystal Structures of the Intermediate Phases in the System Mo-Te

Phase designation	Composition range $X_{Te}$	Structure	Space group	Lattice parameter nm
$Mo_3Te_4$	0.565-0.574 (low temperature)	hexagonal	$R\bar{3}$	$a=1.013$ ; $c=1.170$
	0.559-0.583 (1500 K)			
	0.555-0.608 (1273 K) (3)	monoclinic	-	$a=0.4875$ ; $b=0.5092$ $c=0.7051$ ; $\beta=93.75^\circ$ (16)
	0.539-0.593 (933 K) (3)			
$\alpha-MoTe_2$	0.655-0.666 (1260 K) (9,11)	hexagonal ( $MoS_2$ )	$P6_3/nmc$ (hP6)	$a=0.35182$ ; $c=1.39736$ (17)
	0.652-0.667 (933 K) (3)			
$\beta-MoTe_2$ (high temp)	0.649-0.661 (1273 K) (3)	monoclinic	$P2/m$ (mP12)	$a=0.633$ ; $b=0.3469$ $c=1.386$ ; $\beta=93^\circ 55'$




Fig.13. Phase Diagram of the Molybdenum-Tellurium System.

#### References

1. M.E. Straumanis and R.P. Shodhan, *Z. Metallkde.* **59** (1968) 492.
2. A.A. Opalovsky and V.E. Fedorov, *Izv. Akad. Nauk. SSSR, Neorg. Mater.* **2** (1966) 443; *Dokl. Akad. Nauk SSSR*, **163** (1965) 1163
3. M. Spiesser, J. Rouxel, M. Keriou and M.G. Goureaux, *Bull. Soc. Chim. Fr.* No. 5 (1969) 1427.
4. A. Morrette, *Ann. Chim.* **19** (1944) 130.
5. H. Schafer, *Z. Anorg. Chem.* **286** (1956) 27, 42; **290** (1957) 279; **291** (1957) 221, 254.

6. R.Nitsche, H.U.Bolsterlt and M.Lichensteiger, *J.Phys.Chem.Solids* 21 (1961) 199.
7. D.Puotinen and R.E.Newnham, *Acta Cryst.* 14 (1961) 691.
8. L.H.Brixner, *J.Inorg.Nucl.Chem.* 24 (1962) 257.
9. J.Lepetit, *J De Phys.* 26 (1965) 175.
10. E.Revolinsky and D.J.Berntsen, *J.Phys.Chem.Solids*, 27 (1966) 175.
11. B.E.Brown, *Acta Cryst.* 20 (1966) 268.
12. M.B.Vellinga, R.De Jonge and C.U.Haas, *J.Solid State Chem.* 2 (1970) 299
13. L.Brewer and R.H.Lamoureaux, *Atomic Energy Review, Molybdenum: Physico Chemical Properties of its Compounds and Alloys, Special Issue No.7*  
Ed.L.Brewer, IAEA, Vienna, 1980, p329.
14. A.Opalovsky, V.E.Fedorov, E.U.Lobokov and B.I.Tsikanovskij,  
*Zh.Fiz.Khim.* 45 (1971) 1864
15. R.Flukiger, H.Devantay, J.L.Jordan and J.Muller, *J.Inst.Electr.*  
*Electron.Eng.Trans.Mag.* 13 (1977) 818.
16. R.De Jonge, T.J.A.Popma, G.A.Wiegers, and F.Jellinek, *J.Solid State Chem.*,  
2 (1970) 118
17. O.Knop and R.D.McDonald, *Can.J.Chem.*, 39 (1961) 897