


- 1 A particle P of mass m is attached to one end of a light inextensible string of length a . The other end of the string is attached to a fixed point O . The particle P is held at the point A , where OA makes an angle α with the downward vertical through O , and with the string taut. The particle P is projected perpendicular to OA in an upwards direction with speed $\sqrt{3ag}$. It then starts to move along a circular path in a vertical plane. The string goes slack when P is at B , where OB makes an angle θ with the upward vertical.

Given that $\cos \alpha = \frac{4}{5}$, find the value of $\cos \theta$.

[4]



Let u be speed @ bottom

@A:

$$\frac{mv^2}{r} = \frac{mu^2}{r} - 2mg + 2mg \cos(\alpha)$$

$$\frac{m(3ga)}{a} = \frac{mu^2}{a} - 2mg + \frac{8mg}{5}$$

$$3g = \frac{u^2}{a} - 2g + 1.6g$$

$$u^2 = 3.4ga$$

@B:

$$T = \frac{mv^2}{r} - 2mg + 3mg \cos(180 - \theta)$$

$$\frac{m(3.4ga)}{a} = 2mg - 3mg \cos(180 - \theta)$$

$$3.4 = 2 - 3 \cos(180 - \theta)$$


$$\cos(180 - \theta) = -\frac{7}{15}$$

$$\cos \theta = \frac{7}{15}$$

- 2 One end of a light elastic string, of natural length a and modulus of elasticity λmg , is attached to a fixed point O . The string lies on a smooth horizontal surface. A particle P of mass m is attached to the other end of the string. The particle P is projected in the direction OP . When the length of the string is $\frac{4}{3}a$, the speed of P is $\sqrt{2ag}$. When the length of the string is $\frac{5}{3}a$, the speed of P is $\frac{1}{2}\sqrt{2ag}$.

Find the value of λ .

[4]



Energy in ①:

$$\begin{aligned} & KE + EPE \\ &= \frac{1}{2}(2ga)(m) + \frac{(\lambda mg)(a/3)^2}{2a} \\ &= mga + \frac{\lambda mga}{18} \end{aligned}$$

Energy in ②:

$$\begin{aligned} & KE + EPE \\ &= \frac{1}{2}m\left(\frac{1}{2}\sqrt{2ag}\right)^2 + \frac{\lambda mg\left(\frac{2a}{3}\right)^2}{2a} \\ &= \frac{mga}{4} + \frac{2\lambda mga}{9} \\ &= \frac{mga}{4} + \frac{2\lambda mga}{9} \end{aligned}$$

↳ By CoE:


$$mga + \frac{\lambda mga}{18} = \frac{mga}{4} + \frac{2\lambda mga}{9}$$

$$1 + \frac{\lambda}{18} = \frac{1}{4} + \frac{2\lambda}{9}$$

$$18 + \lambda = 4.5 + 4\lambda$$

$$3\lambda = 13.5$$

$$\lambda = \frac{9}{2}$$




A uniform lamina is in the form of a triangle ABC , with $AC = 8a$, $BC = 6a$ and angle $ACB = 90^\circ$. The point D on AC is such that $AD = 3a$. The point E on CB is such that $CE = x$ (see diagram). The triangle CDE is removed from the lamina.

- (a) Find, in terms of a and x , the distance of the centre of mass of the remaining object $ADEB$ from AC . [4]

↳ find $\bar{x}$

$$P: \left(\frac{x}{3}, \frac{5a}{3} \right)$$

$$Q: \left(2a, \frac{8a}{3} \right)$$



Taking moments about AC:

$$(\bar{x}) \left(\frac{6a}{2} \right) \left(\frac{8a}{2} - \frac{5ax}{2} \right) = \frac{6a}{2} \left(\frac{8a}{2} \right) (2a) + \frac{5ax}{2} \left(\frac{x}{3} \right)$$

$$\bar{x} \left(\frac{9a}{2} \right) (48a - 5x) = \frac{9a}{2} (96a^2 - \frac{5x^2}{3})$$

$$\bar{x} = \frac{288a^2 - 5x^2}{144a - 15x}$$

The object $ADEB$ is on the point of toppling about the point E when the object is in the vertical plane with its edge EB on a smooth horizontal surface.

(b) Find x in terms of a .

[3]

$$\frac{288a^2 - 5x^2}{144a - 15x} = x$$

$$288a^2 - 5x^2 = 144ax - 15x^2$$

$$10x^2 - 144ax + 288a^2 = 0$$

$$5x^2 - 72xa + 144a^2 = 0$$


$$x = \frac{72a \pm \sqrt{5184a^2 - 4(5)(144a^2)}}{10}$$

$$= 7.2a \pm 4.8a$$

$$= 12a \text{ or } 2.4a$$

$$\downarrow$$

$$(as x \leq 6a)$$



Two identical smooth uniform spheres A and B each have mass m . The two spheres are moving on a smooth horizontal surface when they collide with speeds u and $2u$ respectively. Immediately before the collision, A 's direction of motion makes an angle of 30° with the line of centres, and B 's direction of motion is perpendicular to the line of centres (see diagram). After the collision, A and B are moving in the same direction. The coefficient of restitution between the spheres is e .

(a) Find the value of e .

[5]

→ Only horizontal components are affected by collisions:

$$m \cdot u \cos 30^\circ$$

$$m u \cos 30^\circ = m v_{Ax} + m v_{Bx}$$

$$\frac{\sqrt{3}u}{2} = v_{Ax} + v_{Bx} \quad (1)$$

$$\frac{v_{Bx} - v_{Ax}}{\frac{\sqrt{3}u}{2}} = e$$


$$v_{Bx} - v_{Ax} = \frac{\sqrt{3}eu}{2} \quad (2)$$

Adding (1) & (2):

$$2v_{Bx} = \frac{\sqrt{3}u}{2}(e+1)$$

$$v_{Bx} = \frac{\sqrt{3}u}{4}(e+1)$$

$$v_{Ax} = \frac{\sqrt{3}u}{4}(1-e)$$



$$\frac{v_{Ay}}{v_{By}} = \frac{v_{Ax}}{v_{Bx}}$$

$$\frac{\left(\frac{\sqrt{3}u}{4}\right)(1-e)}{u \sin 30^\circ} = \frac{\left(\frac{\sqrt{3}u}{4}\right)(1+e)}{2u}$$

$$\frac{\sqrt{3}}{2}(1-e) = \frac{\sqrt{3}}{8}(1+e)$$

$$4-4e = e+1$$

$$3 = 5e$$

$$e = \frac{3}{5}$$

(b) Find the loss in the total kinetic energy of the spheres as a result of the collision.

[3]

Initially:

$$KE = \frac{1}{2}mv^2 + \frac{1}{2}m(2v)^2 = \frac{5}{2}mv^2$$

Finally:

$$V_A = \sqrt{\left(\frac{v}{2}\right)^2 + \left(\left(\frac{\sqrt{3}v}{4}\right)\left(\frac{2}{5}\right)\right)^2} \quad \left(\frac{3}{16}\right)\left(\frac{4}{25}\right)$$

$$= \sqrt{\frac{v^2}{4} + \frac{3v^2}{100}} = \sqrt{\frac{28v^2}{100}} = \sqrt{\frac{7v^2}{25}}$$

$$V_B = \sqrt{(2v)^2 + \left(\left(\frac{\sqrt{3}v}{4}\right)\left(\frac{8}{5}\right)\right)^2} \quad \left(\frac{3}{16}\right)\left(\frac{4}{25}\right)$$

$$= \sqrt{4v^2 + \frac{12v^2}{25}} = \frac{48}{25}$$

$$= \sqrt{\frac{112v^2}{25}}$$

$$\text{Final KE} = \frac{1}{2}m\left(\frac{112v^2}{25}\right) + \frac{1}{2}m\left(\frac{7v^2}{25}\right)$$

$$= \frac{119}{50}mv^2$$


$$\text{Loss} = \frac{125}{50}mv^2 - \frac{119}{50}mv^2$$

$$= \frac{3}{25}mv^2$$

- 5 One end of a light elastic string, of natural length $12a$ and modulus of elasticity kmg , is attached to a fixed point O . The other end of the string is attached to a particle of mass m . The particle moves with constant speed $\frac{3}{2}\sqrt{3ag}$ in a horizontal circle with centre at a distance $12a$ below O . The string is inclined at an angle θ to the downward vertical through O .

(a) Find, in terms of a , the extension of the string.

[5]



$$T \cos \theta = mg$$

$$T = \frac{mg}{\left(\frac{12a}{12a+x}\right)} = \frac{mg(12a+x)}{12a}$$

$$T \sin \theta = \frac{mv^2}{r}$$

$$\left(\frac{mg(12a+x)}{12a}\right) \left(\frac{\sqrt{x^2+24ax}}{12a+x}\right) = \frac{m\left(\frac{9}{4}\right)(3ag)}{\sqrt{x^2+24ax}}$$

$$\frac{mg(x^2+24ax)}{12a} = \frac{27}{4} mga$$

$$x^2 + 24ax = 81a^2$$

$$x^2 + 24ax - 81a^2$$

$$(x+27a)(x-3a) = 0$$

$$\hookrightarrow x = 3a \text{ or } -27a$$

$$\hookrightarrow \text{not possible}$$

$$\boxed{x = 3a}$$

(b) Find the value of k .

[3]

$$T = \frac{\lambda x}{l}$$

$$\frac{mg(12a+x)}{12a} = \frac{kmg(3a)}{12a}$$

$$\frac{15mg}{12} = \frac{3kmg}{12}$$

$$k=5$$

- 6 A particle of mass m kg falls vertically under gravity, from rest. At time t s, P has fallen x m and has velocity v ms⁻¹. The only forces acting on P are its weight and a resistance of magnitude $kmgv$ N, where k is a constant.

(a) Find an expression for v in terms of t , g and k .

[5]

$$a_{\text{down}} = g - kgv$$

$$\frac{dv}{dt} = g - kgv$$

$$\frac{dv}{1 - kv} = g dt$$

$$\int \frac{1}{1 - kv} dv = \int g dt$$

$$\frac{\ln|1 - kv|}{-k} = gt + c$$

$\Rightarrow$ starts @ rest, so
when $t=0$, $v=0$

$$\frac{\ln 1}{-k} = c = 0$$

$$\frac{\ln|1 - kv|}{-k} = gt$$

$$\ln|1 - kv| = -gtk$$

$$\ln|1 - kv| = -gtk$$

$$1 - kv = e^{-gtk}$$

$$\frac{1 - e^{-gtk}}{k} = v$$

- (b) Given that $k = 0.05$, find, in metres, how far P has fallen when its speed is 12 m s^{-1} . [5]

$$v = 20 - 20e^{-0.05gt}$$

when $v = 12$

$$8 = 20e^{-0.05gt}$$

$$e^{0.05gt} = \frac{5}{2}$$

$$t = \frac{\ln 2.5}{0.05g} = 1.833 \text{ s}$$

$$x = \int_0^{1.833} (20 - 20e^{-0.05gt}) dt$$

$$= \left[20t + \frac{20e^{-0.05gt}}{0.05g} \right]_0^{1.833}$$

$$= 20(1.833) + \frac{20(\frac{2}{5})}{0.05g} - \frac{20e^0}{0.05g}$$


$$= 36.65 + \frac{8(20)}{g} - 40$$

$$\boxed{= 12.7 \text{ m}}$$

- 7 The points O and P are on a horizontal plane, a distance 8 m apart. A ball is thrown from O with speed $u \text{ m s}^{-1}$ at an angle θ above the horizontal, where $\tan \theta = \frac{4}{3}$. At the same instant, a model aircraft is launched with speed 5 m s^{-1} parallel to the horizontal plane from a point 4 m vertically above P . The model aircraft moves in the same vertical plane as the ball and in the same horizontal direction as the ball. The model aircraft moves horizontally with a constant speed of 5 m s^{-1} . After T s, the ball and the model aircraft collide.

(a) Find the value of T .

[6]



@ Time T :

Ball: Horizontal distance = $5T$

Plane: Horizontal distance = $5T \cos \theta = \frac{30T}{5}$

Ball: Height = 4 m

Plane: $y = x \tan \theta - \frac{g x^2}{2u^2} \sec^2 \theta$

$$= \frac{4}{3}x - \frac{125x^2}{9u^2}$$

$$\hookrightarrow = \frac{40T}{5} - 5T^2$$

$$\frac{40T}{5} - 5T^2 = 4$$

$$5T + 8 = \frac{30T}{5}$$

$$\frac{5T^2 + 4}{4} = \frac{5T + 8}{3}$$

$$\sin \theta = \frac{4}{5}$$

$$\cos \theta = \frac{3}{5}$$

$$\tan \theta = \frac{4}{3}$$

$$15T^2 + 12 = 20T + 32$$

$$15T^2 - 20T - 20 = 0$$

$$3T^2 - 4T - 4 = 0$$

$$(3T + 2)(T - 2)$$

$$\hookrightarrow \boxed{T = 2}$$

(b) Find the direction in which the ball is moving immediately before the collision.

[3]

$$5\tau + 8 = \frac{30\tau}{5}$$

$$1.8 = \frac{6v}{5}$$

$$v = 15 \text{ ms}^{-1}$$

for θ from the horizontal:

$$\tan \theta = \frac{v \sin \theta - gt}{v \cos \theta}$$

$$= \frac{15\left(\frac{4}{5}\right) - 20}{15\left(\frac{3}{5}\right)} = \frac{-8}{9}$$

$$\theta = 41.6^\circ \text{ below horizontal}$$

[illegible]

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.