



Cambridge International AS & A Level

CANDIDATE
NAME
CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

FURTHER MATHEMATICS

9231/43

Paper 4 Further Probability & Statistics

May/June 2025

1 hour 30 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [].

This document has **12** pages.



- 1 A person's eye colour may be categorised as "brown", "blue" or "other". A scientist claims that these eye colours are uniformly distributed and hence are equally likely to occur in the population. A survey of 120 people from this population found that 38 people had brown eyes, 52 people had blue eyes and 30 people had eyes which were neither brown nor blue.

Use the data to carry out a goodness of fit test at the 5% significance level to test the scientist's claim.

[6]

H_0 : uniform distribution is a good fit

H_1 : not a good fit

table of expected frequencies

Brown	Blue	other	total
40	40	40	120

$$\text{test statistic} = \sum \frac{(\text{Exp} - \text{Obs})^2}{\text{Exp}}$$

$$= \frac{(38-40)^2}{40} + \frac{(52-40)^2}{40} + \frac{(30-40)^2}{40}$$

$$= 0.1 + 3.6 + 2.5 = 6.2$$

degrees of freedom $\rightarrow n - \overset{\text{estimated parameters}}{v} - 1$
 $\rightarrow 3 - 0 - 1 = 2$

tabular value

$$\rightarrow \chi^2_2(0.95) = 5.991$$

as $6.2 > 5.991 \rightarrow \text{reject } H_0 \rightarrow \text{in critical region}$

sufficient evidence to suggest uniform distribution is not a good fit



- 2 A farmer is investigating whether using a new fertiliser will increase the yield of tomato plants. The farmer selects 40 tomato plants at random and gives them the new fertiliser. The crop mass, x kg, of each of these 40 plants is recorded. The farmer selects a further 60 tomato plants at random and gives them a standard fertiliser. The crop mass, y kg, of each of these 60 plants is recorded. The results are summarised as follows.

$$\Sigma x = 168 \quad \Sigma x^2 = 720 \quad \Sigma y = 228 \quad \Sigma y^2 = 900$$

Find a 90% confidence interval for the difference in mean crop mass associated with each type of fertiliser. [7]

C.I. $\rightarrow$ diff in means $\rightarrow (\bar{x} - \bar{y}) \pm z_{\frac{\alpha}{2}} \sqrt{\frac{s^2_x}{n_x} + \frac{s^2_y}{n_y}}$

as $n > 30$

$$\bar{x} = \frac{168}{40} = 4.2 \quad \bar{y} = \frac{228}{60} = 3.8$$

$$s^2_x = \frac{1}{39} \left(720 - \frac{(168)^2}{40} \right) = \frac{24}{65}$$

$$s^2_y = \frac{1}{59} \left(900 - \frac{228^2}{60} \right) = \frac{168}{295} \quad \text{as } 100(\alpha - 1)\% = 90\%$$

$$\sqrt{\frac{s^2_x}{n_x} + \frac{s^2_y}{n_y}} = \sqrt{0.0187}$$

$$C.I. = (4.2 - 3.8) \pm z_{0.05} (\sqrt{0.0187})$$

$$(4.2 - 3.8) \pm 1.645 (\sqrt{0.0187})$$

$$0.4 \pm 0.225 = [0.175, 0.625]$$

- 3 A continuous random variable X has probability density function f given by

$$f(x) = \begin{cases} kx & 0 \leq x < 1, \\ k(8-x) & 1 \leq x \leq 8, \\ 0 & \text{otherwise,} \end{cases}$$

where k is a constant.

- (a) Show that $k = \frac{1}{25}$.

$$\left(\int_0^1 kx \, dx + k \int_1^8 (8-x) \, dx \right) = 1 \quad \text{area below curve [2]}$$

$$\left[\frac{kx^2}{2} \right]_0^1 + k \left[8x - \frac{x^2}{2} \right]_1^8 = 1$$

$$\frac{k}{2} + \frac{49k}{2} = 1 \quad \rightarrow \quad \boxed{k = \frac{1}{25}}$$

- (b) Find the median value of X .

[3]

$$\int_0^1 kx \, dx + \int_1^m k(8-x) \, dx = 0.5$$

$$\left[\frac{1}{50} x^2 \right]_0^1 + \left[\frac{8}{25} x - \frac{1}{50} x^2 \right]_1^m = 0.5$$

calc gives median

$$\frac{1}{50} + \left(\left(\frac{8}{25} m - \frac{1}{50} m^2 \right) - \left(\frac{8}{25} - \frac{1}{50} \right) \right) = 0.5$$

$$m^2 - 16m + 39 = 0$$

$$(m-13)(m-3) = 0 \quad \rightarrow \quad \boxed{m=3} \quad \text{in range}$$

The random variable Y is defined by $Y = \sqrt[3]{X}$.

(c) Find the probability density function of Y .

$\frac{1}{50}(1) = \frac{8}{25}(1) - \frac{1}{50}(1) = k$ → to maintain continuity
[5]

$k = \frac{-7}{25}$

$$F(x) = \begin{cases} 0 & x < 0 \\ \frac{1}{50}x^2 & 0 \leq x \leq 1 \\ \frac{8}{25}x - \frac{1}{50}x^2 - \frac{7}{25} & 1 \leq x \leq 8 \\ 1 & x > 8 \end{cases}$$

$y = x^{1/3} \quad y^3 = x$

$G(y)$

$$= \begin{cases} 0 & y < 0 \\ \frac{1}{50}y^6 & 0 \leq y \leq 1 \\ \frac{8}{25}y^3 - \frac{1}{50}y^6 - \frac{7}{25} & 1 \leq y \leq 2 \\ 1 & y > 2 \end{cases}$$

limits also follow $y^3 = x$

differentiate $G(y)$ for pdf in $y \rightarrow g(y)$

$$g(y) = \begin{cases} \frac{3}{25}y^5 & 0 \leq y \leq 1 \\ \frac{24}{25}y^2 - \frac{3}{25}y^5 & 1 \leq y \leq 2 \\ 0 & \text{otherwise} \end{cases}$$

- 4 A researcher claims that older people take longer to react to a sudden loud noise than younger people. To investigate this, the researcher randomly selects 6 people over 50 years old and 8 people under 25 years old and records their reaction times, in milliseconds, to a sudden loud noise. The reaction times are as follows.

Over 50 198 212 217 229 235 242

Under 25 178 181 183 192 203 209 223 231

Carry out a Wilcoxon rank-sum test at the 5% significance level to test the researcher's claim. [8]

H_0 : pop medians for <25 and >50 are equal

H_1 : pop medians for <25 is slower than that of >50

Rankings

	Rank	>50	<25
178	1		1
181	2		2
183	3		3
192	4		4
198	5	5	
203	6		6
209	7		7
217	9	9	
223	10		10
212	8	8	
229	11	11	
231	12		12
235	13	13	
242	14	14	
		60	45
		$\rightarrow R_6$	$\rightarrow R_8$





7

as size of > 50 is less than < 25

$$W = \min(60, 6(8.16 - 11) - 60)$$

test
stat

$$= \min(60, 30)$$

$$\boxed{\bar{w} = 30}$$

tabular value $\rightarrow 5\%$, one-tailed, $m = 6$, $n = 8$

$\rightarrow \boxed{31} \rightarrow$ critical value

as $30 < 31 \rightarrow$ reject $H_0 \rightarrow$ shows symmetry

$\therefore$ sufficient evidence to suggest that
older people take longer to read



- 5 A doctor is investigating the concentration of blood glucose in patients at risk of developing type 2 diabetes, where blood glucose is measured in appropriate units. The doctor claims that a particular intervention reduces the concentration by more than k units on average. A group of 8 at risk patients is selected at random and each patient follows the intervention for six months. The blood glucose concentrations before and after the intervention are given in the following table.

Patient	A	B	C	D	E	F	G	H
Before	183	165	172	165	143	176	161	153
After	164	148	164	149	134	153	155	148

- (a) Use a t -test at the 5% significance level to find the range of values of k for which the result of the test is to reject the null hypothesis. [7]

+ - test for paired differences

$$\rightarrow \frac{\bar{d} - k}{\frac{S_d}{\sqrt{n}}}$$

sum of differences

$$\sum d = 19 + 17 + 8 + 16 + 9 + 23 + 6 + 5 = 103$$

$$\sum d^2 = 1641$$

sum of (diff)²

$$\bar{d} = \frac{\sum d}{n} = \frac{103}{8} = 12.875$$

$$S_d^2 = \frac{1}{n-1} \left(\sum d^2 - \frac{(\sum d)^2}{n} \right) = 44.982$$

$$H_0: \mu_b - \mu_a = k \quad H_1: \mu_b - \mu_a > k$$

$$T = \frac{12.875 - k}{\sqrt{\frac{44.982}{8}}}$$

$$\geq t_{0.05, 7} \quad \text{degrees of freedom}$$

one-tailed





$$12.875 - h > 1.895$$

$$\sqrt{44.982}$$

$$8$$

$$12.875 - h \geq 4.49349$$

$$h < 8.38151$$

$$h \leq 8.38$$

→

- (b) State an assumption necessary for the test in part (a) to be valid.

[1]

population of differences are normally distributed → required for a t-test



- 6 A bag contains 7 red balls and 3 blue balls. Kieran selects 2 balls at random, without replacement. The number of red balls selected by Kieran is denoted by X , and the number of different colours present in Kieran's selection is denoted by Y .

(a) Find the probability generating functions, $G_X(t)$ of X and $G_Y(t)$ of Y . [4]

$$P(X=0) = \frac{3}{10} \times \frac{2}{9} = \frac{6}{90} = \frac{1}{15} \quad \text{no. of arrangements}$$

$$P(X=1) = \left(\frac{7}{10} \times \frac{3}{9} \right) \times 2 = \frac{7}{15} \quad \text{due to distinct colours}$$

$$P(X=2) = \left(\frac{7}{10} \times \frac{6}{9} \right) = \frac{7}{15}$$

$$G_X(t) = \frac{1}{15} + \frac{7}{15}t + \frac{7}{15}t^2 \rightarrow \sum t^x P(X=x)$$

$$G_X(1) = 1 \rightarrow \text{so true} \rightarrow \text{double check}$$

$$P(1 \text{ colour}) = P(BB) + P(RR) = \left(\frac{3}{10} \times \frac{2}{9} \right) + \left(\frac{7}{10} \times \frac{6}{9} \right) = \frac{8}{15}$$

$$\begin{aligned} P(2 \text{ colour}) &= P(BR) + P(RB) \\ &= \left(\frac{3}{10} \times \frac{7}{9} \right) + \left(\frac{7}{10} \times \frac{3}{9} \right) \\ &= \frac{7}{15} \end{aligned}$$

$$G_Y(t) = \frac{8}{15}t + \frac{7}{15}t^2$$

$$G_Y(1) = 1 \rightarrow \text{correct} \rightarrow \text{to verify}$$

The random variable Z is the sum of the number of red balls and the number of different colours present in Kieran's selection. Kieran claims that the probability generating function of Z is equal to $G_X(t) \times G_Y(t)$.

(b) Explain why Kieran is incorrect. [1]

X and Y are not independent events but rather directly related



- (c) Find the probability generating function of Z , expressing your answer as a polynomial in t . [4]

$Z = (\text{red balls} + \text{diff colours})$

$$+ P(R + 2 \text{ colour}) + P(RR + 1 \text{ colour}) + P(OR + 1 \text{ colour})$$

$$\rightarrow P(3) = \frac{7}{15} + \frac{7}{15} \quad P(1) = \frac{1}{15}$$

$$P(2) = 0$$

$$G_Z(t) = \frac{1}{15} + 0 + \frac{14}{15}t + 3$$

- (d) Use the probability generating function of Z to find $E(Z)$. [2]

$$G'_Z(t) \text{ where } t=1 \rightarrow E(Z)$$

$$G'_Z(t) = \frac{1}{15} + \frac{14}{5}t^2 \quad E(Z) = \frac{1}{15} + \frac{14}{5} = \frac{43}{15}$$

This image shows a full page of a handwriting practice worksheet. It consists of multiple rows of horizontal dashed lines spaced evenly apart, providing a guide for letter height and placement. The background is plain white, and there are no other markings or text on the page.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.