


- 1 One end of a light elastic string, of natural length a and modulus of elasticity $3mg$, is attached to a fixed point O . The other end of the string is attached to a particle P of mass m . The string hangs with P vertically below O . The particle P is pulled vertically downwards so that the extension of the string is $2a$. The particle P is then released from rest.

- (a) Find the speed of P when it is at a distance $\frac{3}{4}a$ below O . [3]

Initially:



$$PE = 0$$


$$KE = 0$$

$$EPE = \frac{\lambda x^2}{2l} = \frac{3mg(2a)^2}{2a} = 6mga \text{ J}$$

$h=0$

Finally:

→ By CoE, energy of system = $6mga$.



$$\left. \begin{array}{l} EPE = 0 \\ PE = \frac{3}{4}mga \end{array} \right\} \begin{array}{l} KE = \left(6 - \frac{3}{4}\right)mga \\ = \frac{15}{4}mga = \frac{1}{2}m\left(\sqrt{\frac{15}{2}}ga\right)^2 \end{array} \rightarrow \therefore v = \sqrt{7.5ga}$$


- (b) Find the initial acceleration of P when it is released from rest. [2]

$$T = \frac{\lambda x}{l} = \frac{3mg(2a)}{a} = 6mg$$

$$W = mg$$

net force = $5mg$ upward.

$$a = 5g \text{ ms}^{-2}$$



A particle P of mass m is moving with speed u on a fixed smooth horizontal surface. It collides at an angle α with a fixed smooth vertical barrier. After the collision, P moves at an angle θ with the barrier, where $\tan \theta = \frac{1}{2}$ (see diagram). The coefficient of restitution between P and the barrier is e . The particle P loses 20% of its kinetic energy as a result of the collision.

Find the value of e .

[5]

Only horizontal components are affected by this collision



$$\text{Final KE} = 0.8 \left(\frac{1}{2} m u^2 \right)$$

$$= \frac{1}{2} m \left(\frac{2\sqrt{5}}{5} u \right)^2 \Rightarrow \boxed{\text{final } v = \frac{2\sqrt{5}}{5} u}$$

$$v_y = v_y = \frac{2\sqrt{5}}{5} u \cos \theta = \frac{4}{5} u$$

$$\text{we know that } \sqrt{v_x^2 + v_y^2} = v$$

$$\hookrightarrow v^2 = v_x^2 + \frac{16}{25} v^2$$

$$\frac{9}{25} v^2 = v_x^2$$

$$v_x = \frac{3}{5} v$$

$$\sqrt{v_x^2 + v_y^2} = v$$

$$v_x^2 + \frac{16}{25} v^2 = \left(\frac{2\sqrt{5}}{5} v \right)^2$$

$$v_x^2 = \frac{4}{25} v^2$$

$$v_x = \frac{2}{5} v$$

$$v_x = e v_x$$


$$e = \frac{\frac{2}{5} v}{\frac{3}{5} v} = \boxed{\frac{2}{3}}$$

- 3 A particle P of mass m is attached to one end of a light inextensible string of length a . The other end of the string is attached to a fixed point O . The particle P is held at the point A , where OA makes an angle θ with the downward vertical through O , and with the string taut. The particle P is projected perpendicular to OA in an upwards direction with speed u . It then starts to move along a circular path in a vertical plane. The string goes slack when P is at B , where angle AOB is 90° and the speed of P is $\sqrt{\frac{4}{5}ag}$.

- (a) Find the value of $\sin \theta$.

180 - (0+90)

[2]




Considering forces @ B:

$$T = \frac{mv^2}{r} + mg \cos(\theta + 90)$$

$$\frac{mv^2}{r} = mg \sin \theta$$

$$\frac{4}{5}ag = g \sin \theta$$

$$\sin \theta = \frac{4}{5}$$



- (b) Find, in terms of m and g , the tension in the string when P is at A .

[5]

Taking speed at bottom of circle = v

$$T_A = \frac{mv^2}{a} - 2mg + 3mg \cos \theta \Rightarrow \cos \theta = \frac{3}{5}$$

$$= \frac{mv^2}{a} - \frac{mg}{5}$$


$$0 = T_B = \frac{mv^2}{a} - 2mg + 3mg \cos(\theta + 90)$$

$$\frac{mv^2}{a} = 2mg + \frac{12}{5}mg$$

$$v^2 = 4.4ga$$

$$T_A = \frac{m(4.4ga)}{a} - \frac{mg}{5}$$


$$= 4.2mg$$



An object is formed from a solid hemisphere, of radius $2a$, and a solid cylinder, of radius a and height d . The hemisphere and the cylinder are made of the same material. The cylinder is attached to the plane face of the hemisphere. The line OC forms a diameter of the base of the cylinder, where C is the centre of the plane face of the hemisphere and O is common to both circumferences (see diagram). Relative to axes through O , parallel and perpendicular to OC as shown, the centre of mass of the object is $(\bar{x}, \bar{y})$.

- (a) Show that $\bar{x} = \frac{32a^2 + 3ad}{16a + 3d}$ and find an expression, in terms of a and d , for $\bar{y}$. [5]

By symmetry, $(\bar{x}, \bar{y})$ lies on the axis, so we need only consider a 2D image:



$$A: (a, \frac{d}{2})$$

$$B: (2a, -\frac{3a}{4})$$

About x :

$$(\bar{x}) \left(\frac{\pi}{3} \pi (2a)^3 \rho + \pi a^2 d \rho \right) = \left(\frac{\pi}{3} \pi (2a)^3 \rho \right) (2a) + (\pi a^2 d \rho) (a)$$

$$(\bar{x}) (\pi a^2 \rho) \left(\frac{16}{3} a + d \right) = (\pi a^2 \rho) \left(\frac{32}{3} a^2 \right) + (\pi a^2 \rho) (ad)$$

$$\bar{x} = \frac{\frac{32}{3} a^2 + ad}{\frac{16}{3} a + d} = \frac{32a^2 + 3ad}{16a + 3d}$$

About y :


$$\bar{y} (\pi a^2 \rho) \left(\frac{16}{3} a + d \right) = (\pi a^2 \rho) (-4a^2) + (\pi a^2 \rho) \left(\frac{d^2}{2} \right)$$

$$\bar{y} = \frac{\frac{d^2}{2} - 4a^2}{\frac{16}{3} a + d} = \frac{3d^2 - 24a^2}{32a + 6d}$$

The object is placed on a rough plane which is inclined to the horizontal at an angle θ where $\sin \theta = \frac{1}{6}$. The object is in equilibrium with CO horizontal, where CO lies in a vertical plane through a line of greatest slope.

(b) Find d in terms of a .

[3]



$$\text{COR: } \left(\frac{32a^2 + 3ad}{16a + 3d}, \frac{3d^2 - 24a^2}{32a + 6d} \right)$$

$$\rightarrow \text{We need to find } z, \text{ as } z = \frac{32a^2 + 3ad}{16a + 3d}$$

$$\sin \theta = \frac{2a - z}{2a}$$

$$z = 2a - 2a \sin \theta$$

$$= \frac{5a}{3}$$

$$\hookrightarrow \frac{32a^2 + 3ad}{16a + 3d} = \frac{5a}{3}$$

$$96a^2 + 9ad = 80a^2 + 15ad$$

$$16a + 9d = 80a + 15d \quad \text{as } a \neq 0$$

$$16a = 6d$$

$$d = \frac{8a}{3}$$

- 5 A light elastic string of natural length a and modulus of elasticity λmg has one end attached to a fixed point O on a smooth horizontal surface. When a particle of mass m is attached to the free end of the string, it moves with speed v in a horizontal circle with centre O and radius x . When, instead, a particle of mass $2m$ is attached to the free end of the string, this particle moves with speed $\frac{1}{2}v$ in a horizontal circle with centre O and radius $\frac{3}{4}x$.

(a) Find x in terms of a .

[5]

In both cases, Tension = Centripetal force

$$\textcircled{1} \quad \frac{(\lambda mg)(x-a)}{a} = \frac{mv^2}{x}$$

$$\lambda gx^2 - \lambda gax = av^2$$

$$\textcircled{2} \quad \frac{(\lambda mg)(\frac{3}{4}x-a)}{a} = \frac{2m(\frac{1}{2}v)^2}{\frac{3}{4}x}$$

$$\frac{9}{16} \lambda gx^2 - \frac{3}{4} \lambda gxa = \frac{1}{2} av^2$$

Equating:

$$\frac{9}{8} \lambda gx^2 = \frac{3}{2} \lambda gxa = \lambda gx^2 - \lambda gxa$$

$$\frac{9}{8}x - \frac{3}{2}a = x - a$$

$$\frac{1}{8}x = \frac{1}{2}a$$

$$\boxed{x = 4a}$$

(b) Given that $v = \sqrt{12ag}$, find the value of λ .

[2]

$$\lambda g x^2 - \lambda g x a = a v^2$$

$$\hookrightarrow \lambda g (4a)^2 - \lambda g (4a)(a) = a(12ag)$$

$$16\lambda g a^2 - 4\lambda g a^2 = 12g a^2$$

$$12\lambda = 12$$

$$\boxed{\lambda = 1}$$

- 6 A particle P moving in a straight line has displacement x m from a fixed point O on the line and velocity v m s⁻¹ at time t s. The acceleration of P , in m s⁻², is given by $6v\sqrt{v+9}$. When $t = 0$, $x = 2$ and $v = 72$.

(a) Find an expression for v in terms of x .

[4]

$$a = 6v\sqrt{v+9}$$

We know:

$$\frac{dx}{dt} \left(\frac{dv}{dx} \right) = \frac{dv}{dt}$$

$$\frac{dv}{dx} = \frac{a}{v}$$

∴

$$\frac{dv}{dx} = 6\sqrt{v+9}$$

$$\hookrightarrow \frac{1}{\sqrt{v+9}} dv = 6 dx$$

$$\int (v+9)^{-1/2} dv = \int 6 dx$$

$$2(v+9)^{1/2} = 6x + c$$

$$\sqrt{v+9} = 3x + c$$

∴ when $x=2$, $v=72$, so:

$$9 = 6 + c$$

$$\hookrightarrow c = 3$$

$$v+9 = (3x+3)^2$$

$$v = 9x^2 + 18x$$

(b) Find an expression for x in terms of t .

[5]

$$\frac{dx}{dt} = 9x^2 + 18x$$

$$\frac{1}{x^2 + 2x} dx = 9 dt$$

$$\int \frac{1}{x^2 + 2x} dx = \int 9 dt$$

$$\hookrightarrow \frac{1}{x^2 + 2x} = \frac{A}{x+2} + \frac{B}{x}$$

$$\hookrightarrow Ax + Bx + 2B = 1$$

$$\hookrightarrow B = \frac{1}{2}$$

$$A = -\frac{1}{2}$$

$$\Rightarrow \int \frac{1}{2x} - \frac{1}{2x+4} dx = 9t + C$$

$$\int \frac{1}{x} - \frac{1}{x+2} dx = 18t + C$$

$$\ln|x| - \ln|x+2| = 18t + C$$

$$\hookrightarrow \text{when } t=0, x=2$$

$$\hookrightarrow \ln \frac{2}{4} = C$$

$$C = \ln \frac{1}{2}$$

$$\ln \left| \frac{x}{x+2} \right| = 18t + \ln \frac{1}{2}$$

$$\ln \left| \frac{2x}{x+2} \right| = 18t$$

$$\frac{2x}{x+2} = e^{18t}$$

$$2x = e^{18t} x + 2e^{18t}$$

$$x = \frac{2e^{18t}}{2 - e^{18t}}$$

- 7 At time t s, a particle P is projected with speed 40 ms^{-1} at an angle θ above the horizontal from a point O on a horizontal plane and moves freely under gravity. The greatest height achieved by P during its flight is H m and the corresponding time is T s.

(a) Obtain expressions for H and T in terms of θ .

[2]

$$v_y = u \sin \theta - gt \Rightarrow \text{@ max height, } v_y = 0, \text{ so } t = \frac{u \sin \theta}{g} = \boxed{4 \sin \theta}$$

$$\begin{aligned} H &= u \sin \theta T + \frac{1}{2}(-g)(T)^2 = (40 \sin \theta)(4 \sin \theta) - \frac{1}{2}(10)(4 \sin \theta)^2 \\ &= 160 \sin^2 \theta - 80 \sin^2 \theta \\ &= \boxed{80 \sin^2 \theta} \end{aligned}$$

During the time between $t = T$ and $t = 3$, P descends a distance $\frac{1}{4}H$.

(b) Find the value of θ .

[4]

$$\begin{aligned} \text{@ } t = T, \quad H &= 80 \sin^2 \theta \\ \text{@ } t = 3, \quad H &= 60 \sin^2 \theta \end{aligned} \quad \left. \begin{array}{l} \\ \end{array} \right\} \begin{array}{l} 20 \sin^2 \theta \text{ m descended in} \\ (3 - T) \text{ s} \end{array}$$

$$\begin{aligned} \hookrightarrow s &= 20 \sin^2 \theta \\ t &= 3 - 4 \sin \theta \end{aligned} \quad \left(\begin{array}{l} \text{taking down as +} \end{array} \right)$$

$$\begin{aligned} a &= g \\ u &= 0 \end{aligned}$$

$$20 \sin^2 \theta = \frac{1}{2}(g)(3 - 4 \sin \theta)^2$$

$$4 \sin^2 \theta = 9 + 16 \sin^2 \theta - 24 \sin \theta$$

$$0 = 12 \sin^2 \theta - 24 \sin \theta + 9$$

$$= 4 \sin^2 \theta - 8 \sin \theta + 3$$

$$(2 \sin \theta - 3)(2 \sin \theta - 1)$$

$$\hookrightarrow \sin \theta = \frac{1}{2}$$

$$\hookrightarrow \theta = 30^\circ$$

(c) Find the speed of P when $t = 3$.

[3]

$$\begin{aligned}v_y &= 40 \sin \theta - g(3) \\&= 40 \left(\frac{1}{2}\right) - 30 \\&= -10 \text{ ms}^{-1}\end{aligned}$$

$$v_x = v \cos \theta = 40 \left(\frac{\sqrt{3}}{2}\right) = 20\sqrt{3} \text{ ms}^{-1}$$

$$\text{speed} = \sqrt{100 + 1200} = 10\sqrt{13} \text{ ms}^{-1}$$

[illegible]

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.