

Deep Learning Based Cyclone Intensity Estimation using INSAT-3D IR Imagery

Uma N

Computer Science and Engineering
New Horizon College of
Engineering Bengaluru, India.
umamam@gmail.com

Joshua James

Computer Science and Engineering
New Horizon College of
Engineering Bengaluru, India.
joshua.james699@gmail.com

Jithin Vinod

Computer Science and Engineering
New Horizon College of
Engineering Bengaluru, India.
jitvin10@gmail.com

K M Komal

Computer Science and
Engineering New Horizon
College of Engineering
Bengaluru, India.
lamokomal27@gmail.com

Abstract— Tropical cyclones stand out as one of the most financially burdensome natural disasters globally, given the numerous potential risks they pose. Thus, the development of a meticulous assessment model for gauging the severity of tropical cyclones becomes crucial in averting property damage and preserving lives. Various methods and approaches are currently employed, each varying in effectiveness, to assess the atmospheric velocity of tropical cyclones utilizing satellite information at given moment. This research contributes a diagnostic assessment of tropical cyclone strength rooted in deep learning from infrared satellite data, demonstrating RMSE measuring 13.24 knots. Moreover, the study introduces part of the inaugural visualization portals of its kind in a production system. This portal showcases contextual information and the outcomes, providing end-users with valuable insights.

Keywords—Deep learning, Advanced learning, velocity prediction, transition from study to application, lifecycle of automated learning, interpretation of models

I. INTRODUCTION

Tropical cyclones are to blame for over 100,000 fatalities globally and have the potential to kill up to 1000 people in an incident [1]. When inflation is taken into consideration, the projected economic losses from many hurricanes in recent years exceed \$50 billion [1]. In the past, some storms have caused damage exceeding \$100 billion. Insurance Institute, covered seaside assets in the United States alone is worth beyond trillions and is susceptible to storms that make landfall. Numerous risks are connected to storms, and numerous variables may influence the ultimate fatality count, property damage estimates. Models of hurricane damage calculate danger by multiplying the gustiness velocity by an $\exp[2]$ or pow . Its, crucial for disaster planning and response and be able to precisely determine the strength, which is the clear cut highest sustained surface wind speed. There are a handful of clear observations in the winds among the tc, especially when it's over open water. Thus, satellite data is utilized in advance diagnosis of strength. The NHC's acknowledges a potential 10–20% margin of error in their post-analysis based on observations". This poses a challenge in precisely assessing the strength using satellite data. Similarly, The NHC states that the customary 24-hour

evaluation deviation in intensity for individual storms are between 10 - 20 percent. Therefore, increasing Onset force estimations employing satellite imagery may result in notable advancements in short-term intensity forecasts that would enhance our country's readiness and prompt response to disasters. When direct data from planes are not available, meteorologists generally count on variants in Dvorak approach, A method utilizing satellite imagery established in the 70s [3], for predicting tropical cyclone strength. These methods make use of human visual evaluation of cloud field characteristics including the bending of the outer rain belts and the symmetries of the central core and the infrared satellite-recorded brightness temperatures. Dvorak-based methods have some acknowledged drawbacks, one of which being the interpretation of human storm intensity classification on cloud-centric characteristics. Using this method, two skilled analysts can simultaneously determine differing strength assessments for identical instances of storm. Wind speed estimates for hurricane-forced storms may vary by approximately 12 knots (hereafter referred kts) or roughly 6 ms^{-1} , although when they exhibit agreement within 0.5 (T-Number), reflecting precision of Dvt estimates. Additionally, anecdotal data proposes that significant variations can occasionally be observed while contrasting estimates from several agencies and in situations involving complicated cloud formations.

II. MATERIALS AND METHODS

A. Dvorak Technique

This Technique is a method utilized to forecast the strength of tropical cyclones, like hurricanes and typhoons, relying on satellite-derived predictions from numerous images over an extended duration [4]. It was formulated by Dr. Vernon Dvorak in the early 1970s and is widely employed by meteorologists to assess strength of tc. Based on its imaging, the Technique is a method used to appraise the severity. Meteorologists use it frequently to gauge strong cyclones. Providing a brief explanation of the Dvorak Technique's operation: 1) Imagery from satellites: Meteorologists examine cloud patterns linked to the cyclones using probe photography. The arrangement

and form of cloud patterns surrounding the storm core is the primary characteristic that is analyzed.

2) Pattern Classification: Using a scale, cloud patterns are categorized into various tropical numbers (T-No). Each of these numbers represents a different degree of storm strength and is connected to particular cloud formation Intensity.

3) Estimation: Meteorologists use the assigned T-Number to determine strength of tropical cyclone. The scale typically ranges from T1.0 to T8.0, where greater numbers correspond to more severe storms. A T3.0 might correspond to a tropical storm, while a T6.0 could indicate a powerful hurricane.

4) Adjustments: to the estimated intensity based on additional factors, such as magnitude of storm, its structure, and any environmental conditions that might affect its development.

5) Final Intensity Estimate: The final intensity estimate is then used in official forecasts and reports to communicate the current strength of the tropical cyclone. Dvorak Technique is a valuable tool for estimating storm intensity, it has limitations. It relies on visual interpretation of satellite imagery and may not be as accurate in certain situations, such as when a storm is rapidly intensifying or undergoing structural changes.

Advanced technologies and mathematical models are also used in conjunction for more accurate and timely predictions.




Fig 1. sample of Tropical Cyclone Patterns

B. Advanced Dvorak Technique

In the development of the human operated Dvorak method, The ADT was introduced which signifies a technological progression in meteorology, particularly in evaluating the evaluation of the strength of tropical cyclones by examining satellite data images. It was enhanced by Olander and Velden [5], the change was an automation approach where data images were directly feed into an algorithm which is automated, to determine T-number.

C. DAVT

The deviation-angle variance method was explained and employed by Ritchie in the North Pacific and by Pineros in North Atlantic [6]. Through utilizing (IR) satellite, the evenness of tropical cyclone is quantified. The luminosity of the photosis statistically analysed using a directed gradient method. The gradient vector's degree of alignment may be measured as its departure through the ideal spoke axis. Quantification of cyclone's is rooted in variability of its deviation angle. The two primary drawbacks are that it requires photos with appropriately labelled (tc) centres and it employs various modelling and fitting coefficients for (tc) in various locations, which limits its wider, worldwide use.




Fig.2. Convolutional Neural Network Architecture

D. Tropical Cyclone Strength Assessment Derived from Passive Microwave Imagery

Since microwaves may pass through clouds and expose precipitation within the storm, passive microwave satellite observations provide more comprehensive details about the interior structural formation of the cyclone [7]. Infrared photography proved more successful in predicting and analyzing tropical cyclone strength than passive microwave imaging in the beginning. However, by including microwave data in the ADT, has decreased estimates errors. Only when a storm is intensifying and the cyclone does not exhibit a visible eye, Microwave data is included in the ADT. When contrasted with infrared geostationary satellite observations, the main drawbacks of employing unassertive microwave data for operational assessment of cyclone strength is their restricted frequency over time of observations.

E. CNN

Deep learning, which combines Algorithms involving the essence of acquiring characteristics in machine learning, computer vision, and pattern recognition relies on the underlying principle of multiple strata, each engaged in feature detection. Bridging the conceptual gap between using low-level characteristics (picture elements or just pixels) for constructing high-level abstractions is the core problem in picture classification. The tiered hierarchical learning method of deep learning makes an

effort to seal the semantic divide. Newer cutting-edge picture attained concludes in classification in several areas by recent advances in deep learning techniques, primarily using convolutional neural networks (CNN). Methodology has been employed for a wide spectrum of computer Visual challenges, including object identification, visual prominence detection, picture categorization [8]. Tailored to the objective, every task utilizes a marginally varied network topology. Convolutional, pooling, and fully linked layers make up each of their fundamental parts, which are all quite comparable.

Convolutional layers are mostly used to acquire feature interpretations for the inputs [9]; a feature is any kind of input pattern, cloud streak on a satellite picture. The chief purpose in this level is convolution, where each network iteration involves learning the filter values of the convolution filters, leveraged to convolve the input, result is then subjected applying a non-linear activation function element-wise to produce feature maps the rectified linear unit (ReLU) serves as the network's nonlinearity introduced by activation function.

Typically, the pooling layers are been placed between the convolutional layers. mainly employed to shrink the representation's spatial size, which lowers the scale of network calculations and parameters. With deploying average pooling or the max pooling, the input is resized geographically while working independently on each feed. At higher levels of representation, the network may be able to produce increasingly abstract features by piling on multiple layers. The densely connected layer's job is to carry out the classification and high-level reasoning, usually with help of a classifier, Soft Max. During training, the most used back propagation approach for updating parameters is (SGD). To assist lower variation in updating of parameter's and promote more consistent convergence, each SGD parameter update is resolved with regard to a small batch. The pace of convergence is dictated by the learning rate. As our prior research has shown, tropical storm category assessment using CNN is feasible.

Our findings demonstrate that employing CNNs is superior than current methods. Estimating wind velocity at intervals of a range till 5 kt using data from the Research Laboratory of Navy. By adding fresh GOES images, helps in improving to predict wind velocity at 1 kt intervals, conducting thorough evaluations, and developing and implementing a production system, we expand on earlier work. CNN's are also been inducted in other research projects for hurricane intensity estimation. CNN track data, 3D reanalysis data, and other parameters including location data and maximum sustained windspeed. They define the tracking issue as calculating the displacement of the eye between its present location and its future location. Hurricanes and other climate-related occurrences are also effectively tracked and forecasted using other CNN variants, such as LSTM and U-Net [10]. The model described is exclusively a diagnostic tool for evaluating the strength of tropical cyclones; It does not consider Path and predictions of storm.

III. METHODOLOGY

Hurricanes and other climate-related occurrences are also effectively tracked and forecasted using other CNN variants, such as LSTM and U-Net. The model described is exclusively a diagnostic tool for evaluating the strength of tropical cyclones; It does not consider Path and predictions of storm.

A. Machine Learning Lifecycle

The issue formulation, data gathering and evaluating model building and assessment, and rollout to a production environment are the four primary stages of the ml lifecycle. Plenty of ml projects come to an end when it can be demonstrated that the model exhibits enhanced accuracy when compared. Therefore, the rigorous assessment stage in the machine learning lifecycle which helps understand the procedure how, the model renders decisions, and the model's Implementation in a live environment, where it is possible to be tested using fresh real-time data, are the stages of the lifecycle that are typically skipped. Learned to construct own system to produce, using the entire ml's lifecycle as a roadmap.

B. Problem Definition

The challenge is to use satellite imagery to impartially assess the velocity of wind. The primary causes of the concluding fatality figure or amount of damage amidst a tc are frequently linked to the hurricane's wind speed. Therefore, precise cyclone strength (wind speed) estimation is crucial for disaster response and preparation. Because of the individual interpretation of Dvorak approach and factual criteria used to regulate the evolution of cyclone strength over time, there are native setbacks with the present methods used to measure cyclone winds. Furthermore, as present methods operate at hourly temporal frequencies or less, Intensity estimates at a peak frequency made in prediction advisories would provide projection and predictive experts an additional piece of information to consider when determining the cyclone's maximum wind speed.

C. Analysis and Data Collection

The INSAT-3D satellite provides infrared photos of the Indian subcontinent, which are fed into the prototype. Algorithm is trained using a sizable collection of historical photos of cyclones together with estimations of their intensities.

D. Model Creation and Assessment

We create a DL model for the unbiased appraisal of TC's strength using satellite pictures and CNN utilizing this training dataset. It took several rounds of the ML's different stages for the CNN's development to reach the conclusive model. Further iterations with the specimen after Pradhan's 2018 study showed that unbent output at 1 kt wind speeds performed better for image categorization to Saffir-Simpson scale storm severity and 5 kt wind speed intervals. As a result, this model takes in learning samples at intervals of 5 kt and outputs the highest momentum at

resolution of 1 kt; the clarity of the model, however, cannot be exceeding the 5 kt detail level of the input training data. For every storm that occurred in the East Pacific maritime regions in 2018 and 2019, The overall (RMSE) was 13.62 knots.

E. System Design




Fig.3. System Design

F. Proposed Methodology LSTM an Additional Approach

Cyclones, formidable atmospheric occurrences with swirling winds and low-pressure centres, present challenges for precise forecasting during their dynamic trajectory. Understanding and predicting these ever-changing patterns are crucial for proactive disaster preparedness.




Fig.4. Implementation

Long Short-Term Memory, excels in gauging cyclone intensity during continuous motion by apprehending dependencies and temporal intricacies in sequential data.

Unlike traditional models, LSTM networks efficiently manage information, discerning intricate connections in dynamic cyclone behaviour. The model's recurrent nature enables effective time-series analysis, crucial for tracking continuously moving cyclones and providing precise intensity predictions. Its ability to alleviate the vanishing gradient problem enhances effectiveness in capturing cyclone intensity changes. The architecture, including memory cells and gating mechanisms, empowers the network in situations marked by continuous movement and evolving patterns.

To encapsulate, its capacity to handle sequential data and address training challenges positions it as a superior option for estimating intensity in continuously moving cyclones. Accurate model like this helps minimize damage and protectives.

G. Advanced Utilization of Visualization Portal Technology

The visualization portal enhances storm intensity estimation by integrating real-time data feeds, satellite visuals, and numerical weather models. Advanced techniques, like 3D mapping, allow users to explore complex atmospheric patterns. Customizable filters focus on essential parameters, supports accurate predictions. Collaborative features, including shared data and annotations, enhance communication among meteorologists.

Real-time Integration: Aggregating data from diverse sources, including satellites and weather stations, ensures access to the latest cyclone-related information.

Dynamic Visualization: Using advanced techniques like 3D mapping, the portal offers an immersive representation of meteorological data, enabling intuitive exploration of atmospheric patterns.

Customizable Filters: Users can apply tailored filters to focus on relevant parameters, aiding meteorologists in informed assessments.

Machine Learning: Integration with predictive analytics improves forecasting by analyzing historical data and identifying patterns for more accurate predictions.

Collaborative Tools: Facilitating collaboration, the portal allows data sharing and annotations among meteorologists and researchers, fostering a comprehensive understanding of cyclone behavior.

In summary, it employs cutting-edge tech for powerful intensity estimation, empowering users to interpret data, make accurate predictions, and collaborate effectively in monitoring evolving weather conditions.

H. Optimizing Cyclone Intensity Estimation CNN vs LSTM

Selection between CNN and LSTM hinges on data characteristics and inherent features of the problem. CNNs excel in spatial data analysis, ideal for tasks with satellite imagery capturing crucial spatial patterns. Conversely,

LSTMs handle sequential data adeptly, valuable for tracking cyclones' temporal evolution. In practical applications a synergistic approach combining CNN and LSTM known as hybrid models proves advantageous. CNNs extract features, informing LSTMs to capture temporal dependencies for accurate intensity predictions. Optimal model selection hinges on data type, intensity estimation goals, and overall architectural considerations.

IV. RESULT




Fig.5. Detecting Formation of early-stage eye




Fig.6. Satellite IR image of Cyclone michaung



Fig.7 Inspection of intensity gradient

V. CONCLUSION AND FUTURE WORK

A complete Deep Learning based method for predicting and estimating the real time and spontaneous occurring tropical cyclone by utilizing its wind velocity activated in real is provided in this study. New Convolutional neural network models were devised as component of the system to objectively assess the tropical

cyclones wind velocity utilizing solely satellite imagery. Features found in CAMs are compared with Dvorak (T-no) pictures to thoroughly ensure, assess the model and methodically move it into production. Additionally, we introduce a unique approach to initiate the workflow to give real-time Wind pace evaluations through simultaneous information platform and monitor assessment for future storms. Furthermore, we discover that developing an algorithm takes a lot less time compared to compiling sizable, trustworthy learning dataset models of pictures and corresponding feeds. It takes many repetitive procedures with updated data and parameters to deploy the ML model, is a difficult process.

The specimen was versioned correctly and handled similarly to source code from the standpoint of software engineering. Ultimately, a diversified team of ML specialists, experts in that particular domain, native users, software engineers in that specific field, user interface designers are required in effective Implementation of a complete Machine Learning project.

REFERENCES

- [1] E. Blake, E. Rappaport, and C. Landsea, "The deadliest, costliest, and most intense United States tropical cyclones from 1851 to 2006 ((and other frequently requested hurricane facts)," NOAA Technical Memo. NWSTPC-5, 2007.
- [2] Z. Huang, D. Rosowsky, and P. Sparks, "Long-term hurricane risk assessment and expected damage to residential structures," Reliability Engineering & System Safety, vol. 74, pp. 239-249, 2001.
- [3] V. Dvorak, "A technique for the analysis and forecasting of tropical cyclone intensities from satellite pictures." NOAA Technical Memo, NESS 36, 15 pp. 14, 1972.
- [4] T. Olander and C. Velden, "The current status of the UW-CIMSS Advanced Dvorak Technique (ADT)," 30th Conference on Hurricanes and Tropical Meteorology, American Meteorological Society, 2012.
- [5] T. Olander and C. Velden, "The Advanced Dvorak Technique: Continued Development of an Objective Scheme to Estimate Tropical Cyclone Intensity Using Geostationary Infrared Satellite Imagery," Weather and Forecasting, vol. 22, pp. 287-298, 2007.
- [6] M. Pineros, E. Ritchie, and J. Tyo, "Objective measures of tropical cyclone structure and intensity change from remotely sensed infrared image data," IEEE Transactions on Geoscience and Remote Sensing, vol. 46(11), 2008.
- [7] J. Hawkins, T. Lee, J. Turk, C. Sampson, J. Kent, and K. Richardson, "Real-time inter-net distribution of satellite products for tropical reconnaissance," Bulletin of American Meteorological Society, vol. 82, pp. 567- 578, 2001.
- [8] Y. Bengio and A. Courville, "Deep Learning of Representations," Handbook on Neural Information Processing, vol. 49, pp. 1-28, 2013.
- [9] A. Krizhevsky, I. Sutskever, and G. Hinton, "ImageNet Classification with Deep Convolutional Neural Networks," Advances in Neural Information Processing Systems, pp. 1097-1105, 2012.
- [10] S. Kim, H. Kim, J. Lee, S. Yoon, S. Kahou, K. Kashinath, and Mr. Prabhat, "Deep-hurricane-tracker: Tracking and forecasting extreme climate events," In Proceedings 2019 IEEE Winter Conference on Applications of Computer Vision (WACV), pp. 1761-1769, 2019.