

डायबेटिक रेटिनोपैथी विश्लेषण

एआई-समर्थित रेटिना मूल्यांकन रिपोर्ट

रिपोर्ट दिनांक:	27-07-2026	रिपोर्ट आईडी:	DEA2D708-945
पूर्वानुमान:	Moderate DR	विश्लेषण समय:	17:56:59

क्लिनिकल व्याख्या

- The current retinal imaging assessment reveals findings consistent with a diagnosis of Moderate Non-Proliferative Diabetic Retinopathy (NPDR). Specifically
- the presence of numerous intraretinal dot-and-blot hemorrhages
- scattered microaneurysms
- and several distinct cotton wool spots are observed within multiple quadrants of both retinas. Furthermore
- there is mild venous beading and evidence of intraretinal microvascular abnormalities (IRMA) noted
- indicating localized areas of capillary non-perfusion and increased vascular permeability without the development of new blood vessels.

रोग सारांश

- Moderate Diabetic Retinopathy signifies a significant stage of vascular compromise within the retina due to sustained hyperglycemia
- leading to more extensive microvascular damage than typically seen in mild DR. This stage is characterized by increased retinal ischemia and vascular leakage
- elevating the risk for future visual impairment and progression to more advanced forms of the disease. While vision may currently be unaffected
- the structural changes indicate a clear need for intensified management.

संभावित चिकित्सीय चर्चाएँ

- Primary medical concerns for Moderate DR include the potential for rapid progression to Severe Non-Proliferative Diabetic Retinopathy or even Proliferative Diabetic Retinopathy
- characterized by the growth of fragile new blood vessels that can lead to severe hemorrhage and retinal detachment. There is also an increased likelihood of developing diabetic macular edema (DME)

- a condition where fluid accumulates at the macula
- directly impacting central vision and requiring prompt intervention to preserve sight.

उपचार मार्गदर्शन

- Optimal management at this stage centers on stringent systemic control of underlying risk factors
- including aggressive management of blood glucose
- blood pressure
- and cholesterol levels. While direct ocular treatments such as laser photocoagulation or anti-VEGF injections are generally reserved for more advanced stages or the presence of significant macular edema
- close monitoring is essential. Collaboration with an endocrinologist or primary care physician is highly recommended to achieve and maintain target metabolic parameters.

जीवनशैली सफ़िरशें

- To mitigate further progression of diabetic retinopathy
- it is imperative to adopt and sustain a disciplined healthy lifestyle. This includes strict adherence to a diabetes-friendly diet
- engaging in regular moderate physical activity as advised by your physician
- and complete cessation of smoking
- which significantly exacerbates retinal damage. Consistent self-monitoring of blood glucose levels and active participation in your diabetes management plan are crucial.

अनुवर्ती परामर्श

- Given the findings of Moderate DR
- a comprehensive ophthalmic re-evaluation
- including dilated fundus examination and repeat retinal imaging
- is strongly advised within 3-4 months to monitor for any signs of progression. Earlier re-assessment is warranted if you experience any new visual symptoms
- such as blurred vision
- floaters
- or flashes of light
- to ensure timely intervention and preserve visual function.

चकित्सा अस्वीकरण

- This report is a professional assessment based solely on the provided retinal imaging and does not substitute for a complete clinical ophthalmic examination and diagnosis. It is intended for informational purposes only and should not be considered as definitive medical advice or treatment. Always consult with a qualified healthcare professional for personalized medical guidance

- diagnosis

- and treatment options pertaining to your specific health condition.

CareSense AI द्वारा उत्पन्न
पेशेवर नेत्र रोग विश्लेषण रिपोर्ट
संस्करण 1.0