

डायबेटिक रेटिनोपथी विश्लेषण

एआय-समर्थित रेटिना मूल्यांकन अहवाल

अहवाल दिनांक:	27-07-2026	अहवाल आयडी:	01F3E929-75A
पूर्वानुमान:	Moderate DR	विश्लेषण वेळ:	17:57:21

क्लिनिकल समज

- Retinal imaging reveals distinct features consistent with Moderate Diabetic Retinopathy
- characterized by numerous microaneurysms and intraretinal dot-blot hemorrhages distributed across multiple quadrants. Notably
- early signs of venous caliber irregularities and scattered cotton wool spots are observed
- signifying an advancement in microvascular damage without evidence of neovascularization or significant macular edema at this juncture. This stage indicates a heightened risk for further progression.

आजार सारांश

- Moderate Diabetic Retinopathy represents an intermediate stage of diabetic eye disease
- where sustained elevated blood glucose levels have led to more widespread damage to the small blood vessels within the retina. While vision may remain unaffected
- the increased number and severity of microvascular changes signal a greater potential for progression to more advanced stages
- including severe non-proliferative or proliferative forms of retinopathy.

संभाव्य वैद्यकीय चर्चा

- The primary concern with Moderate DR is the elevated risk of advancing to severe non-proliferative or proliferative diabetic retinopathy
- which carries a substantial threat of irreversible vision loss due to vitreous hemorrhage or tractional retinal detachment. Furthermore
- there is an ongoing risk for the development or worsening of diabetic macular edema (DME)
- even in the absence of current significant macula involvement
- which could impair central vision.

उपचार मार्गदर्शन

- Management at this stage is primarily centered on rigorous systemic control to mitigate disease progression. Strict optimization of glycemic control (HbA1c)
- blood pressure
- and lipid profiles is paramount
- often requiring close collaboration with an endocrinologist or primary care physician. While direct ocular interventions like laser or anti-VEGF injections are not typically indicated for uncomplicated moderate DR
- they may become necessary if macular edema develops or if the disease progresses to a severe or proliferative stage.

जीवनशैली शफिरसी

- Proactive lifestyle modifications are essential in slowing the progression of diabetic retinopathy and preserving vision. Adhering to a balanced
- low-glycemic diet
- engaging in regular physical activity
- maintaining a healthy weight
- and complete cessation of smoking are critical steps. These efforts directly contribute to better systemic health and can significantly impact the long-term health of your retinal microvasculature.

पुढील सल्ला

- Given the current diagnosis of Moderate DR
- consistent and timely ophthalmic follow-up is crucial
- generally recommended every 4 to 6 months
- or more frequently if deemed necessary by clinical assessment. These regular examinations will allow for diligent monitoring of retinal changes
- early detection of any progression to more severe stages or the onset of macular edema
- and timely intervention to preserve visual acuity.

वैद्यकीय अस्वीकरण

- This report provides an assessment based on the retinal imaging findings and does not constitute a substitute for a comprehensive clinical ophthalmological examination and personalized medical advice. It is imperative to discuss these findings with your treating ophthalmologist and other healthcare providers to formulate an individualized management plan. This information is for educational purposes and should not be used for self-diagnosis or self-treatment.

CareSense AI द्वारा तयार
व्यावसायिक डोळ्यांचा रोग विश्लेषण अहवाल
आवृत्ती 1.0