राष्ट्रीय कृषि मौसम परामर्श सेवा बुलेटिन

National Agromet Advisory Service Bulletin



शुक्रवार, दि. 21 अगस्त 2026

(अवधि: 21 अगस्त से 25 अगस्त 2026)



Friday, 21st August 2026



(For the period 21st August to 25th August, 2026)





कृषि मौसम विज्ञान प्रभाग,
भारत मौसम विज्ञान विभाग, पुणे
द्वारा जारी


Issued by
Agricultural Meteorology Division,
India Meteorological Department, Pune.


NOAA/VIIRS/BLENDED NDVI composite for the week number 33 (13.08.2026 to 19.08.2026) over Agricultural regions of India	Difference in NOAA/VIIRS/BLENDED NDVI between week number 33, 2026 and 32, 2026
	
Agricultural vigour was good at many places Jammu & Kashmir, Punjab & Himachal Pradesh; at a few places of Uttarakhand, West Uttar Pradesh & North Eastern states and at isolated places of Haryana, east Uttar Pradesh & Kerala.	Difference in agricultural vigour was good at few places of Uttarkhand, Himachal Pradesh, East uttar Pradesh, Bihar, North Eastern states, Sub Himalayan West Bengal & Vidarbha and at isolated places of Punjab, Haryana, West Uttar Pradesh, Jharkhand, Gangetic West Bengal, North Coastal Andhra Pradesh, Telangana & Konkan.


Sub-Divisionwise Rainfall Forecast and weather warning for next 5 days	
	



SUBDIVISION RAINFALL FORECAST FOR DAY 2
2026-08-22
Arabian Sea
Bay of Bengal
Indian Ocean
0 250 500 km



SUBDIVISION WEATHER WARNING FOR DAY 2
2026-08-22
Arabian Sea
Bay of Bengal
Indian Ocean
0 250 500 km



SUBDIVISION RAINFALL FORECAST FOR DAY 3
2026-08-23
Arabian Sea
Bay of Bengal
Indian Ocean
0 250 500 km


SUBDIVISION WEATHER WARNING FOR DAY 3
2026-08-23
Arabian Sea
Bay of Bengal
Indian Ocean
0 250 500 km









Legend	Category	% Stations
WS	Widespread/Most Places	76-100
FWS	Fairly Widespread/Many Places	51-75
SCT	Scattered/ A Few Places	26-50
ISOL	Isolated Places	1-25
DRY	No Rain	0

No heavy Rainfall	Heavy Rainfall / Snowfall
Heavy	
Very heavy	
Exceptionally Heavy	
No TS	Thunderstorm / Squall / Hailstorm
Light TS	
Moderate TS	
Severe TS / Very Severe TS / Hailstorm	

Weather Warning during next 5 days

21 August 2026 (Day 1)

- **Heavy to Very Heavy Rainfall** very likely at isolated places over East Madhya Pradesh.
- **Heavy Rainfall** very likely at isolated places over Andaman & Nicobar Islands, Arunachal Pradesh, Assam & Meghalaya, Chhattisgarh, East Uttar Pradesh, Haryana, Chandigarh & Delhi, Himachal Pradesh, Nagaland, Manipur, Mizoram and Tripura, Punjab, Uttarakhand, West Bengal & Sikkim and West Madhya Pradesh.
- **Thunderstorm accompanied with lightning & gusty winds(40-50kmph)** very likely at isolated places over Andhra Pradesh, North Interior Karnataka and Tamil Nadu Puducherry & Karaikal.
- **Thunderstorm accompanied with lightning & gusty winds(30-40kmph)** very likely at isolated places over Coastal Karnataka, Jammu-Kashmir-Ladakh-Gilgit-Baltistan-Muzaffarabad, Rajasthan, South Interior Karnataka and West Uttar Pradesh.
- **Thunderstorm accompanied with Lightning** very likely at isolated places over Chhattisgarh, Gangetic West Bengal, Madhya Pradesh and Uttarakhand.
- **Strong Surface Winds** very likely at isolated places over Andhra Pradesh and Karnataka.

22 August 2026 (Day 2)

❖ **Heavy to Very Heavy Rainfall** very likely at isolated places over East Madhya Pradesh.

- **Heavy Rainfall** very likely at isolated places over Assam & Meghalaya, Bihar, Chhattisgarh, East Uttar Pradesh, Haryana, Chandigarh & Delhi, Himachal Pradesh, Jammu-Kashmir-Ladakh-Gilgit-Baltistan-Muzaffarabad, Nagaland, Manipur, Mizoram and Tripura, Odisha, Punjab, Sub Himalayan West Bengal & Sikkim, Uttarakhand and West Madhya Pradesh.
- **Thunderstorm accompanied with lightning & gusty winds(40-50kmph)** very likely at isolated places over Andhra Pradesh, North Interior Karnataka and Tamil Nadu Puducherry & Karaikal.
- **Thunderstorm accompanied with lightning & gusty winds(30-40kmph)** very likely at isolated places over Coastal Karnataka, Jammu-Kashmir-Ladakh-Gilgit-Baltistan-Muzaffarabad, Rajasthan, South Interior Karnataka and West Uttar Pradesh.
- **Thunderstorm accompanied with Lightning** very likely at isolated places over Chhattisgarh, Madhya Pradesh and Uttarakhand.
- **Strong Surface Winds** very likely at isolated places over Andhra Pradesh and Karnataka.

23 August 2026 (Day 3)

- **Heavy Rainfall** very likely at isolated places over Arunachal Pradesh, Assam & Meghalaya, Bihar, Chhattisgarh, Coastal Karnataka, Jharkhand, Kerala & Mahe, Madhya Pradesh, Nagaland, Manipur, Mizoram and Tripura, North Interior Karnataka, Odisha, Sub Himalayan West Bengal & Sikkim and Uttarakhand.
- **Thunderstorm accompanied with lightning & gusty winds(40-50kmph)** very likely at isolated places over Andhra Pradesh, North Interior Karnataka and Tamil Nadu Puducherry & Karaikal.
- **Thunderstorm accompanied with lightning & gusty winds(30-40kmph)** very likely at isolated places over Coastal Karnataka, East Rajasthan, Jammu-Kashmir-Ladakh-Gilgit-Baltistan-Muzaffarabad and South Interior Karnataka.
- **Thunderstorm accompanied with Lightning** very likely at isolated places over Chhattisgarh, Madhya Pradesh and Uttarakhand.
- **Strong Surface Winds** very likely at isolated places over Andhra Pradesh and Karnataka.

24 August 2026 (Day 4)

- **Heavy to Very Heavy Rainfall** likely at isolated places over Odisha.
- **Heavy Rainfall** likely at isolated places over Arunachal Pradesh, Assam & Meghalaya, Bihar, Chhattisgarh,

Coastal Karnataka, Kerala & Mahe, Madhya Pradesh, Nagaland, Manipur, Mizoram and Tripura, North Interior Karnataka, Tamil Nadu Puducherry & Karaikal, Uttarakhand and West Bengal & Sikkim.

- **Thunderstorm accompanied with lightning & gusty winds(40-50kmph)** likely at isolated places over Andhra Pradesh and North Interior Karnataka.
- **Thunderstorm accompanied with lightning & gusty winds(30-40kmph)** likely at isolated places over Coastal Karnataka, East Rajasthan, Jammu-Kashmir-Ladakh-Gilgit-Baltistan-Muzaffarabad and South Interior Karnataka.
- **Thunderstorm accompanied with Lightning** likely at isolated places over Chhattisgarh, Madhya Pradesh and Uttarakhand.
- **Strong Surface Winds** likely at isolated places over Andhra Pradesh and Karnataka.

25 August 2026 (Day 5)

- **Heavy to Very Heavy Rainfall** likely at isolated places over Odisha.
- **Heavy Rainfall** likely at isolated places over Arunachal Pradesh, Assam & Meghalaya, Bihar, Chhattisgarh, Kerala & Mahe, Madhya Pradesh, Tamil Nadu Puducherry & Karaikal, Uttarakhand and West Bengal & Sikkim.
- **Thunderstorm accompanied with lightning & gusty winds(40-50kmph)** likely at isolated places over Andhra Pradesh.
- **Thunderstorm accompanied with lightning & gusty winds(30-40kmph)** likely at isolated places over Coastal Karnataka and South Interior Karnataka.
- **Thunderstorm accompanied with Lightning** likely at isolated places over Chhattisgarh, Madhya Pradesh and Uttarakhand.
- **Strong Surface Winds** likely at isolated places over Andhra Pradesh, Coastal Karnataka and South Interior Karnataka.

Warning for Fishermen

- As **Squally wind with speed 45-55 kmph gusting to 65 kmph** is likely to prevail along and off Somalia and Oman coasts, over most parts of westcentral adjoining some parts of southwest, and few parts of northwest & eastcentral Arabian Sea. **Squally wind with speed 45-55 kmph gusting to 65 kmph** is likely to prevail over few parts of Gulf of Mannar adjoining Comorin area and southwest Bay of Bengal adjoining south Sri Lanka coast, over some parts of southeast Bay of Bengal. **Squally Weather with wind speed 40-50 kmph gusting to 60 kmph** is likely to prevail over Andaman Sea on 21st; **Squally wind with speed 45-55 kmph gusting to 65 kmph** is likely to prevail along and off Somalia and Oman coasts, over most parts of westcentral adjoining some parts of southwest, and few parts of north & eastcentral Arabian Sea. **Squally wind with speed 45-55 kmph gusting to 65 kmph i**s likely to prevail over few parts of Gulf of Mannar adjoining Comorin area and southwest Bay of Bengal adjoining south Sri Lanka coast. **Squally Weather with wind speed 40-50 kmph gusting to 60 kmph** is likely to prevail off north Andhra Pradesh coasts adjoining some parts of westcentral Bay of Bengal, over Andaman Sea on 22nd; **Squally wind with speed 45-55 kmph gusting to 65 kmph** is likely to prevail along and off Somalia. **Squally Weather with wind speed 35-45 kmph gusting to 55 kmph** is likely to prevail along and off Kerala, Lakshadweep Area adjoining Comorin area. **Squally wind with speed 45-55 kmph gusting to 65 kmph** is likely to prevail over Gulf of Mannar adjoining Comorin area and southwest Bay of Bengal adjoining south Sri Lanka coast. **Squally Weather with wind speed 40-50 kmph gusting to 60 kmph** is likely to prevail along and off North Andhra Pradesh coasts, over some parts of westcentral Bay of Bengal, over Andaman Sea a and Oman coasts, over most parts of westcentral adjoining some parts of southwest, adjoining eastcentral Arabian Sea on 23rd; **Squally wind with speed 45-55 kmph gusting to 65 kmph** is likely to prevail along and off Somalia. **Squally Weather with wind speed 35-45 kmph gusting to 55 kmph** is likely to prevail along and off Kerala, Lakshadweep Area adjoining Comorin area. **Bay of Bengal: Squally wind with speed 45-55 kmph gusting to 65 kmph** is likely to prevail over Gulf of Mannar adjoining Comorin area and

southwest Bay of Bengal adjoining south Sri Lanka coast. **Squally Weather with wind speed 40-50 kmph gusting to 60 kmph** is likely to prevail along and off North Andhra Pradesh coasts, over some parts of westcentral Bay of Bengal, over Andaman Sea a and Oman coasts, over most parts of westcentral adjoining some parts of southwest, adjoining eastcentral Arabian Sea on 24th and **Squally wind with speed 45-55 kmph gusting to 65 kmph** is likely to prevail along and off Somalia and Oman coasts, over most parts of westcentral adjoining some parts of southwest, adjoining eastcentral Arabian Sea. **Squally Weather with wind speed 35-45 kmph gusting to 55 kmph** is likely to prevail along and off Kerala, Lakshadweep Area adjoining Comorin area. **Squally wind with speed 45-55 kmph gusting to 65 kmph** is likely to prevail over Gulf of Mannar adjoining Comorin area and southwest Bay of Bengal adjoining south Sri Lanka coast on 25th August, 2026, **fishermen are advised not to venture into these areas.**

Salient Advisories

- As **rain/thundershowers at many/most places** likely over Arunachal Pradesh and Himachal Pradesh during next 48 hours, over Jammu & Kashmir during next 72 hours, over Assam & Meghalaya, Nagaland- Manipur-Mizoram-Tripura, Uttarakhand, West Bengal & Sikkim, Jharkhand, Chhattisgarh, Madhya Pradesh, Konkan & Goa, Coastal Karnataka, Keralam, Lakshadweep and Andaman & Nicobar Islands during next 4-5 days, over East Uttar Pradesh and Bihar during 22nd to 24th and **Heavy to very heavy rainfall** is likely at isolated places over East Madhya Pradesh during next 48 hours, over Odisha on 24th & 25th; **Heavy rainfall** is likely at isolated places over Andaman & Nicobar Islands during next 24 hours, over East Uttar Pradesh, Punjab, Haryana, Himachal Pradesh during next 48 hours, over North Eastern State, Sub-Himalayan West Bengal, West Madhya Pradesh Chhattisgarh & Uttarakhand during next 4 to 5 days, over Jammu & Kashmir on 22nd, over Odisha on 22nd & 23rd, over Coastal & North Interior Karnataka on 23rd & 24th, over Kerala & East Madhya Pradesh during 23rd to 25th, over Jharkhand on 23rd and over Tamil Nadu on 24th & 25th August, 2026, postpone irrigation, intercultural operations and application of plant protection measures & fertilizers to the standing crops. Provide adequate drainage in standing crop fields to avoid waterlogging.
  - **East Madhya Pradesh -** Ensure proper drainage in soybean, cotton, maize and sugarcane fields so as to prevent waterlogging. Avoid transplanting of vegetables during heavy rainfall period. Provide support to young fruit plants. Postpone planting of new saplings. In tall sugarcane crops, carry out propping (tying adjacent canes together) to minimize lodging.
  - **Odisha** - Make necessary arrangements to drain out excess water from *kharif* rice, maize, pulses, cotton, pigeon pea, banana, vegetable fields and nursery beds. Keep field drains, outlets and bunds clear and functional. In waterlogged areas, postpone sowing/re-sowing of maize and green gram and prevent water stagnation in pearl millet. Provide staking/support to vegetables, banana, papaya and new horticultural plants.
  - **Gangetic West Bengal -** In **Laterite and red soil zone**, in areas prone to water accumulation, prepare broad bed and furrow systems to improve drainage and aeration for groundnut. Harvest matured crops promptly when conditions permit and store in a dry, safe place.
- As **Thunderstorm accompanied with lightning & gusty winds(40-50kmph)** very likely at isolated places over Tamil Nadu Puducherry & Karaikal during next 72 hours, over North Interior Karnataka during next 4 days, over Andhra Pradesh during next 5 days; **Thunderstorm accompanied with lightning & gusty winds(30-40kmph)** very likely at isolated places over West Uttar Pradesh during next 48 hours, over Jammu-Kashmir-Ladakh-Gilgit-Baltistan-Muzaffarabad and West Rajasthan during next 4 days, over South Interior Karnataka and Coastal Karnataka during next 5 days, over East Rajasthan on 23rd & 24th August, 2026; provide mechanical support to banana & papaya plantations and staking to young plants and vegetables. Undertake propping in sugarcane.
- **Sowing/Transplanting/Harvesting**
  - sowing of irrigated maize & fodder sorghum; transplanting of rice and nursery sowing of *Samba* rice in

Tamil Nadu

- harvesting of matured green gram; transplanting of rice & chilli; planting of areca nut in Karnataka
- sowing of green gram, black gram & horse gram; planting of papaya, turmeric & ginger in Chhattisgarh
- sowing of pigeon pea, cow pea & castor; planting of banana in Gujarat
- sowing of carrot & radish; transplanting of cauliflower and planting of citrus, mango, litchi, guava, loquat, *ber and amla* in Punjab
- harvesting of mature brinjal, capsicum & tomato; sowing of buckwheat, oats & *toria* in Himachal Pradesh
- sowing of pearl millet, sesame, pigeon pea & aromatic rice; transplanting of brinjal, tomato, chilli & early cauliflower in Uttar Pradesh
- transplanting of *Aman* rice in Gangetic West Bengal
- transplanting of scented rice in Bihar
- harvesting of jute; sowing of pigeon pea, green gram, black gram, cow pea & sesame in Assam

➢ **Pest and Disease**

- In Odisha, spray Fipronil 5% SC @ 400ml or Profenophos 50% EC @ 400ml or Chlorantraniliprole 18.5% SC @ 60ml per acre during clear weather to control leaf folder and case worm infestation in paddy.
- In Tamil Nadu, spray Carbosulfan @ 400ml/ac or Cartap hydrochloride @ 600g/ac during morning or evening hours to manage leaf folder and stem borer infestation in paddy.
- In Karnataka, spray Hexaconazole @ 1ml / litre water during clear weather to control powdery mildew in green gram and black gram
- In Gujarat, spray Imidacloprid 17.8% SL @ 3-4 ml in 10 litres of water to control aphids, leaf hopper and mealy bug in cotton.
- In Punjab, spray Azoxystrobin 18.2% + Difenoconazole 11.4% SC @ 100 ml in 200 litres of water during clear weather as soon as banded leaf and sheath blight appears in maize field.
- In Chhattisgarh, apply Beta Cyfluthrin + Imidacloprid @ 350ml/ha or Thiamethoxam + Lambda cyhalothrin @ 125 ml/ha after rainfall subsides to manage attack of leaf-eating caterpillars, white fly and stem fly in soybean.

Detailed Agromet Advisories

SOUTH INDIA [TAMIL NADU, TELANGANA, ANDHRA PRADESH, KARNATAKA, KERALAM]

Realised Rainfall: Significant amount of rainfall occurred in all the districts of Kerala except Alappuzha, Kollam and Thiruvanantapuram; in Coimbatore, Sivaganga districts of Tamilnadu; in Alluri Sitharamraju, B.R. Ambedkar Konaseema, East Godavari, Eluru, Yanam districts of Andhra Pradesh; B. Kothagudem, Kamareddy, Mahabubabad, Mancherial, Ranjanna Sircilla, Warangal districts of Telangana and Dakshina Kannada, Udupi, Uttara Kannada, Chikkamagaluru, Kodagu & Shivamogga districts of Karnataka.

Advisories:

➢ As **rain/thundershowers at many/most places** likely over Coastal Karnataka, Keralam, Lakshadweep during next 4-5 days and **Heavy Rainfall** very likely at isolated places over Coastal Karnataka & North Interior Karnataka on 23rd & 24th, over Keralam during 23rd to 25th and over Tamil Nadu Puducherry & Karaikal on 24th & 25th August, 2026; postpone irrigation, intercultural operations and application of plant protection measures & fertilizers to the standing crops. Provide adequate drainage in standing crop fields to avoid waterlogging.

➢ As **Thunderstorm accompanied with lightning & gusty winds(40-50kmph)** very likely at isolated places over Tamil Nadu Puducherry & Karaikal during next 72 hours, over North Interior Karnataka during next 4 days, over Andhra Pradesh during next 5 days, **Thunderstorm accompanied with lightning & gusty winds(30-40kmph)** very likely at isolated places over South Interior Karnataka and Coastal Karnataka during next 5 days; provide mechanical support to banana & papaya plantations and staking to young plants and

vegetables. Undertake propping in sugarcane.

Tamil Nadu

- In North Eastern Zone, continue transplanting of rice varieties utilizing the expected rainfall and provide proper drainage facility to the nursery field. Due to prevailing weather condition leaf folder and stem borer infestation is found in paddy crop, to manage them, spray Carbosulfan @ 400ml/ac or Cartap hydrochloride @ 600g/ac during morning or evening hours.
- In Western Zone, continue sowing of irrigated maize for reaping potential yield and fodder sorghum. Also anticipating drizzling rainfall, prepare raised bed samba season nursery for sowing of medium duration rice. Prevailing high RH and cloudy weather condition is favorable for the tikka leaf spot incidence in groundnut and monitor the incidence; if necessary, spray Carbendazim @ 500g or Mancozeb @ 1000g per ha.
- In Cauvery Delta Zone, considering favorable weather, continue nursery preparation & sowing for samba rice. Also, by utilizing weather condition land can be prepared for samba rice cultivation before rain.
- In High Altitude Hilly Zone, to manage wilt in pepper, apply Trichoderma Asperellum / Bacillus subtilis @ 25g and Neem cake @ 500g / vine is recommended. Applications can be made during the rain-free period.
- In High Rainfall Zone, weather is congenial for spiraling white fly in coconut. Install Yellow sticky traps @ 5/acre and spray neem oil 3% along with sticking agent.
- In Coastal Zone, to manage Phomopsis blight in brinjal, spray Pseudomonas fluorescens or Bacillus subtilis @ 5g or 5ml/ litre water.

Animal Husbandry

- In Tamil Nadu, due to very high and prolonged day temperatures, farmers are advised to feed their cattle and small ruminants with sufficient quantities of concentrates and green fodder for the maintenance of body weight and milk production.

Keralam

- In High Altitude Zone, present weather is favorable for incidence of soft rot and shoot borer in ginger, drench soil with Copper oxychloride 50WP @ 2g/litre of water or Copper hydroxide with 50WP @ 1.5g /litre of water against soft rot disease. Spray Chlorantraniliprole 18.5 SC @ 3ml/10 litres of water against shoot borer.
- In Southern Zone, weather is congenial for rice bug infestation in paddy from the flowering to maturity stages. As a precautionary measure, keep the field and bunds free from weeds and grasses. If rice bug infestation is observed, spray fish-jaggery extract @ 20ml per litre of water. In case of severe infestation, spray Cypermethrin 10% EC @ 0.5 ml per litre of water during clear weather.
- In Central Zone, weather is congenial for leaf fall in nutmeg. As a prophylactic measure, spray 1% Bordeaux mixture. Proper drainage facilities have to be ensured.
- In Northern Zone, weather is favourable for leaf folder in paddy, spray Flubendiamide 39.35% SC @ 1ml/10 litres of water. Avoid excessive Nitrogenous fertilizers.
- In Problem Area Zone, ensure good drainage around palms. For bud rot in coconut, remove affected tissues and apply 1% Bordeaux mixture as a protective measure according to recommendations.

Andhra Pradesh

- In High Altitude tribal Zone, apply gypsum @ 200 kg/acre at flowering stage in groundnut and sugarcane at grand growth stage. To reclaim the Iron deficiency in the rice, spray with ZnSo4 @ 5g/L. To avoid moisture stress in nursery due to prolonged dry spells, spray multi K or 19:19:19 @ 1g/L during morning or evening times.
- In Scarce rainfall zone, where dry weather conditions continue, spray 19:19:19 @ 5 g/litre of water as a foliar spray in standing groundnut, castor, cotton, maize and red gram to minimize moisture stress. The current weather conditions are favourable for sucking pest incidence. Advised to setting up of yellow sticky traps @ 10 per acre for monitoring whitefly incidence and blue sticky traps @ 10 per acre for monitoring thrips incidence.
- In North Coastal Zone Zone, continue transplanting of rice by clipping the leaf tips of rice nurseries. Conserve soil moisture through intercultivation in red gram. Due to prevailing dry conditions, spray 19:19:19

@ 5-10g per litre of water as a foliar spray to help the standing crops of red gram, groundnut and direct sown rice.

- In Krishna Godavari Zone, undertake intercultivation /hand weeding to control weeds and conserve soil moisture in red gram at 25–30 DAS. Where chemical control is required, apply Imazethapyr @ 750 mL/ ha at around 3 weeks after sowing.
- In Southern Zone, due to continuous dry spells, if the leaves of the paddy crop show wilting symptoms, spray multi K @ 10g or Potassium nitrate @ 10 g/ L of water in 15 days interval on need basis

Telangana

- In Northern Telangana, sesame sowing can be done up to 30 August. For late Kharif sowing, suitable varieties such as JCS 2854, JCS 3202, JCS 3287 and Swetha may be selected. Farmers are advised to install yellow sticky traps @ 10 traps per acre to monitor the incidence of sucking pests in cotton.
- In Southern Telangana, undertake top dressing of Nitrogen (25 kgs of Urea/acre) and Potassic (10 kgs MOP/acre) fertilizers at 20, 40 and 60 DAS after receipt of rains in rainfed crops. In rainfed mango orchards go for mulching around the tree using local available material to conserve the moisture and to control the weeds. Advised to go for the spraying of micro nutrient mix Formula 4 @ 2g per liter of water on newly emerged flush of leaves.

Karnataka

- In Hill Zone, continue transplanting of paddy and planting of healthy areca nut saplings during suitable field conditions; continue raising areca nut saplings. Provide the drainage in maize field to remove the excess water from the field. Ensure proper drainage in ginger and remove excess water from maize fields. For control of *katte roga* in black pepper as a precautionary measure drench copper sulphate solution @ 2 to 4 litre per vine (@ 3g/litre of water). After that spray 1% Bordeaux mixture solution.
- In Coastal Zone, due to continuous rainfall there may be chances of *Kole roga* in arecanut, spray 1% Bordo mixture in clear weather condition to avoid yield losses. Provide drainage facility to drain excess rain water from black pepper, arecanut and coconut. To manage T-Mosquito bug in cashew, spray Monocrotophos @ 1.7ml /litre or Lambda cyhalothrin @ 0.5ml/litre. Conserve available soil moisture in paddy by strengthening field bunds and minimizing water losses.
- In North East Dry Zone, undertake paddy and chilli transplanting where adequate soil moisture and suitable field conditions are available. Spray Hexaconazole @ 1ml / litre water for controlling powdery mildew in green gram and black gram.
- In North Dry Zone, to manage moisture in pigeon pea, pearl millet and maize crops at vegetative stage, take up repeated intercultivation to conserve soil moisture. Due to cloudy condition and rainfall, there is chance of occurrence of Anthracnose disease in grapes, spray Thiophanate methyl @ 1g /litre of water. To manage mealy bug in grapes, spray Imidacloprid 17.8 S.L. @ 0.4 ml/litre of water.
- In North Transition Zone, harvest early-sown green gram during rain-free periods and protect harvested produce from rainfall. Provide light irrigation to vegetables only where required based on soil-moisture conditions. Spray Hexaconazole/Propiconazole @1ml per litre of water to manage rust in sugarcane. Undertake intercultivation in black gram, soybean, maize and groundnut for weed control and soil-moisture conservation.
- In Central Dry Zone, complete sowing and transplanting operations wherever adequate soil moisture is available. Carry out intercultivation and earthing-up in maize to support plants, suppress weeds and conserve soil moisture.
- In Eastern Dry Zone, undertake sowing of finger millet if sufficient soil moisture available in crop field. Continue intercultural operations in standing crops. Based on field conditions, provide protective irrigation to early-sown crops where rainfall/soil moisture is inadequate.
- In South Transition Zone, undertake transplanting of paddy due to good rainfall received during the last week. Maintain enough standing water in the already transplanted paddy. Prevailing weather favours Mahali (fruit rot) in arecanut, remove fallen infected nuts and spray 1% Bordeaux mixture or Copper oxychloride @ 3g/L

during a rain-free period. Continuous rainfall and high humidity favour rhizome rot and leaf spot in ginger. Ensure proper drainage and drench with Metalaxyl + Mancozeb @ 2g/L or Copper oxychloride @ 3g/L if symptoms appear. For managing root rot provide adequate drainage to remove excess of water from the field.

- In Southern Dry Zone, in paddy, provide protective irrigation wherever irrigation is assured. For delayed transplanting, avoid long-duration varieties without assured water and prefer short-duration varieties where suitable. In ginger, avoid excessive irrigation and maintain suitable soil moisture around the rhizome zone. Undertake earthing-up and moisture conservation in maize. If severe moisture stress persists and recovery is poor, utilize the crop as green fodder or silage. Provide protective irrigation to standing crops only where required and adopt soil-moisture conservation through mulching, intercultivation and field-bund maintenance.

NORTHEAST INDIA [ARUNACHAL PRADESH, ASSAM & MEGHALAYA, NMM&T]

Realized rainfall: Significant rainfall occurred over Changlang, East Kameng, East Siang, Lohit, Lower Subansari, Papumpare, Tawang, Tirap, Upper Siang and West Siang districts of Arunachal Pradesh; over Baksa, Biswanath, Bongaigaon, Cachar, Charaideo, Darrang, Dhemaji, Dhubri, Dibrugarh, Goalpara, Golghat, Hailakandi, Hojai, Jorhat, Kamrup (Metro), Karbi Anglong, Kokrajhar, Lakhimpur, Morigaon, Sonitpur, Sibsagar, Sribhumi, Tinsukia, Udalgiri and West Karbi Anglong districts of Assam; over East Garo Hills, East Jaintia Hills, East Khasi Hills, Ri-bhoi, South Garo Hills, South West Khasi Hills, West Khasi Hills and West Jaintia Hills districts of Meghalaya; over Aizwal, Kolasib, Lawngtlai and Saiha districts of Mizoram; over Bishnupur, Imphal West and Tamenglong districts of Manipur; over Dimapur, Kohima, Mokokchung, Mon, Paren, Phek, Tuensang, Wokha, Zunheboto districts of Nagaland and over North Tripura and South Tripura districts of Tripura.

Advisories:

- As **rain/thundershowers at many/most places** likely over Arunachal Pradesh during next 48 hours, over Assam & Meghalaya and Nagaland- Manipur-Mizoram-Tripura during next 4-5 day and **Heavy Rainfall** very likely at isolated places over the region during next 5days, postpone irrigation, intercultural operations and application of plant protection measures & fertilizers to the standing crops. Provide adequate drainage in standing crop fields to avoid waterlogging.

Arunachal Pradesh

- Ensure proper drainage in rice, maize, finger millet, soybean, cowpea, vegetables and orchards to remove excess water and prevent waterlogging following rainfall. Millets during panicle initiation and flowering stage should be kept well-drained during light to moderate rain, with proper spacing and regular weeding. Monitor for ear head bugs and caterpillars, remove visible insects and maintain field sanitation. If infestation appears, spray neem seed kernel extract (NSKE) 5% (50 g crushed neem seeds/litre water) during dry weather. Regularly monitor Jhum paddy for blast, particularly for eye-shaped spots with grey centres and brown margins on leaves. Maintain proper spacing and drainage and avoid excess nitrogen to reduce disease risk. If symptoms appear, spray Trichoderma viride @ 5g/litre water during clear weather.

Assam

- In Hill Zone, if sugarcane crop shows symptoms of red rot and stem borer, rouged out and buried or burned the infested plants. Adequate drainage should be provided in the sugarcane plot. During the tillering stage, monitor for stem borer and case worm; if infestation is observed, spray Fipronil 5 SC @ 50 ml/bigha or Thiamethoxam 25 WG @ 13 g/bigha only during a rain-free period.
- In Lower Brahmaputra Valley Zone, continue sowing of pigeon pea, green gram and sesame wherever field conditions are suitable. Harvest jute at the small pod stage for better fibre quality. Monitor *Sali* rice for rice swarming caterpillar; undertake hand collection or kerosene application at the initial stage, and if infestation exceeds 25% of the field, spray Fipronil 5 SC @ 2ml /litre of water only during clear weather. Maintain proper drainage in crop fields.
- In Upper Brahmaputra Valley Zone, complete sowing of sesame, green gram and black gram by the end of August. Farmers may also undertake direct sowing of pre-germinated seeds in puddled fields.
- In Barak Valley Zone, continue sowing of sesame, pigeon pea, green gram and black gram under favourable

field conditions. Adopt direct seeding of short-duration rice varieties such as Luit and Disang to minimize the risk of delayed crop establishment and moisture stress. Due to the prevailing and predicted moderate rainfall over the next five days, postpone top-dressing of urea. During fair weather, apply Urea @ 4.5 kg/bigha for semi-dwarf varieties and 1.5 kg/bigha for tall varieties at 20–30 days after transplanting. Application should preferably be done during the morning or evening hours.

- In Central Brahmaputra Valley Zone, continue sowing of sesame, pigeon pea, green gram and black gram under suitable field conditions. Continue nursery-bed preparation and sowing of early cauliflower and radish. Continue harvesting of jute. Ensure proper drainage in crop fields and avoid fertilizer or pesticide application during rainfall or thunderstorms.
- In North Bank Plain Zone, continue sowing of sesame, pigeon pea, cow pea, green gram and black gram. August is very conducive for heavy infestation of rice (which is at tillering stage) with Rice Hispa, army worm and other insect pest. Spray NSKE (Neem Seed Kernel Extract) 5% (4 ml/litre) twice. Release Trichogramma japonicum to control stem borer and leaf folder. Release it 6 times, at 10-day intervals. Start the first release at 25 days after transplanting. Use about 6,700 Trichogramma per bigha for each release.

Meghalaya

- Harvest matured cobs of maize during available clear weather periods and dry the harvested maize promptly to avoid deterioration due to continuous humid/rainy conditions. In transplanted rice, maintain a thin water layer of 2–3 cm and keep field bunds properly repaired to retain the desired water level. Ensure proper drainage and remove excess water from maize, cowpea, okra and ginger fields to prevent waterlogging, particularly after rainfall.

Manipur

- Undertake harvesting of colocasia. Harvest mature ginger rhizomes during available rain-free periods and ensure proper drainage in ginger fields to prevent water stagnation. Monitor ginger for soft rot and, where infestation is observed, apply Metalaxyl or Ridomil @ 1ml /litre of water during a suitable rain-free period. Monitor Kharif rice at the tillering stage for leaf folder and thrips, apply Fipronil 5% SC for control where infestation is observed, preferably during a suitable rain-free window.

Mizoram

- Start field preparation for sowing of black gram, green gram, soybean, sesamum and lentil in harvested fields of maize. Due to cloudy weather condition & high humidity, initial incidence of blast may occur in lowland rice, apply Tricyclazole @ 0.6g per litre of water during sunny days. Apply Cypermethrin 25 EC @ 2ml /litre of water during sunny days in case of heavy infestation of gundhi bug in Upland or *Jhum* rice. Continuous wet weather conditions are favorable for the spread of soft rot and rhizome rot in ginger. Avoid water stagnation and remove affected clumps in case of heavy infestation, and drench the affected and surrounding areas with Mancozeb 75% WP @ 3g per litre of water.

Nagaland

- Maintain 3 to 5 cm water level in wet rice cultivation fields. Undertake harvesting of matured cobs of maize. Prevailing humid weather with frequent rain and cloudy days, is favourable for leaf spot during the vegetative stage of soybean. Bury immediately or burn the infested plant.

Tripura

- Adequate soil moisture to be maintained during panicle initiation and booting stage of *Aus* rice, as these are critical water requirement stages of rice. A shallow water layer of about 2–5 cm, depending on field condition and water availability to be maintained in the field. There is chance of attack of Bacterial Leaf Blight (BLB) in *Aman* rice which may stunned or reduce the Tillering. Manage the attack by fresh cow dung extract 20% twice (starting from initial appearance of the disease and another at fortnightly interval). In groundnut crop, maintain the drainage particularly in lowland where chances of stagnant water due to rain is more. There may be attack of White Grub on groundnut. The attack may be controlled by soil application of 10% Phorate @ 10 kg/ha.

Animal Husbandry

- Prevailing temperature may uncomfortable to the poultry birds. Ensure proper ventilation in poultry houses and also protect birds from heavy rains. Make sure to deworm older flocks every month against round-worms. Replace the litters frequently and keep the houses clean and ensure clean drinking water to the birds. Increase protein by 2% in poultry diet. Ensure vaccination against Ranikhet disease.

EAST INDIA [JHARKHAND, BIHAR, ODISHA, WEST BENGAL & SIKKIM, ANDAMAN & NICOBAR ISLANDS]

Realised Rainfall: Significant rainfall occurred in all districts of Odisha**;** all districts of Jharkhand except Godda & Sahebganj; in all districts of Andaman & Nicobar Islands except Nicobar; in all districts of West Bengal & Sikkim except Cooch Behar, North & South Dinajpur and Malda; in Arwal, Aurangabad, Bhabua, Buxar, Gaya, Gopalganj, Jahanabad, Monghyar, Nalanda, Patna, Rohtas, Saran, Siwan and Vaishali districts of Bihar.

Advisories:

- As **rain/thundershowers at many/most places** likely over West Bengal & Sikkim, Jharkhand, Andaman & Nicobar Islands during next 4-5 days, over Bihar during 22nd to 24th and **Heavy to Very Heavy Rainfall** very likely at isolated places over Odisha on 24th & 25th; **Heavy Rainfall** very likely at isolated places over Andaman & Nicobar Islands during next 24 hours, over Gangetic West Bengal on 21st, 24th & 25th, over Sub Himalayan West Bengal & Sikkim during next 5 days, over Odisha on 22nd & 23rd, over Bihar during 22nd to 25th, over Jharkhand on 23rd August, 2026; postpone irrigation, intercultural operations and application of plant protection measures & fertilizers to the standing crops. Provide adequate drainage in standing crop fields to avoid waterlogging.

Bihar

- In North East Alluvial Zone, after harvesting of mango, apply 1 kg urea + 500g phosphorus + 500 g potash + 20–25 kg well-decomposed cow dung manure to trees above 10 years old by placing the inputs in a 1-metre-wide circular trench around the tree.
- In North West Alluvial Plain zone and North East Alluvial Zone, if transplanted paddy seedlings in the field have died due to dry weather, transplant new plants in the vacant spots. If 25–30 days have passed since transplantation, apply the recommended dose of urea for the first top-dressing @ 25 kg per acre.
- In the South Bihar Alluvial Zone, complete the transplanting of scented rice varieties during this month. Apply algal culture @ 10 kg/ha in the transplanted rice fields. Maintain adequate standing water in the transplanted rice fields. Apply the remaining dose of nitrogen as urea after removing weeds from the maize field for proper growth & yield.

Jharkhand

- In the Western Plateau Zone, keep the harvested produce at safer places. Maintain proper drainage and prevent waterlogging in all crops. Cover the vegetable seedling with polythene to protect them from heavy rainfall. To protect rice crop from stem borer infestation, remove and destroy 'dead hearts' and white ears in the field. In case of severe infestation, spray Chlorantraniliprole 18.5% SC @ 0.3 ml per litre of water; avoid excessive use of nitrogen (urea). Drain out excess rainwater from red gram to avoid waterlogging; drench the base of affected or early wilting plants with Carbendazim @ 1g per litre of water during clear weather. Monitor the incidence of fruit fly, in bottle guard, sponge guard, bitter guard, cucumber & ridge guard, infected fruits should be picked and buried in the soil; use 20-25 Pheromone trap per acre. Maintain proper drainage in maize, monitor crop regularly for stem borer symptoms; remove and destroy severely affected plants and if infestation is severe, apply Carbofuran 3G @ 3 Kg per acre in the centre whorls of the plant.

Odisha

- In Western Undulating Agroclimatic Zone, postpone sowing operations, top dressing of fertilizers and herbicide application until excess moisture drains and suitable dry weather prevails. Hoeing, weeding and 1st top dressing followed by earthing up at the three-week stage of cotton should be undertaken only when the soil becomes workable. Monitor maize fields regularly and remove infected plant parts.
- In East & South Eastern Coastal Plain Zone, go for gap filling in rice crop using seedling of same age or clonal tillers to have a uniform distribution. Make arrangements for draining out excess water from the field

of Pulses, maize, groundnut and vegetables. Complete all inter-culture operation in groundnut, before flowering starts. For control of leaf eating caterpillar in black gram and green gram, apply neem oil formulation 300 ppm @ 1 litre per acre.

- In West Central Table Land Zone, ensure proper drainage in groundnut fields, as the crop is highly sensitive to stagnant water. In rice, install 4–5 pheromone traps per acre to monitor yellow stem borer (Scirpo lure) and leaf folder (Cnaphalo lure); if male moth population reaches 4–5 per trap, apply the recommended insecticide only after rainfall subsides, under clear and calm weather conditions. Ensure proper drainage in orchard and avoid prolonged water stagnation around the tree basin.
- In North Eastern Ghat Zone, postpone beushaning and urea application in direct-seeded rice until excess water drains and field conditions become suitable. In early-sown maize, postpone hand weeding, earthing up and application of urea @ 55 kg/acre at 20–25 DAS until the soil becomes workable after rainfall.
- In North Central Plateau Zone**,** drain out excess water from the green gram field, remove the wilted plants and destroy them away from the field. After cessation rain, keep yellow sticky traps in the rice field @ 12-15 nos. of cards per ha, to control insect population. Postpone the transplanting of tomato in water logging areas, let the field dry before transplanting.
- In Mid Central Table Land Zone, due to cloudy weather there is a chance of leaf folder and case worm infestation in paddy, spray Fipronil 5% SC @ 400ml per acre or Profenophos 50% EC @ 400ml per acre or Chlorantraniliprole 18.5% SC @ 60ml per acre during clear weather. To manage Maydis Leaf Blight disease in maize, spray Mancozeb 75 % WP @ 400g per litre or Zineb 75 % WP @ @ 400g per litre or Carbendazim 12 % + Mancozeb 63 % WP @ 400g per litre. Carry out thinning and hand weeding if the sesame crop is 15 days old; apply 6 to 8 kg of nitrogen per acre.
- In South Eastern & Eastern Ghat High Land Zones**,** monitor rice for stem borer and leaf folder and, when pest population reaches ETL, apply Chlorantraniliprole 18.5 SC @ 150 ml/ha only during a suitable clear and calm weather window. Ensure drainage from maize fields to prevent waterlogging and root damage. Clean all drainage channels immediately in ginger, turmeric, vegetables, pulses and low-lying maize fields. Provide staking/support to tomato, chilli and other tall crops against strong wind.

Sub-Himalayan West Bengal & Sikkim

- In Hill Zone, clear drainage channels in vegetable fields, orchards, rice and finger millet nurseries to prevent waterlogging. Closely monitor tomato, chilli, ginger and cucurbits for bacterial wilt, damping-off and other fungal diseases under prolonged humid conditions. In chilli, for fruit rot/anthracnose, apply Azoxystrobin 23 EC @ 1ml/litre of water or Copper Oxychloride 50 WP @ 2.5 g/litre of water with sticker only during clear weather. To minimize fruit drop in mandarin orange, spray Planofix @ 1 ml/4.5 litre of water only when weather permits.
- In Terai Zone, in Aman paddy affected by blast, avoid top dressing with nitrogenous fertilizer during the prevailing wet conditions. Do not start retting matured jute in running water and use stagnant water in ponds or small water bodies. Protect chilli seedlings with plastic cover from heavy rain. Improve drainage/raise soil around bitter gourd plants to prevent water stagnation. To save the okra crop from heavy rain, raised up the ground level near the stem with soil, and support the plant with stick or PVC pipe.
- In Old Alluvial Zone, maintain proper drainage in rice and vegetable fields. In rice, undertake weed management and judicious nitrogen application only when field conditions become suitable. For brinjal fruit and shoot borer, spray Cartap Hydrochloride 50% SP @ 1 g/litre of water only during a clear-weather window. For the attack of thrips and mites in chilli crop, spray Spiromesifen 22.9% S.C. @ 0.75 ml per litre of water during afternoon.

Gangetic West Bengal

- In New Alluvial Zone, for fungal diseases in rice such as brown spot, spray Carbendazim 12% + Mancozeb 630% WP @ 2g per litre of water only when disease symptoms appear. Harvesting of jute may be taken up at appropriate maturity. Drainage channels should be maintained before sowing of mung and kalai, which are highly susceptible to waterlogging.

- In the Coastal Saline Zone, Aman rice transplanting may continue only where field conditions permit; retain a portion of seedlings in the nursery for gap filling/re-transplanting if required, avoid urea application in the main field during the active wet spell, and keep drainage channels clear and weed-free to rapidly remove excess water from furrows.
- In Laterite and red soil zone, during transplanting of rice, keep one row free after every 3 rows for good penetration of sunlight so that disease can be prevented. In areas prone to water accumulation, prepare broad bed and furrow systems only after excess water drains to improve drainage and aeration for groundnut. Harvest matured crops promptly when conditions permit and store in a dry, safe place. Remove and destroy infected fruits and leaves of chilli plant promptly to limit disease spread.

Andaman & Nicobar Islands

- In paddy, maintain proper drainage to remove excess water and maintain a water depth of 2–3 cm in the field. Avoid fertilizer and plant protection applications immediately before or during rainfall. Keep harvested produce and livestock feed in clean, dry and well-ventilated places. In ginger and turmeric fields where the soil has been disturbed by previous week rainfall, carry out earthing up to restore the plant base and cover exposed roots.

WEST INDIA [GOA, MAHARASHTRA, GUJARAT]

Realised rainfall: Significant rainfall occurred in all the districts of Konkan & Goa; in Kolhapur, Nasik, Pune districts of Madhya Maharashtra; in Bhandara and Gondia districts of Vidarbha; in Dadra and Nagar Haveli, Daman, Dangs, Mahisagar, Navsari, Sabarkantha, Tapi and Valsad districts of Gujarat. Weather remained mainly dry over rest of the region.

Advisories:

Maharashtra

- In Konkan, in view of light to moderate rainfall during next 5 days, undertake earthing up in turmeric. maintain 3 to 5 cm water level in transplanted rice fields. Undertake weeding operations in one month old transplanted finger millet / proso millet followed by application of second split dose of nitrogen fertilizer @ about 870 g urea per guntha. Roll an empty drum over 50–55 days old groundnut fields. Undertake application of paclobutrazol in mango orchards. High humidity conditions favour the incidence of bud rot disease in coconut, to control spray either 1% Bordeaux mixture or Copper Oxychloride @ 0.25% (25 g per 10 litre of water).
- In Madhya Maharashtra, In view of light to moderate rainfall during next 5 days, maintain 3-5 cm water level in rice fields. Continue intercultural operations in maize, cotton, green gram and black gram. If soybean crop turns yellow, spray a solution of 2% urea or 10 g of 19:19:19 fertilizer per litre of water during clear weather. Due to cloudy weather incidence of sap-sucking pests may observe in soybean, to control spray a 5% neem extract during clear weather. For control of sap-sucking pests in green gram and black gram, spray 5% Neem extract during clear weather and use 8 to 10 yellow sticky traps per acre. For effective control of sap-sucking pests (aphids, jassids, and thrips) in okra spray Imidacloprid 70% WG (1 g) or Thiamethoxam 25 WG (2 g) per 10 litres of water during clear weather. In view of light rainfall during next 5 days, continue intercultural operations in cotton, pigeon pea, green gram, black gram and pearl millet. Apply need based irrigation to groundnut, pigeon pea and soybean. Apply second dose of top dressing to 60 days old cotton crop 50 kg N/ha to BT or 40 kg N /ha respectively. If water stress is observed in green gram and black gram during flowering to pod-filling stage, apply light irrigation or spray 2% urea.

  In view of light to moderate rainfall during next 5 days, maintain 3-5 cm water level in rice (tillering stage). If water stress observed in finger millet, spray 19:19:19 water-soluble fertilizer at a 2% concentration. Spray Phule Micronutrient Grade II at 5–7 ml per litre of water in maize. In transplanted rice, Apply fertilizer dose of Urea @ 500 g per guntha in rice fields. Spray Phule Micronutrient Grade-II @ 40 ml per pump. Due to high humidity and frequent rain there may be incidence of Blight and Leaf Blight disease in rice thus to control spray the crop with Hexaconazole 5% E.C. @ 20 ml per 10 litres of water.

  In view of light to moderate rainfall during next 5 days, maintain 1-2 cm water level during seed

establishment stage and 2-3 cm during initial growth stage in rice. Undertake gap filling in groundnut and finger millet. Continue transplanting of kharif onion. Continue intercultural operations in groundnut and pearl millet.

- In Marathwada, in view of light rainfall during next 5 days, Apply need based application to sugarcane, cotton, maize, soybean, sorghum, pearl millet, orchards and vegetables. continue intercultural operations in *kharif* crops like cotton, maize, soybean, sorghum, pearl millet and sapota. if water stress is observed in sorghum, spray 1% (100 g/10 litres of water) Potassium nitrate (13:00:45). Undertake nipping operation in 45 days old pigeon pea crop. To control oily spot disease in Mrig Bahar pomegranate, spray Streptocycline @ 2 g or Copper Oxychloride @ 25 g per 10 litres of water. If infestation of leaf eating cater pillar & stem borer observed in soybean, spray Profenofos 50% @ 400 ml or Ethion 50% @ 600 ml or Thimethoxam 12.6% + Lambda Cyhalothrin 9.5% @ 50 ml or if incidence is more take a spray of Chlorantraniliprole 18.5% @ 60 ml per acre.
- In Vidarbha, apply need based application to soybean, cotton, pigeon pea, sorghum, maize, green gram and black gram. Continue intercultural operations (weeding / hoeing) in kharif crops like soybean, cotton, pigeon pea, sorghum and maize. In green gram and black gram, avoid intercultural operation during flowering stage. To control early stage incidence of aphids / jassids in cotton apply 5% neem extract or Azadirachtin 0.03% (300 ppm) @ 50 ml per 10 litres of water. To control early stage incidence of stem fly in soybean remove and destroy damaged parts of plant, for above ETL level (about 10% damaged plants), spray Chlorantraniliprole 18.5% @ 3 ml per 10 litres of water.

  In view of light to moderate rainfall during next 5 days, maintain optimum water level of 3 to 5 cm in transplanted rice field. Undertake weeding operation in 30 days old transplanted rice followed by application of 2nd dose of Nitrogen 25 kg/ha (54 Kg Urea per ha). Due to cloudy weather condition incidence of fruit borer is observed in okra, to control spray Quinolphos 25 EC @ 16 ml per 10 litres of water. Due to cloudy weather condition, sucking pest incidence is observed in brinjal and tomato, for control foliar spray of Dimethoate 30 % SC @ 20 ml per 10 litres of water.

Gujarat

- In Middle Gujarat Zone, carry out planting of banana; select tissue culture plant of varieties like Grand Naine, Rubusta, Basrai and Anand Vaaman. To control bacterial leaf blight disease in kharif rice, spray mixture of Streptocycline @ 1g + Copper oxychloride @ 20g fungicides in 400-500 litres of water per hectare. To control sucking pest in cotton, spray neem seed extract @ 500g or Neem oil @ 50ml in 10 litres of water.
- In Bhal and Coastal Zone, undertake interculturing and weeding operations in kharif crops. To control white grubs in groundnut, spray Chlorpyrifos 20 EC with 4 litres per hectare through drip irrigation or along with irrigation. To control aphids, whiteflies, leafhoppers and thrips in green gram and black gram, spray neem seed kernel powder (5% extract) @ 500 g or fungal biocontrol agent Verticillium lecanii powder @ 40g in 10 litres of water.
- In North West Zone, carry out sowing of cow pea crops. To control caterpillar in soybean, spray Chlorpyriphos 20% EC @ 25ml or Methomyl 40% SP @ 10 ml or Triazophos 40% EC @ 16ml in 10 litres of water. To control of aphids, leaf hopper and mealy bug in cotton, spray Imidacloprid 17.8% SL @ 3-4 ml in 10 litres of water. Carry out transplanting of vegetable crops.
- In South Saurashtra Zone, interculturing and hand weeding should be carried out in cotton; to control Stem borer infestation apply Carbofuran 3G @ 30 kg/ha and earthing up to support crop. To control white fly and leaf minor in vegetables, spray neem seed extract 5% or 40ml neem oil with Thiamethoxam 25 % WG @ 5g or Imidacloprid 30.5 % SC @ 5ml in 10 litres of water.
- In South Gujarat Zone, carry out sowing of pigeon pea and castor. Carry out planting of banana. Maintain 2–3 cm standing water in rice field; apply 40 kg and 48 kg of Nitrogen per hectare. To control sucking pests, pyrilla and whitefly in sugarcane, spray Thiamethoxam 25% WG @ 4g or Imidacloprid 17.8% SL @ 3ml in 10 litres of water.
- In North Saurashtra Zone, carry out inter culturing and weeding in black gram, green gram and cow pea. To

control leaf spot infestation in groundnut, spray Tebuconazole @ 10 ml in 10 litres of water. Incidence of jassids, aphids and white fly is observed in cotton, spray Acephate 75 % SP @ 15g with 1500 ppm neem oil 40 ml in 10 litres of water.

CENTRAL INDIA [M.P, CHHATTISGARH]

Realised Rainfall: Significant rainfall occurred in all districts of East Madhya Pradesh except Chhindwara and Seoni; in Asoknagar, Bhind, Bhopal, Datia, Guna, Gwalior, Morena, Narmadapuram, Neemuch, Rajgarh, Sehore, Shajapur, Sheopur Kalani, Shivpuri and Vidisha districts of West Madhya Pradesh; in all districts of Chhattisgarh except Kondagaon, Mungeli and Narayanpur.

Advisories:

- As **rain/thundershowers at many/most places** likely over the region during next 4-5 days and **Heavy to Very Heavy Rainfall** very likely at isolated places over East Madhya Pradesh during next 48 hours; **Heavy Rainfall** very likely at isolated places over West Madhya Pradesh and Chhattisgarh during next 5 days, over East Madhya Pradesh during 23rd to 25th August, 2026; postpone irrigation, intercultural operations and application of plant protection measures & fertilizers to the standing crops. Provide adequate drainage in standing crop fields to avoid waterlogging.

Madhya Pradesh

- In Malwa Plateau Zone, drain out excess water from maize, soybean. To manage stem rot in maize, ensure proper field drainage and avoid excessive application of nitrogen fertilizer.
- In Central Narmada Valley, drain out excess water from sugarcane, soybean, maize and vegetables. Undertake application of nitrogen fertilizers in maize, rice, and vegetables after the rain recedes.
- In Vindhya Plateau Zone, provide extensive drainage in soybean and vegetables. Undertake intercultural operation to remove weeds in kharif crops when soil is in workable conditions.
- In Bundelkhand Zone, provide extensive drainage in vegetable, pulse and oilseed crops. Weather is conducive for occurrence of rhizome rot disease in colocasia and ginger crops; monitor the crops for the same and if noticed undertake appropriate plant protection measures.
- In Jhabua Hill Zone, in 30–40-day-old maize, carry out earthing-up by placing soil around the plant base to prevent lodging. For effective management of caterpillars in black gram during flowering stage, at the early stage of infestation, install 20–25 'T'-shaped bird perches (2–2.5 feet high) and 8 pheromone traps per hectare. Strong winds may cause young and tender vegetable plants to bend or lodge. To prevent this, carry out earthing-up around the base of the plants and provide proper staking with bamboo sticks or wooden stakes to keep the plants upright and protect them from lodging due to strong winds.
- In the Satpura Plateau Zone, ensure proper drainage in maize, cotton, sugarcane and vegetable crops. Provide staking / physical support to banana, tomato & chillies. In some areas due to excess moisture in maize crop, stem rot disease has been reported, for its prevention do not use nitrogen fertilizer in large amounts. Spraying with copper oxychloride (COC) @ 3g per litre of water at an interval of 12-15 days is recommended.

Chhattisgarh

- In the Bastar Plateau Zone, ensure proper drainage from rice, maize, groundnut, minor millets and pulses fields. Undertake sowing of green gram, black gram & horse gram at optimum soil moisture condition by selecting improved varieties after seed treatment. Carry out weeding in *kodo, kutki, ragi,* groundnut & red gram. Carry out earthing up in timely sown Kharif maize at vegetative growth, install ten T-shaped bamboo perches per acre in the field so that birds can sit on them and easily prey on caterpillars. Maintain proper water level in rice field; spray Trifloxystrobin 25% +Tebuconazole 50% WG or Azoxystrobin @ 1g per litre of water during clear weather at 12–15 days of intervals as soon as the initial symptoms of blast appear.
- In the North Hill Zone, ensure proper drainage to avoid water stagnation in rice, maize, black gram, pigeon pea, cucurbitaceous crop, tomato, capsicum & okra. In black gram, undertake weeding only when soil conditions permit. In cucurbitaceous crops, maintain drainage and provide ladder-like bamboo support. In tomato and capsicum, harvest ripe fruits promptly, maintain proper staking and support plants during flowering and fruit formation. In papaya and banana, provide support to fruit-bearing plants.

- In the Chhattisgarh Plain Zone, drain out the excess water from red gram, green gram, black gram & groundnut. Undertake planting of papaya, turmeric & ginger; maintains the drainage channel in the field. If symptoms of powdery mildew are seen in green gram & black gram, spray soluble Sulphur @ 3g/ litre of water during clear weather. In soybean, monitor leaf-eating caterpillars, white fly and stem fly, where necessary, apply Beta Cyfluthrin + Imidacloprid @ 350ml/ha or Thiamethoxam + Lambda cyhalothrin @ 125 ml/ha after rainfall subsides.

Animal Husbandry

- In Chhattisgarh, keep livestock and poultry in safe, dry shelters. Maintain adequate ventilation in animal sheds. Keep animals away from waterlogged areas. Provide a balanced diet to animals Ensure vaccination against anthrax and foot-and-mouth disease.

NORTH INDIA [JAMMU & KASHMIR (UTs), HIMACHAL PRADESH, UTTARAKHAND, PUNJAB, HARYANA, DELHI, UTTAR PRADESH & RAJASTHAN]

Realised Rainfall: Significant rainfall occurred in Banswara, Baran, Bharatpur, Bhilwara, Chittorgarh, Dausa, Dholpur, Dungarpur, Jhalawar, Karauli, Pratapgarh & Udaipur districts of East Rajasthan; in all districts of East Uttar Pradesh except Chandauli, Deoria, Kanpur Dehat, Kushi nagar, Mau; in all districts of West UP except Aligarh, Baghpat, Bulandsahar, Firozabad, Ghaziabad, Hapur, Hathras, Mainpuri, Mathura & Shamli; in most of the districts of Himachal Pradesh except Kinnaur and Lahaul & Spiti; in Doda, Jammu, Kathua, Kulgam, Poonch, Reasi, Samba and Udhampur districts of Jammu & Kashmir and in all the districts in Uttarakhand; in Fatehgarh Saheb, Gurdaspur, Hoshiarpur, Pathankot, Patiala, Rupnagar, SBS Nagar districts of Punjab; in Ambala, Chandigarh, Kurukshetra, Panchkula, Panipat, Sonepat, Yamunanagar districts of Haryana and in North Delhi and West Delhi districts of New Delhi.

Advisories:

- As **rain/thundershowers at many/most places** likely over Himachal Pradesh during next 48 hours, over Jammu & Kashmir during next 72 hours, over and **Heavy Rainfall** very likely at isolated places over East Uttar Pradesh, Haryana-Chandigarh-Delhi, Himachal Pradesh, Punjab during next 48 hours, over Uttarakhand during next 5 days, over Jammu-Kashmir-Ladakh-Gilgit-Baltistan-Muzaffarabad on 22nd August, 2026;postpone irrigation, intercultural operations and application of plant protection measures & fertilizers to the standing crops. Provide adequate drainage in standing crop fields to avoid waterlogging.
- As **Thunderstorm accompanied with lightning & gusty winds(30-40kmph)** very likely at isolated places over West Uttar Pradesh during next 48 hours, over Jammu-Kashmir-Ladakh-Gilgit-Baltistan-Muzaffarabad and West Rajasthan during next 4 days; provide mechanical support to banana & papaya plantations and staking to young plants and vegetables. Undertake propping in sugarcane.

Himachal Pradesh

- In High Hill Temperate Dry Zone, keep the cucumber and tomato plants upright by providing support. Undertake sowing of buckwheat, oats and toria. Remove weeds from rajmash & potato and apply irrigation as per requirement. Remove diseased plants in barley field and destroy them. Apply irrigation as per requirement. For effective management of aphids in cauliflower and cabbage, spray Imidacloprid 17.8% SL @ 0.5 ml per litre of water or Thiamethoxam 25% WG @ 0.3g per litre of water. Apply the spray during clear weather and ensure thorough coverage of the affected plant parts.
- In Mid Hills Sub Humid Zone, provide support to cucumber vines. Protect the cucumber, tomato and brinjal fruits from making direct contact with soil. Undertake timely harvesting of mature brinjal, capsicum and tomato fruits. Ensure drainage channels are clean in ginger and turmeric. Do not allow rain water to stand in the maize crop as it is highly sensitive to standing water and promotes bacterial stalk rot.
- In Sub Montane and Low Hills Sub Tropical Zone, maintain 2-5 cm standing water in transplanted rice field. Ensure quick drainage from maize fields. Provide staking/support to tomato and other climbing vegetables and keep the crop canopy well ventilated. Avoid plucking tea leaves during or just after rain to reduce fungal contamination in processed leaves. Ensure proper surface drainage in tea fields.
- In High Hills Sub Temperate Wet Zone, harvest cucumber, bottle gourd, bitter gourd, snake gourd, ridge gourd,

and sponge gourd regularly (every 2-3 days, depending on the crop and season) to encourage continuous flowering and higher yields. Maize is sensitive to oversupply of water. Therefore, make proper drainage channels to remove the extra water from the fields. Carry out hoeing and weeding in finger millet during clear weather.

U.T. of Jammu & Kashmir

- In Sub Tropical Zone, carry out intercultural operations like hoeing, weeding and earthing up in maize during clear weather. To control stalk rot in maize, spray Mancozeb @ 0.25% during clear weather. To control stem borer, apply Cartap Hydrochloride/Carbofuran granules @ 5g per plant during clear weather.
- In Intermediate Zone, provide mechanical support to young vegetable plants to prevent lodging (bending or falling over) and ensure a better crop stand against strong winds. To control leaf blight in maize, spray Propiconazole @ 0.1% during clear weather. To control blast in rice, spray Tricyclazole @ 0.06 % or 1 ml in 1.5 litres of water on a non-rainy day. To prevent water logging in pulse crop, ensure proper drainage by creating channels and drains to remove excess water.
- In Valley Temperate Zone, after the semi-dough stage of rice, follow alternate wetting and drying up to physiological maturity to optimize water use. Regularly monitor rice fields and bunds for rodent activity and destroy active burrows to minimize crop damage. Keep field bunds and surrounding areas free from weeds and grasses to reduce rodent shelter and breeding sites. In fields with severe rodent infestation, adopt community-based rodent management using recommended baiting methods. Undertake harvesting of early-sown maize wherever cobs have attained physiological maturity. Avoid moisture stress in all vegetable crops especially at flowering stage. Maintain weed free conditions in all vegetables.

U.T. of Ladakh

- In Cold Arid Zone, give light irrigation to wheat in soft dough stage, if the soil is dry, as water is still needed to fill the grain. Stop watering 10 to 15 days before harvest (in hard dough stage) so the plants dry evenly and are ready to cut. Pick cucumbers every 2 to 3 days in the morning. Do not let them turn yellow or oversized, which lowers the quality. Keep the soil evenly moist with regular watering. Dry soil makes cucumbers taste bitter. Look for white powdery spots on leaves (powdery mildew) caused by cool night air. Remove severely infected leaves immediately. Apply light irrigation a day before harvesting radish. This makes it easier to pull up the roots without breaking them and keeps the tubers fresh.

Uttarakhand

- In Bhabar and Tarai Zone, avoid irrigation in rice. Provide proper drainage in sugarcane. Tie the crop at 5 feet height to avoid damage to the tall crop. Maintain drainage and carry out hoeing and weeding in maize at optimum soil moisture condition. To control Bacterial blight infestation in rice, spray Copper Oxychloride @ 500g and Streptocycline @ 15g in 500 litres of water at an interval of 7-10 days during clear weather conditions.
- In Hill Zone, ensure rapid drainage of excess rainwater from the barnyard millet and kidney beans field to avoid prolonged water stagnation. Remove the mulch (if applied) in ginger, in order to prevent water stagnation and rotting of developing rhizomes.
- In Sub Humid and Sub Tropic Zone, drain out excess water from rice field and retain shallow standing water. Divert excess runoff in barnyard millet through stable terrace drains to protect emerging ears and prevent soil erosion. Clear terrace outlets in maize field and strengthen bunds to reduce erosion, waterlogging and lodging. Maintain terrace drainage in finger millets and protect terrace risers from runoff without cutting unstable slopes. Secure stakes and trellises in tomato, harvest ripe fruits safely and drain promptly. Inspect for fungal symptoms after drying; avoid applying fertilizer or pesticides during rainy weather.

Punjab

- In Western Zone, prevailing rainy and humid weather is highly favourable for the development of Phytopthora diseases in citrus orchards. Drench the soil around each tree with Ridomil Gold or Curzate M-8 @ 25 g per 10 litres of water. Spray 80 ml Coragen 18.5 SC or 80g Proclaim 5 SG or 100 ml Sumicidin 20 EC or 200 ml Ripcord 10 EC in 100-125 litres of water against fruit and shoot borer of brinjal. In brinjal,

spider mite attack can be minimized by spraying O-mite 57 EC @ 300 ml per acre in 150 litres of water. Sheath blight may be noticed during the month of August in paddy. The disease can be checked by spraying Amistar Top 325 SC @ 200 ml or Nativo 75 WG @ 80 g in 200 litres of water. Keep the bunds clean by removing weeds.

- In Western Plain Zone, maintain appropriate water management in transplanted rice. Apply need based irrigation to sugarcane crop. For top borer in sugarcane, apply Ferterra 0.4 GR @ 10 kg/acre or Chlorantraniliprole 18.5 SC @ 150 ml in 400 litres of water/acre only if damage exceeds 5% and during clear weather. Monitor guava for fruit fly and use PAU-recommended fruit-fly traps.
- In Undulating Plain Zone, undertake transplanting of cauliflower and sowing of carrot and radish where soil conditions permit. If borer damage appears on maize, spray Coragen 18.5 SC @ 30 ml/acre in 60 litres of water during a rain-free period. Carry out weeding in soybean and groundnut to save from moisture and nutrient stress. Whitefly spreads mungbean yellow mosaic virus in soyabean. At first appearance, spray 1 L homemade neem extract mixed with 80–100 L water per acre using a knapsack sprayer. Prepare the extract by boiling 5 kg neem leaves and fruits in 10 L water for 30 minutes, then filter through a muslin cloth. Repeat the spray after one week if needed. Regularly monitor the maize crop for signs of fall armyworm infestation.
- In Sub Montane Undulating Zone, planting of evergreen fruit plants such as citrus, mango, litchi, guava, loquat, ber, amla and papaya where field conditions permit. Maintain proper drainage in maize to prevent water stagnation. Maize borer can be controlled by spraying Chlorantraniliprole 18.5 SC @ 30ml in 60 litres of water per acre at 2-3 weeks old crop in both irrigated and rainfed crops. To control whitefly, jassids and aphids in green gram, spray the crop upon pest appearance with 1 litre of homemade neem extract using 80-100 litres of water per acre with manually operated knapsack sprayer.
- In Central Plain Zone, undertake planting of evergreen fruit plants such as citrus, mango, guava, litchi, loquat and sapota where soil conditions permit. Maintain proper drainage in sugarcane and remove excess rainwater promptly. Prevailing weather is highly favourable for the development of Phytopthora diseases in citrus orchards. Drench the soil around each tree with Ridomil Gold or Curzate M-8 @ 25g per 10 litres of water during this month. For the management of sugarcane pyrilla, use Chlorpyriphos 20EC @ 600ml /acre in 400 litre of water. To control banded leaf and sheath blight of maize, spray Amistar Top 325 SC @ 100 ml in 200 litres of water as soon as it appears in the field.

Haryana

- In Haryana, drain out the excess rain water from the cotton crop. Ensure proper drainage in standing crop to prevent waterlogging and root damage (due to high rainfall) & immediate remove excess rainwater from furrows from the field. High humidity level, favour insect infestation in cotton, keep vigil on white fly infestation. If infestation of white fly is observed above the economical threshold level (ETL), spray of 1 litre Nimbecidine mixed with 200-250 litres of water per acre, during clear weather. For control of leafhopper infestation, spray Imidacloprid 200SL @ 40 ml mixed with 120-150- liter water per acre, during clear weather. Due to high moisture and favourable conditions, keep vigil on root weevil in rice crop and if infestation occurs, apply 10 kg Carbaryl 4G/Sevidol 4G per acre, during clear weather. Maintain water level in transplanted paddy fields.

Delhi

- Undertake sowing of carrot, sweet corn and baby corn in raised bed. Transplant brinjal, tomato, chilli and early cauliflower on ridges where field conditions permit. Transplant onion seedlings in the prepared fields. Undertake sowing of sarson saag, radish, spinach, coriander. Farmers are advised to install pest specific pheromone traps @ 3-4/acre in the paddy fields for monitoring of stem borer. In vegetables (brinjal, cauliflower, cabbage, tomato etc), for monitoring of borer insect, install pheromone trap @ 3-4/acre.

Uttar Pradesh

- In the Central Zone, maintain proper drainage in paddy and other crop fields and postpone fertilizer, herbicide and pesticide application during rainy conditions. Continue nursery raising of brinjal, chilli and early cauliflower and sowing of cucurbits and okra only where field conditions permit. Continue nursery sowing of

recommended paddy varieties in areas where transplanting has been delayed or additional nursery is required, subject to suitable field conditions; if Khaira disease symptoms appear, apply zinc sulphate with the recommended dose of urea only during a rain-free period. Stem borer and shoot fly may attack the maize crop, monitor regularly; if infestation is noticed, spray Emamectin Benzoate 5% SG @ 200g /ha mixed in 500–600 litres of water during clear weather. To control weeds in black gram and green gram, spray Imazethapyr 10% SL @ 1 litre/ha mixed in 500–600 litres of water, about 15–20 days after sowing when the weather is clear.

- In the Eastern Plain Zone, continue sowing of recommended early varieties of pigeon pea and raising additional rice nurseries or direct sowing of short-duration (110–125 days) improved paddy varieties only where the soil is suitably moist but not waterlogged. In autumn-planted sugarcane, provide support to prevent lodging; undertake earthing-up after weeding only when field conditions permit. Continue transplanting of brinjal, tomato, chilli and early cauliflower. Sow aromatic rice varieties during August so that the grain develops good quality and aroma. If Khaira disease is seen in paddy, spray a solution of 5 kg zinc sulphate + 2.5 kg slaked lime in 1000 litres of water per hectare. Urea may also be used as recommended. The present weather is suitable for sowing okra (ladyfinger) and cowpea; complete sowing at the earliest. Plant mango, guava, jackfruit and lemon seedlings in raised beds or polythene bags for better growth and drainage.
- In the Western Plain Zone, ensure proper drainage in pigeon pea and prevent waterlogging in pulse, oilseed, turmeric, ginger fields and vegetable crops. Uproot and destroy sugarcane plants infected with Red Rot Disease, including their roots. After removal, apply 0.2% Thiophanate Methyl solution as a soil drench. Farmers who have kept their fields vacant for autumn sugarcane planting, should sow Dhaincha (Sesbania) and incorporate it into the soil after 50–55 days as green manure. Due to high humidity and rainfall, rice may be affected by Bacterial Leaf Blight (BLB); for control, spray Streptocycline Sulphate 90% + Tetracycline Hydrochloride 10% @ 15 g with Copper Oxychloride 50% WP @ 500g in 500–750 litres of water per hectare during clear weather.
- In the Bundelkhand Zone, continue sowing of pearl millet (bajra), sesame and pigeon pea only where adequate but non-saturated soil moisture is available. Continue raising paddy nurseries where transplanting has been delayed. In sesame, postpone application of 60–65 kg urea and 25 kg sulphur/ha until rainfall subsides and soil conditions become suitable. Undertake weed control in pigeon pea, green gram and black gram only when the soil is workable. Continue transplanting of tomato, brinjal, chilli, early cauliflower, cabbage and marigold and establishment of new orchards of mango, guava, litchi, aonla, jackfruit, lemon, jamun, ber and papaya should be taken up only after excess water drains and suitable field conditions prevail.

Rajasthan

- In Hyper Arid Partially Irrigated Western Plain Zone, undertake hoeing and weeding in pearl millet. Manage weeds below threshold level at early growth stage in cluster bean, sesame and moth bean and apply Imidacloprid @ 0.3ml/l or Thiamethoxam @ 0.3 g/l or Dimethoate @ 1.5 ml/l of water for control of thrips. Irrigate horticultural fruit crops such as date palm, kinnow, orange and lemon as per crop requirement, considering the prevailing soil moisture conditions.
- In Flood Prone Eastern Plain Zone, to control downy mildew and blast disease due to high humidity spray Mancozeb (0.2%) or Metalaxyl in bajra crop.
- In Southern Humid Plain Zone, after the weather clears, apply Tembotrione 42 SC @ 120g a.i./ha in 500 L water at 25 DAS for weed control in maize. Maintain plant population in cotton; keep soybean and cotton field free from weeds. Undertake transplanting of paddy in main fields.
- In Transitional Plain Zone of Inland Drainage, spray Carbendazim @ 0.5g /L of water in clear weather to prevent and manage leaf spot disease in groundnut. Postpone irrigation in orchards and utilize the expected rainfall to meet plant water requirements. Undertake transplanting of early cauliflower in well-prepared fields, preferably when soil moisture is adequate. Harvest mature marigold and rose flowers during dry weather to maintain flower quality; ensure proper drainage and air circulation.
- In Semi-Arid Eastern Plain Zone, to protect pearl millet crop from white grub infestation, apply Imidacloprid

17.8 SL @ 500 ml per hectare mixed with 100 kg of soil, followed by light irrigation. Transplant papaya seedlings and before transplanting, apply 10 kg of well-rotted FYM and 300g of SSP per pit. Harvest marketable fruits of tomato, chilli, brinjal and okra regularly and maintain proper drainage in fields to avoid water logging.

- In Arid Western Plain zone, regularly monitor the mung bean crop against infestation of whiteflies or other sucking pests, spray Malathion 50 EC or Dimethoate 30 EC @ 1 litre/ha as a foliar spray only during morning or evening hours.
- In Southeastern Humid Plain Zone, ensure proper drainage in paddy, soybean, maize and vegetable fields to prevent waterlogging. Continuously monitor the rice crop for bacterial leaf blight, blast, and stem borer pests and avoid applying nitrogen during prolonged wet weather. Avoid irrigation during wet weather and postpone spraying operations until the weather clears up.
- In Irrigated North Western Plain Zone, for control of early stem borer and root borer in sugarcane crop, use Furadan dust 3% @ 6 kg per bigha or Phorate 10% @ 4 kg per bigha. Keep the cotton field free from weeds and apply weeding after first irrigation and maintain plant population.
- In Sub-Humid Southern Plain and Aravali Hill Zone, control incidence of Fall Army Worm and whitefly infestations at egg stage by release of Trichogramma @ 1.5 lack/ha in maize crop. Excessive moisture and continuous rainfall can increase the incidence of blight and leaf spot diseases in tomato; prevent waterlogging in the field, and remove and destroy diseased leaves.

Animal Husbandry

- In Rajasthan, during the rainy season, keep animals in a clean, dry, and well-ventilated area. Do not allow water to stagnate around the shed and provide them with clean drinking water. Protect the animals from cold, rain and monitor them for foot problems and infections. Keep their fodder dry and clean

The Agromet Products in this Bulletin are generated by Agrimet Division, weather forecast issued by NWFC, New Delhi and Agromet Advisories are prepared based on the DAAS Bulletins issued by different Agro-meteorological Field Units (AMFUs).