

Basic Information

Age Range
21-30 years

Sex
Male

Weight
65 kg

Height
-- cm

Procedure Information

Type of Procedure
Emergency

Surgical Branch
Vascular surgery, Orthopedic surgery

Procedure
Popliteal bypass - Femur/ Tibia Nailing / Plating

ASA Grade
E, II

Incident Narrative Description

Planned for GA with ETT
After induction - laryngoscopy revealed CL IIb with BURP - bougie inserted - ETT size 8.0mm could not be inserted - attempted with size 7.5 mm - again could not be inserted. BMV initiated - senior called. Re-attempted with C-Mac - again bougie used for intubation - size 8.0 mm inserted.
No desaturation - no hemodynamic instability.

Anesthesia Technique

Primary
> General Anesthesia
> ETT

Monitoring Used

ECG

Capnograph

Pulse Oximeter

IBP

AGM

CVP

Urine output

Incident Demographics

Time of Incident

> Night hours

Place of Incident

> Operating room

Timing of Event

> On induction

Incident Relation

> To Anesthesia

Airway Incident

Intubation Problems

> Difficult Intubation

Patient Outcome Category

Error, No Harm

> Category D - D2 - Required intervention to preclude harm

Final Notes

Attributed to inexperience of first operator.

Reviewer's Comments

Review #1

Incident & Outcome

Difficult Intubation

Contributing Factors

none

Root Cause & Prevention

No mention of airway assessment

Suggested Course of Action

Well done

Review #2

Incident & Outcome

Unanticipated difficult intubation, with successful intubation being done in third attempt following change of hands

Contributing Factors

Operator inexperience, inadequate training , unanticipated difficulty requiring expert intervention

Root Cause & Prevention

-Robust preanesthetic airway assessment
-being prepared for unanticipated difficulties
- Call for help early