

Multivariate statistical dependence measure based on characteristic functions

povilas.daniusis, povilasd@neurotechnology.com

November 2021

Abstract

In this paper we propose multivariate statistical dependence measure based on difference between joint and marginal characteristic functions. We discuss simulated examples, and applications for feature selection/extraction, causal inference and conduct corresponding experiments with diverse collection of multivariate data sets.

1 Introduction

2 Previous work

3 Proposed Method

We will derive independence measure relying on property of characteristic functions (also known as Kac theorem [1]), that independence of two random vectors $X \in R^{d_x}$ and $Y \in R^{d_y}$ is equivalent to $\forall \alpha \in R^{d_x}, \beta \in R^{d_y}$,

$$\mathbb{E}_{X,Y} e^{i\langle \alpha, X \rangle + i\langle \beta, Y \rangle} = \mathbb{E}_X e^{i\langle \alpha, X \rangle} \mathbb{E}_Y e^{i\langle \beta, Y \rangle}, \quad (1)$$

where $i = \sqrt{-1}$, d_x and d_y are dimensions of X and Y , respectively.

This motivates the construction of a novel statistical dependence measure, which we further refer to as Kac independence measure (KacIM):

$$\kappa(X, Y) = \max_{\alpha, \beta} |\mathbb{E}_{X,Y} e^{i\langle \alpha, X \rangle + i\langle \beta, Y \rangle} - \mathbb{E}_X e^{i\langle \alpha, X \rangle} \mathbb{E}_Y e^{i\langle \beta, Y \rangle}| \quad (2)$$

It is easy to see that $0 \leq \kappa(X, Y) \leq 1$, $\kappa(X, Y) = \kappa(Y, X)$.

3.1 Estimation

Having i.i.d. data (x_j, y_j) , $j = 1, 2, \dots, n$ an empirical estimator of KacIM (2) is defined via corresponding empirical characteristic functions:

$$\hat{\kappa}(X, Y) = \max_{\|\alpha\|=\|\beta\|=1} \left| \frac{1}{n} \sum_{j=1}^n e^{i(\langle \alpha, x_j \rangle + \langle \beta, y_j \rangle)} - \frac{1}{n^2} \sum_{j=1}^n e^{i\langle \alpha, x_j \rangle} \sum_{k=1}^n e^{i\langle \beta, y_k \rangle} \right|. \quad (3)$$




Figure 1: Dependence detection in additive (left) and multiplicative (right) noise scenario.

Empirical estimator also admits $0 \leq \hat{\kappa}(X, Y) \leq 1$. Normalisation of parameters α and β on to unit sphere is included due to stability issues. The estimator (3) can be calculated by using gradient ascend.

4 Experiments

Dependence measures have board area of applications. For example, regularization [?, ?], feature selection and extraction [?], information bottleneck methods [?], causal inference [?], among others. Further we will reformulate some key ideas in these topics for KacIM, and experimentally investigate corresponding empirical scenarios.

4.1 Generated data

We begin with simple example, which demonstrates the efficiency of KacIM for simulated multivariate data with additive and multiplicative noise.

In Figure 4.1 reflects KacIM values during iterative adaptation (500 iterations). In the case of independent data, both x_i and y_i ($d_x = 1024$, $d_y = 32$) are sampled from gaussian distribution, independently. In the case of dependent data, an additive noise (left graph) and multiplicative noise (right graph), the dependent variable is generated according to $y_i = \sin(Px_i) + \cos(Px_i) + \lambda\epsilon_i$ ($\lambda = 0.15$) and $y_i = (\sin(Px_i) + \cos(Px_i))\epsilon_i$, respectively, where P is $d_x \times d_y$ random projection matrix, $\epsilon_i \sim N(0, 1)$ and $\epsilon_i \perp x_i$.

When data is independent (blue graph), both in additive and multiplicative cases, due to independence, estimator (3) is resistant to maximization, and oscillates near zero. On the other hand, when data is not independent (orange graph), the condition of Kac theorem is violated and maximization of estimator (3) is possible.




Figure 2: Noise level (x axis) vs final iteration KacIM value (y axis). KacIM values for larger noise levels saturates to the level similar to its final values.

4.2 Influence of noise level λ to KacIM estimator value

In this simulation we use the same additive noise setting as in previous paragraph, but evaluate all noise levels $\lambda \in [0.0, 2.4]$, with step 0.1. Figure 4.2 empirically shows that value of KacIM correlates with noise level, and therefore the proposed measure is able not only to detect whether independence is present, but also to quantitatively evaluate it, which enables to use KacIM to derive cost functions for various other learning-based algorithms.

4.3 Feature Extraction

We conduct linear feature extraction by seeking

$$W^* = \arg \max_W \kappa(Wx, y). \quad (4)$$

Afterwards, feature extraction is conducted by $f = W^*x$ and k -nearest neighbor classification with Euclidean distance is performed, comparing unmodified inputs x and features of all possible dimensions up to d_x .

5 Discussion

References

- [1] David Applebaum, B.V. Rajarama Bhat, Johan Kustermans, J. Martin Lindsay, Michael Schuermann, Uwe Franz: Quantum Independent Increment Processes I: From Classical Probability to Quantum Stochastic Calculus