Supplementary Information

A natural ionic liquid: low molecular mass compounds of aggregate glue droplets on spider orb webs

Yue Zhao [1,2,*], Takao Fuji [1], Masato Morita [3], Tetsuo Sakamoto [3]

[1] *Laser Science Laboratory, Toyota Technological Institute, 2–12–1 Hisakata, Tempaku-ku, Nagoya 468–8511, Japan.*

[2] *Collaborative Open Research Center, Kogakuin University, 2665-1 Nakano, Hachioji, Tokyo, 192-0015 Japan*

[3] *Department of Applied Physics, School of Advanced Engineering, Kogakuin University, 2665-1 Nakano, Hachioji, Tokyo, 192-0015 Japan*

* Correspondence to Yue Zhao.

e-mail: zhaoyue@toyota-ti.ac.jp

List of Contents

Supplementary Figure S1

Mass spectrum of positive ions of the aggregate glue droplets attached to the substrate.

Supplementary Figure S2

Mass spectrum of negative ions of the aggregate glue droplets attached to the substrate.

Supplementary Figure S3

Secondary positive ion maps of the aggregate glue droplets attached to the substrate.

Supplementary Figure S4

Secondary negative ion maps of the aggregate glue droplets attached to the substrate.

Supplementary Figure S5

Topographic nanopatterns of the suspended and attached aggregate glue droplets.

Supplementary Figure S6

Corrected infrared transmission spectrum of choline dihydrogen phosphate.

Supplementary Figure S7

The photograph of the spider and orb web.

Supplementary Figure S8

The spectrum of excitation ultraviolet light.

Supplementary Figure S9

Optical setup of the second harmonic generation imaging system.



Supplementary Figure S1 Mass spectrum (m/z 1 to 300) of positive ions of the aggregate glue droplets attached to the substrate.