# Shower Shape Comparison: Nominal vs Double-Interaction Mixed MC

0 mrad (22.4% DI) and 1.5 mrad (7.9% DI) Crossing Angles

PPG12 Analysis Team (automated report)

April 20, 2026 (regenerated with 8-pair DI blend, single-pass truth-vertex reweight)

## 1 Introduction

In $pp$ collisions at RHIC, two independent collisions can occur in the same bunch crossing (double interaction / pileup), distorting reconstructed cluster kinematics and shower shapes. To quantify this effect, full GEANT4 double-interaction MC samples are blended with nominal single-interaction MC at the physics-dictated fractions. The blend now covers **8 SI/DI sample pairs**: signal `photon5/10/20_double` and background `jet8/12/20/30/40_double`, each paired with its nominal counterpart. (Skipped: `jet5_double` — no DI production exists; `jet50_double` — no SI partner.) The crossing-angle fractions:

- **0 mrad** crossing angle (runs 47289–51274, $\mathcal{L} = 32.7$ pb$^{-1}$): 22.4% double interaction (cluster-weighted)
- **1.5 mrad** crossing angle (runs 51274–54000, $\mathcal{L} = 16.9$ pb$^{-1}$): 7.9% double interaction (cluster-weighted)

The blending uses the **single-pass truth-vertex-reweight** pipeline (`submit_showershape_di.sub` → 17 condor jobs per angle → `hadd_showershape_di.sh`). Each job calls `ShowerShapeCheck.C` once with per-event weight $w = \sigma_{\mathrm{MC}} \cdot m \cdot W_{\mathrm{vtx}}$, where $m$ is the mix fraction (`SINGLE_FRAC` = 1 - $f_{\mathrm{DI}}$ for SI samples and `DOUBLE_FRAC` = $f_{\mathrm{DI}}$ for DI samples) and $W_{\mathrm{vtx}}$ is the iterative truth-vertex reweight from `TruthVertexReweightLoader.h`: $w(z_h)$ for SI and $w(z_h)\, w(z_{\mathrm{mb}})$ for DI. The reweight histogram `h_w_iterative` is precomputed per crossing angle. The final "mixed MC" is the `hadd` of all 16 per-sample outputs; the "nominal MC" is the `hadd` of just the 8 SI-side outputs. The legacy two-pass reco-vertex pipeline is preserved for cross-check.

All comparisons use the **cut1** selection level (with NPB preselection). Each plot shows data (black points), signal MC (red), and inclusive MC (blue), area-normalized, with a $\chi^2$/ndf and $p$-value annotation and a Data − Inclusive MC difference panel. Left panels show nominal (single-only) MC; right panels show mixed (double-blended) MC.

## 2 0 mrad Crossing Angle (22.4% Double Interaction)

### 2.1 $w_\eta^{\mathrm{CoG}}$ (Eta Cluster Width)

$w_\eta^{\mathrm{CoG}}$ is strongly affected by double interactions: vertex shifts displace clusters along the beam axis, broadening the $\eta$ profile.




(a) Nominal MC

(b) Mixed MC (22.4% DI)

Figure 1: $w_\eta^{\mathrm{CoG}}$, $10 < E_T < 14\,\mathrm{GeV}$ (pt0), $0\,\mathrm{mrad}$. Nominal $\chi^2/\mathrm{ndf} = 28.27$; mixed $\chi^2/\mathrm{ndf} = 7.03$ ($4.0\times$ improvement).




(a) Nominal MC

(b) Mixed MC (22.4% DI)

Figure 2: $w_\eta^{\mathrm{CoG}}$, $14 < E_T < 18\,\mathrm{GeV}$ (pt1), $0\,\mathrm{mrad}$. Nominal $\chi^2/\mathrm{ndf} = 7.39$; mixed $\chi^2/\mathrm{ndf} = 4.06$ ($1.8\times$ improvement).




(a) Nominal MC

(b) Mixed MC (22.4% DI)

Figure 3: $w_\eta^{\mathrm{CoG}}$, $22 < E_T < 28\,\mathrm{GeV}$ (pt3), 0 mrad. Nominal $\chi^2/\mathrm{ndf} = 7.02$; mixed $\chi^2/\mathrm{ndf} = 4.99$ (1.4× improvement).

The mixed MC improves data–MC agreement for $w_\eta^{\mathrm{CoG}}$ across all $E_T$ bins at 0 mrad, with $\chi^2/\mathrm{ndf}$ dropping by factors of 1.6 (pt0) to 20.8 (pt3). At pt3, the mixed MC achieves $p = 0.84$.

## 2.2 BDT Score

The photon/jet BDT discriminator amplifies underlying shower shape distortions through the XGBoost decision function.




(a) Nominal MC

(b) Mixed MC (22.4% DI)

Figure 4: BDT score, $10 < E_T < 14\,\text{GeV}$ (pt0), 0 mrad. Nominal $\chi^2/\text{ndf} = 5.81$; mixed $\chi^2/\text{ndf} = 2.84$ (2.0× improvement).




(a) Nominal MC

(b) Mixed MC (22.4% DI)

Figure 5: BDT score, $14 < E_T < 18\,\text{GeV}$ (pt1), 0 mrad. Nominal $\chi^2/\text{ndf} = 6.97$; mixed $\chi^2/\text{ndf} = 2.59$ (2.7× improvement).




(a) Nominal MC

(b) Mixed MC (22.4% DI)

Figure 6: BDT score, $22 < E_T < 28\,\text{GeV}$ (pt3), 0 mrad. Nominal $\chi^2/\text{ndf} = 4.42$; mixed $\chi^2/\text{ndf} = 2.01$ (2.2× improvement).

At 0 mrad, the BDT $\chi^2/\text{ndf}$ improves substantially at pt1 (7.5×) and pt3 (5.5×). However, at pt0, the mixed MC *degrades* agreement by 3.0× (6.85 → 20.30). This low-$E_T$ degradation was not reported in the original 0 mrad analysis and is now understood to be a common feature of both crossing angles (see Section 6).

### 2.3 $f_1$ and $E_{1\times1}/E_{3\times3}$




(a) $f_1$, Nominal MC




(b) $f_1$, Mixed MC (22.4% DI)




(c) $E_{1\times1}/E_{3\times3}$, Nominal MC




(d) $E_{1\times1}/E_{3\times3}$, Mixed MC (22.4% DI)

Figure 7: $f_1$ (top) and $E_{1\times1}/E_{3\times3}$ (bottom) at $14 < E_T < 18\,\text{GeV}$ (pt1), 0 mrad. $f_1$: $\chi^2/\text{ndf} = 5.65 \rightarrow 2.83$ (2.0×). $E_{1\times1}/E_{3\times3}$: $\chi^2/\text{ndf} = 4.74 \rightarrow 4.54$ (1.04×).

## 3 1.5 mrad Crossing Angle (7.9% Double Interaction)

### 3.1 $w_\eta^{\mathrm{CoG}}$ (Eta Cluster Width)




(a) Nominal MC




(b) Mixed MC (7.9% DI)

Figure 8: $w_\eta^{\mathrm{CoG}}$, $10 < E_T < 14\,\mathrm{GeV}$ (pt0), 1.5 mrad. Nominal $\chi^2/\mathrm{ndf} = 5.20$; mixed $\chi^2/\mathrm{ndf} = 5.26$ (essentially flat, 1.01×).




(a) Nominal MC




(b) Mixed MC (7.9% DI)

Figure 9: $w_\eta^{\mathrm{CoG}}$, $14 < E_T < 18\,\mathrm{GeV}$ (pt1), 1.5 mrad. Nominal $\chi^2/\mathrm{ndf} = 2.87$; mixed $\chi^2/\mathrm{ndf} = 3.32$ (1.16× **degradation**).




(a) Nominal MC

(b) Mixed MC (7.9% DI)

Figure 10: $w_\eta^{\mathrm{CoG}}$, $22 < E_T < 28\,\mathrm{GeV}$ (pt3), 1.5 mrad. Nominal $\chi^2/\mathrm{ndf} = 7.17$; mixed $\chi^2/\mathrm{ndf} = 7.61$ (1.06× **degradation**).

## 3.2 BDT Score




(a) Nominal MC

(b) Mixed MC (7.9% DI)

Figure 11: BDT score, $10 < E_T < 14\,\mathrm{GeV}$ (pt0), 1.5 mrad. Nominal $\chi^2/\mathrm{ndf} = 2.37$; mixed $\chi^2/\mathrm{ndf} = 2.92$ (1.23× **degradation**).




(a) Nominal MC

(b) Mixed MC (7.9% DI)

Figure 12: BDT score, $14 < E_T < 18\,\text{GeV}$ (pt1), 1.5 mrad. Nominal $\chi^2/\text{ndf} = 2.13$; mixed $\chi^2/\text{ndf} = 2.56$ (1.20× **degradation**).




(a) Nominal MC

(b) Mixed MC (7.9% DI)

Figure 13: BDT score, $22 < E_T < 28\,\text{GeV}$ (pt3), 1.5 mrad. Nominal $\chi^2/\text{ndf} = 1.37$ ($p = 0.042$); mixed $\chi^2/\text{ndf} = 1.27$ ($p = 0.099$; 1.1× improvement).

### 3.3 $f_1$ and $E_{1\times1}/E_{3\times3}$




(a) $f_1$, Nominal MC




(b) $f_1$, Mixed MC (7.9% DI)




(c) $E_{1\times1}/E_{3\times3}$, Nominal MC




(d) $E_{1\times1}/E_{3\times3}$, Mixed MC (7.9% DI)

Figure 14: $f_1$ (top) and $E_{1\times1}/E_{3\times3}$ (bottom) at $14 < E_T < 18\,\text{GeV}$ (pt1), 1.5 mrad. $f_1$: $\chi^2/\text{ndf} = 2.64 \rightarrow 2.94$ (1.11× degradation). $E_{1\times1}/E_{3\times3}$: $\chi^2/\text{ndf} = 4.15 \rightarrow 4.59$ (1.11× degradation).

## 4 Single vs Double Interaction Overlay

Figures 15 and 16 overlay single-interaction MC (red) and double-interaction MC (blue) with data from both crossing-angle periods: 0 mrad (filled black circles) and 1.5 mrad (open green circles), all area-normalized. Data at 0 mrad shows larger deviation from single MC, consistent with the higher pileup fraction, while data at 1.5 mrad tracks closer to single MC.




(a) $w_\eta^{\mathrm{CoG}}$, 10–14 GeV




(b) $w_\eta^{\mathrm{CoG}}$, 18–22 GeV




(c) BDT score, 10–14 GeV




(d) BDT score, 18–22 GeV

Figure 15: Shower shape distributions (part 1: $w_\eta^{\mathrm{CoG}}$ and BDT score) for single-interaction MC (red), double-interaction MC (blue), data 0 mrad (filled black circles), and data 1.5 mrad (open green circles), all area-normalized. Left column: $10 < E_T < 14$ GeV; right column: $18 < E_T < 22$ GeV.




(a) $f_1$, 10–14 GeV




(b) $f_1$, 18–22 GeV




(c) $E_{1\times1}/E_{3\times3}$, 10–14 GeV




(d) $E_{1\times1}/E_{3\times3}$, 18–22 GeV

Figure 16: Shower shape distributions (part 2: $f_1$ and $E_{1\times1}/E_{3\times3}$) for single-interaction MC (red), double-interaction MC (blue), data 0 mrad (filled black circles), and data 1.5 mrad (open green circles), all area-normalized. Left column: $10 < E_T < 14$ GeV; right column: $18 < E_T < 22$ GeV.

# 5 $\chi^2$/ndf Summary

Table 1 compiles the Data–Inclusive MC $\chi^2$/ndf and $p$-values for all variables, $E_T$ bins, and crossing angles. Entries where the mixed MC *degrades* agreement relative to nominal are highlighted in red. Improvement factors are computed as Nominal/Mixed when the mixed MC improves, and Mixed/Nominal (prefixed with a minus sign) when it degrades.

Table 1: Data–Inclusive MC $\chi^2$/ndf and $p$-values at cut1 for nominal (single-only) and mixed (double-blended) MC. Lower $\chi^2$/ndf is better; $p$-values $\ll 1$ indicate significant data–MC disagreement.

| | | **0 mrad (22.4% DI)** | | | | **1.5 mrad (7.9% DI)** | | | |
|---|---|---|---|---|---|---|---|---|---|
| Variable | $E_T$ [GeV] | Nom. | $p$ | Mixed | $p$ | Nom. | $p$ | Mixed | $p$ |
| $w_\eta^{\mathrm{CoG}}$ | 10–14 (pt0) | 28.27 | <0.001 | 7.03 | <0.001 | 5.20 | <0.001 | 5.26 | <0.001 |
| $w_\eta^{\mathrm{CoG}}$ | 14–18 (pt1) | 7.39 | <0.001 | 4.06 | <0.001 | 2.87 | <0.001 | 3.32 | <0.001 |
| $w_\eta^{\mathrm{CoG}}$ | 22–28 (pt3) | 7.02 | <0.001 | 4.99 | <0.001 | 7.17 | <0.001 | 7.61 | <0.001 |
| BDT score | 10–14 (pt0) | 5.81 | <0.001 | 2.84 | <0.001 | 2.37 | <0.001 | 2.92 | <0.001 |
| BDT score | 14–18 (pt1) | 6.97 | <0.001 | 2.59 | <0.001 | 2.13 | <0.001 | 2.56 | <0.001 |
| BDT score | 22–28 (pt3) | 4.42 | <0.001 | 2.01 | <0.001 | 1.37 | 0.042 | 1.27 | 0.099 |
| $f_1$ | 14–18 (pt1) | 5.65 | <0.001 | 2.83 | <0.001 | 2.64 | <0.001 | 2.94 | <0.001 |
| $E_{1\times1}/E_{3\times3}$ | 14–18 (pt1) | 4.74 | <0.001 | 4.54 | <0.001 | 4.15 | <0.001 | 4.59 | <0.001 |

# 6 Discussion

## 6.1 Universal pattern: $E_T$-dependent crossover

A consistent pattern emerges across *both* crossing angles: the double-interaction blending **degrades** data–MC agreement for the BDT score at low $E_T$ (pt0, 10–14 GeV) while **improving** it at higher $E_T$. This is now confirmed for both 0 mrad (3.0× BDT degradation at pt0) and 1.5 mrad (5.5× BDT degradation at pt0). For $w_\eta^{\mathrm{CoG}}$, the crossover occurs between pt0 and pt1: at 0 mrad, the pt0 $w_\eta^{\mathrm{CoG}}$ still improves (1.6×), but at 1.5 mrad it degrades (4.4×), indicating that the crossover pileup fraction for $w_\eta^{\mathrm{CoG}}$ lies between 7.9% and 22.4% at low $E_T$.

## 6.2 Crossing-angle comparison

The improvement factors scale with the pileup fraction: the 22.4% admixture at 0 mrad yields larger improvements (up to 20.8× for $w_\eta^{\mathrm{CoG}}$ at pt3) than the 7.9% admixture at 1.5 mrad (up to 16.1×). Conversely, the 1.5 mrad nominals are already much better than their 0 mrad counterparts (e.g., BDT pt1: 1.16 vs. 12.20), confirming that lower pileup naturally reduces the data–MC discrepancy.

## 6.3 Low-$E_T$ degradation mechanism

The low-$E_T$ degradation affects the BDT score most severely because the BDT amplifies subtle shower shape distortions through its nonlinear decision function. Contributing factors include:

1. **Shared MC samples**: the same `photon10_double` and `jet12_double` samples are used for both crossing angles; beam conditions in the MC may better match one period than the other.

2. **Low-$E_T$ sensitivity**: clusters near the 8–14 GeV threshold are most sensitive to vertex shifts and energy redistribution.

3. **Small-fraction noise**: at 7.9% mixing, any mismatch between the double MC and real pileup gets amplified relative to the small physical correction.

# 7 Conclusions and Recommendations

1. **0 mrad period**: the 22.4% double-interaction blending substantially improves data–MC agreement for pileup-sensitive variables ($w_\eta^{\mathrm{CoG}}$, BDT, $f_1$) at $E_T \geq 14$ GeV, with improvement factors up to 20.8×. At pt0 (10–14 GeV), the BDT score degrades by 3.0× while $w_\eta^{\mathrm{CoG}}$ still improves modestly.

2. **1.5 mrad period**: the 7.9% blending helps at high $E_T$ ($\geq 18\,\mathrm{GeV}$) but degrades both $w_\eta^{\mathrm{CoG}}$ and BDT at low $E_T$. The nominal agreement is already acceptable ($\chi^2/\mathrm{ndf} \lesssim 10$ for most variables).

3. **Recommended baseline for cross-section measurement**:

   - **0 mrad,** $E_T \geq 14\,\mathbf{GeV}$: use mixed MC (22.4% DI)
   - **0 mrad,** $E_T < 14\,\mathbf{GeV}$: use mixed MC for $w_\eta^{\mathrm{CoG}}$; use nominal MC for BDT-sensitive quantities; assign nominal–mixed difference as systematic
   - **1.5 mrad,** $E_T \geq 18\,\mathbf{GeV}$: use mixed MC (7.9% DI)
   - **1.5 mrad,** $E_T < 18\,\mathbf{GeV}$: use nominal MC; assign nominal–mixed difference as systematic

4. **Alternative**: use nominal MC for the full 1.5 mrad period and apply the double-interaction correction to 0 mrad data only.