

Cluster E_T Response and Resolution for Tight + Isolated Prompt Photons

Double-interaction mixed MC, 0 mrad + 1.5 mrad + lumi-weighted combined

PPG12

May 12, 2026

Setup

What is plotted. Per-truth- E_T -bin distributions and summary curves of the cluster energy response

$$R = \frac{E_T^{\text{cluster}}}{E_T^{\gamma, \text{truth}}}$$

for the tight-photon-ID + reco-isolated cluster arm of the analysis, with the matched truth particle restricted to direct or fragmentation prompt photons ($p_{\text{pid}} = 22$, $\text{photonclass} < 3$, $E_T^{\text{iso, truth}} < 4 \text{ GeV}$ in cone $R = 0.3$). The reco-side selection matches the cross-section analysis nominal: parametric tight BDT $\text{bdt} > 0.86 - 0.02 E_T^{\text{cluster}}$, parametric topo-cluster isolation $E_T^{\text{iso, reco}} < 0.49 + 0.037 E_T^{\text{cluster}}$ at $R = 0.4$, $|\eta^\gamma| < 0.7$, plus the standard common-region cuts.

Samples. Double-interaction blended signal MC (`photon5`, `photon10`, `photon20`; `SI = _nom`, `DI = _double` samples) weighted by the cluster-weighted double-interaction fractions $f_2^{0 \text{ mrad}} = 0.224$ and $f_2^{1.5 \text{ mrad}} = 0.079$. Per-event weight carries $\sigma \cdot f_{\text{mix}} \cdot (L_{\text{period}}/L_{\text{target}}) \cdot w_{\text{truth vtx}}$, with $L_{\text{target}} = 64.37 \text{ pb}^{-1}$ (all-range, allz). The lumi-weighted combined response is the `hadd` of the two period TH2 histograms, which reproduces the all-range distribution at L_{target} exactly because $L_{0 \text{ mrad}} + L_{1.5 \text{ mrad}} = L_{\text{target}}$.

Binning. Truth photon E_T in the 13 analysis truth-pT bins from the nominal config (`analysis.pT_bins_truth`): 2-GeV bins from 8 to 28 GeV, then [28, 32), [32, 36), and [36, 45) GeV. Response axis binned at 0.005 per bin in [0, 2.0]. The fill site is intentionally placed *outside* the analysis pT-range gate (the `cluster_Et` between `pTmin` and `pTmax` cut applied to the downstream ABCD histograms) so the [8, 10) and [36, 45) truth bins reflect the true detector response rather than an artificial truncation by histogram-fill cuts.

Fit conventions. Gaussian fit performed in $[\mu_{\text{hist}} - 2 \text{ RMS}, \mu_{\text{hist}} + 2 \text{ RMS}]$ to capture the core response. A double-sided Crystal Ball (DSCB) overlay is included on the QC grid to expose the low-side punch-through tail; we quote Gaussian (μ, σ) as the primary metric and histogram (mean, RMS) on [0.3, 1.7] as a cross-check.

Per-bin response distributions




Figure 1: Cluster E_T response distributions for the tight + iso prompt-photon arm, 0 mrad (22.4% DI) blended MC. One panel per truth E_T bin. Black: histogram projection. Blue dashed: Gaussian fit on $[\mu - 2\sigma, \mu + 2\sigma]$. Green: double-sided Crystal-Ball fit.




Figure 2: As Fig. 1, for the 1.5 mrad (7.9% DI) blended MC.





Figure 3: As Fig. 1, for the lumi-weighted combined sample ($L = 64.37 \text{ pb}^{-1}$).


Response and resolution vs truth photon E_T




(a) Gaussian fit mean.

(b) Gaussian fit width.

Figure 4: Response (left) and resolution (right) vs truth photon E_T for the tight + iso prompt-photon arm. Markers: 0 mrad (red squares), 1.5 mrad (blue triangles), lumi-weighted combined (black circles).



(a) Histogram mean on $[0.3, 1.7]$.



(b) Histogram RMS on $[0.3, 1.7]$.

Figure 5: Cross-check using the histogram mean and RMS on a fixed window $[0.3, 1.7]$ (insensitive to fit-window choice).







Figure 6: Four-panel summary: Gaussian (μ, σ) and histogram (mean, RMS) overlaid for the three samples.

Numbers

Per-bin Gaussian μ/σ from the fit on $[\mu_{\text{hist}} - 2 \text{ RMS}, \mu_{\text{hist}} + 2 \text{ RMS}]$, and histogram (mean, RMS) on the fixed window $[0.3, 1.7]$. Lumi-weighted combined values for the analysis.

$E_T^{\gamma, \text{truth}}$ [GeV]	Gaussian core fit				Histogram on [0.3, 1.7]			
	$\mu_{0 \text{ mrad}}$	$\sigma_{0 \text{ mrad}}$	$\mu_{1.5 \text{ mrad}}$	$\sigma_{1.5 \text{ mrad}}$	μ_{all}	σ_{all}	$\langle \text{Hist} \rangle_{\text{all}}$	RMS_{all}
[8, 10)	1.005	0.079	1.005	0.077	1.005	0.078	1.006	0.083
[10, 12)	0.994	0.082	0.995	0.079	0.995	0.081	0.993	0.087
[12, 14)	0.993	0.081	0.995	0.078	0.994	0.080	0.992	0.085
[14, 16)	0.994	0.079	0.995	0.076	0.994	0.077	0.992	0.083
[16, 18)	0.995	0.076	0.996	0.074	0.995	0.076	0.993	0.080
[18, 20)	0.994	0.076	0.996	0.072	0.995	0.074	0.993	0.079
[20, 22)	0.995	0.073	0.994	0.070	0.994	0.072	0.992	0.077
[22, 24)	0.995	0.072	0.996	0.069	0.995	0.071	0.993	0.076
[24, 26)	0.994	0.071	0.995	0.067	0.995	0.069	0.992	0.074
[26, 28)	0.992	0.069	0.992	0.066	0.992	0.068	0.990	0.072
[28, 32)	0.987	0.068	0.987	0.065	0.987	0.067	0.984	0.070
[32, 36)	0.974	0.066	0.973	0.063	0.974	0.065	0.972	0.067
[36, 45)	0.953	0.066	0.952	0.063	0.953	0.065	0.951	0.067

Table 1: Cluster E_T response and resolution for the tight + iso prompt-photon arm. Statistical uncertainties on the Gauss fit are $\lesssim 10^{-3}$ for most bins (see `resp_summary_numbers.txt`).

Observations

- Across $E_T^{\gamma, \text{truth}} \in [10, 28]$ GeV the Gaussian response sits at $\mu = 0.992$ – 0.996 (within 1 % of unity), consistent with the energy scale set by the EMCal calibration after the analysis selection. The [8, 10) bin is unbiased at $\mu = 1.005$.
- Above $E_T^{\gamma, \text{truth}} = 28$ GeV the response trends downward — $\mu = 0.987$ in [28, 32), 0.974 in [32, 36), and 0.953 in [36, 45). This $\sim 5\%$ drop in the highest truth bin is consistent with the known EMCal high-energy non-linearity and the steeply-falling truth- E_T spectrum (the bin average is dominated by entries near the lower bin edge where the response itself is closer to 1.0, so the observed μ understates the response at $E_T = 45$ GeV).
- Resolution improves monotonically from $\sigma \approx 0.080$ at $E_T = 13$ GeV to $\sigma \approx 0.065$ at $E_T = 35$ GeV. The [8, 10) resolution sits at $\sigma = 0.078$, in line with the [10, 12) bin once the histogram-fill bias is removed.
- Per-period: 0 mrad is consistently ~ 3 – 5% *wider* than 1.5 mrad (σ ratio in mid-bins $\sim 0.073/0.069 \approx 1.06$), matching the physical expectation that the higher double-interaction fraction at 0 mrad (22.4 % vs 7.9 %) broadens the response via the shifted reconstructed vertex.
- The combined-period μ and σ track the lumi-weighted average $(L_0 \mu_0 + L_{1.5} \mu_{1.5})/L_{\text{target}}$ closely, as expected — the hadd takes care of this automatically because the per-event $L_{\text{period}}/L_{\text{target}}$ pre-scaling is baked in by `RecoEffCalculator_TTreeReader.C`.

Reviewer pass matrix

Artifact	physics	cosmetics	re-derivation	fix-validator
<code>RecoEffCalculator</code> edit	PASS	N/A	N/A	PASS
<code>plot_cluster_response.C</code>	N/A	PASS	N/A	PASS
μ, σ in 3 representative bins	N/A	N/A	PASS	N/A
Report plots	N/A	PASS	N/A	N/A

Open WARN items (non-blocking): minor text-marker overlap in the `resp_summary_4panel` top-left panel and in the single-panel mean plots near $E_T \approx 11$ GeV; the sPHENIX info block sits close to the lowest- E_T markers due to the upward bias in that bin.

Implementation notes

- **Fill site:** `efficiencytool/RecoEffCalculator_TTreeReader.C`. Inside the `tight+iso` arm of the `truth-photon-matched` block (line 2893), fills a new TH2 `h_pT_truth_reco_tightiso_{ieta}` with $(E_T^{\gamma, \text{truth}}, E_T^{\text{cluster}}/E_T^{\gamma, \text{truth}})$.
- **Configs:** `config_bdt_clusterresp_0rad.yaml` and `_1p5mrad.yaml` (clones of `config_bdt_nom_*.yaml` with a unique `var_type` so the new histogram fills into dedicated files and leaves the nominal production intact).
- **Driver:** `efficiencytool/submit_clusterresp.sub` — 12 condor jobs (6 photon SI/DI samples $\times$ 2 periods), via the per-job wrapper `run_clusterresp_job.sh`.
- **Hadd:** `efficiencytool/hadd_clusterresp.sh` — builds per-period combined signal MC, then the all-range lumi-weighted combination.
- **Plot:** `plotting/plot_cluster_response.C`.