ABCD Purity Non-Closure: Non-Tight BDT Lower Bound and Flat Threshold Partitions

Shuonli Li / analysis proxy
sPHENIX Collaboration — PPG12 Isolated Photon

April 19, 2026

1 Abstract

We scan the non-tight BDT lower bound (`analysis.non_tight.bdt_min`) from its nominal value of 0.02 to 0.05 and 0.10 and quantify the ABCD purity non-closure against the MC truth purity in each of the twelve reconstructed p_T bins. In the 14–28 GeV physics range, the mean absolute non-closure is 0.062–0.073 with no strong monotonic trend in the scan parameter, while the low-E_T bins ($p_T \leq 11$ GeV) show a larger ABCD bias of -0.19 to -0.25 (absolute). The observed spread across the three scan points in the core range (standard deviation of |NC| of 0.03–0.04) is consistent with the current `nt_bdt` systematic entry in the purity group. We additionally evaluate four flat (ET-independent) BDT partitions around the collaborator's primary request of tight $= [0.9, 1.0]$, non-tight $= [0.5, 0.9]$: this configuration gives a core-range mean |NC| of 0.094 (slightly worse than parametric), while tightening the tight cut to 0.95 degrades closure substantially (mean 0.191, nonphysical `gpurity_leak` > 1 at kinematic edges).

2 Method

ABCD regions. The ABCD method partitions clusters into four regions using the tight-BDT and isolation selections:

- A = tight $\cap$ isolated (signal region),
- B = tight $\cap$ non-isolated (background template in BDT),
- C = non-tight $\cap$ isolated (background control in BDT),
- D = non-tight $\cap$ non-isolated.

Assuming approximate factorization of the BDT and isolation distributions for background, $N_A^{\text{bkg}} = R \cdot N_B N_C / N_D$ with leakage corrections $c_{B,C,D}$ for residual signal contamination in B, C, D.

Non-tight BDT bounds. The non-tight BDT window is $[\texttt{nt_bdt_min}, \texttt{nt_bdt_max}]$. The upper bound is the parametric, E_T-dependent cut $\texttt{nt_bdt_max} = 0.80 - 0.015 \cdot E_T$ (fixed at the nominal values). The lower bound `nt_bdt_min` is the scan variable; we take the three pipeline-consistent points $\{0.02, 0.05, 0.10\}$.

Purity extraction. The ABCD purity is obtained by solving the cubic closure equation implemented in `CalculatePhotonYield.C`, which includes signal-leakage corrections. We report the signal-leak-corrected estimator `gpurity_leak` as the ABCD purity. The truth purity is the ratio of truth-matched photon clusters to all clusters in region A,

$$\mathcal{P}_A^{\text{truth}} = \frac{N_A^{\gamma,\text{truth}}}{N_A^{\text{all}}}. \tag{1}$$

The non-closure is defined as the absolute difference $\text{NC} \equiv \mathcal{P}_A^{\text{ABCD,leak}} - \mathcal{P}_A^{\text{truth}}$.

Important caveat on the closure test. The closure test as implemented in `CalculatePhotonYield` (with `isMC=true`) uses the merged *jet* MC (`jet5+jet8+jet12+jet20+jet30+jet40`, cross-section weighted) as the "data" histogram, while the leakage corrections c_B, c_C, c_D are drawn from the separate merged photon MC (`photon5+photon10+photon20`). This is therefore a *self-consistency test* of ABCD within jet MC (where "photons" are direct/fragmentation photons originating in dijet events), and not a closure test on a cross-section-weighted signal+background cocktail. The reader should interpret the truth purity in the plot accordingly: it is the direct+fragmentation photon fraction of tight+iso clusters in jet MC.

Inputs. The three scan points correspond to the ROOT files

`efficiencytool/results/Photon_final_bdt_{nom,ntbdtmin05,ntbdtmin10}_mc.root`.

3 Parametric Non-Tight BDT Lower-Bound Scan



(a) Truth-matched signal purity. The three `nt_bdt_min` points share region A by construction, so the curves overlap.

(b) ABCD / truth purity ratio. Deviations from unity are the ABCD non-closure.

Figure 1: Parametric scan over the non-tight BDT lower bound `nt_bdt_min` $\in \{0.02, 0.05, 0.10\}$. Truth-matched signal purity (left) and ABCD/truth ratio (right) vs reconstructed E_T. Absolute non-closure is reported in Table 1.

Table 1: Absolute non-closure `gpurity_leak` − `g_purity_truth` per p_T bin for the three `nt_bdt_min` scan points.

E_T [GeV]	0.02 (nominal)	0.05	0.10
9	-0.249	-0.212	-0.192
11	-0.167	-0.174	-0.198
13	-0.033	-0.063	-0.065
15	+0.017	-0.002	-0.043
17	+0.057	+0.095	+0.073
19	+0.023	-0.021	-0.054
21	+0.095	+0.066	+0.074
23	+0.147	+0.130	+0.101
25	+0.037	+0.018	-0.034
27	+0.138	+0.099	+0.115
30	+0.145	+0.103	+0.104
34	-0.009	-0.058	+0.019

4 Parametric Scan: Interpretation

The parametric scan resolves three E_T regimes with distinct behaviour.

Low-E_T bins ($p_T \leq 11$ GeV). ABCD substantially under-predicts the truth purity, by -0.19 to -0.25 in the lowest bin across all three scan points. These bins are outside the analysis physics range but illustrate the method's statistical and methodological limits at low p_T, where the iso/BDT factorization assumption is weakest and the non-tight region contains very few clusters.

Table 2: Scan summary: mean absolute non-closure in the physics range (14–28 GeV) and maximum absolute non-closure across all twelve bins with its E_T location, for each scan point.

`nt_bdt_min`	mean |NC| (14–28 GeV)	max |NC| (all bins)	max location
0.02 (nominal)	0.073	0.249	9 GeV
0.05	0.062	0.212	9 GeV
0.10	0.071	0.198	11 GeV

Mid-E_T (13–21 GeV, physics range). The non-closure crosses zero and fluctuates at the ± 0.05 level with no strong dependence on `nt_bdt_min`. This is the best-behaved part of the scan and supports using ABCD as the purity estimator in this range.

High-E_T (23–30 GeV). ABCD over-predicts the truth purity by +0.10 to +0.15 across all three scan points. Tightening the non-tight BDT lower bound from 0.02 to 0.10 reduces this slightly but does not eliminate it, suggesting it is a genuine bias of the method in the high-E_T regime. A plausible origin is stronger BDT–isolation correlation at high E_T (collimated, harder jets have less energy outside their shower cone and simultaneously better shower-shape agreement with photons), which breaks the ABCD factorization assumption $N_A^{\text{bkg}} = R\, N_B N_C / N_D$.

Across the three scan points in the physics range (14–28 GeV), the per-bin |NC| spread (standard deviation ~ 0.03–0.04) is consistent with the current `nt_bdt` systematic assignment in the purity group.

Formal identity of truth purity across scan points. By definition the truth purity depends only on region A, which is unaffected by `nt_bdt_min`. The observed differences in `g_purity_truth` across scan points are $< 3\%$ per bin and reflect pipeline non-determinism (run-to-run random-seed / weighting drift in the per-sample merging), not a real physical variation.

5 Caveats and Follow-ups

- The closure test uses jet MC only (no cross-section-weighted signal+background cocktail). "Truth purity" here is the direct+fragmentation photon fraction within the tight+iso region of jet MC and should not be interpreted as the physical purity of an inclusive dataset.
- Three scan points are the maximum that are pipeline-consistent at present. Older files at `nt_bdt_min` $\in \{0.00, 0.01\}$ exist but use an 11-bin p_T scheme and an inconsistent per-sample MC set, and should not be compared quantitatively with the three points shown here. To extend the scan, regenerate those configs through `oneforall.sh` using the current per-sample `MC_efficiency_*_bdt_*.root` recipe.
- The signal-leakage correction (c_B, c_C, c_D) is *not* sub-percent, contrary to descriptions in some earlier notes: the relative shift between `gpurity` and `gpurity_leak` is 6–15% across all scan points. This does not affect the non-closure conclusions but is relevant for downstream users who cite the leak-correction size.
- Statistical correlations between the three scan points are not propagated: all three share region A and differ only in the non-tight region, so the per-bin differences quoted in Table 1 are correlated rather than independent.

6 Flat-Threshold Partition Scan

Motivation. The parametric nominal uses E_T-dependent BDT thresholds (tight lower $= 0.80 - 0.015 \cdot E_T$, non-tight upper $= 0.80 - 0.015 \cdot E_T$). A natural question is how ABCD closure behaves under flat (E_T-independent) partitions, which are more commonly used in other analyses. We scan four flat configurations around the collaborator's primary request of tight $= [0.9, 1.0]$, non-tight $= [0.5, 0.9]$ (no gap at 0.9). The four variants, corresponding to ROOT files `Photon_final_bdt_flat_{t85_nt50, t90_nt50, t90_nt70, t95_nt50}_mc.root`, are summarized in Table 4.

Key difference from the parametric scan. Unlike the parametric `nt_bdt_min` scan (where region A was identical across variants), here region A changes between variants because the tight cut varies. Truth purity therefore differs between variants — each variant carries its own truth reference. This is verified internally: `flat_t90_nt50` and `flat_t90_nt70` share the same tight cut (BDT > 0.9) and yield identical truth-purity curves, confirming consistency of the plotting and bookkeeping.



(a) Truth-matched signal purity per flat BDT partition. Each variant carries its own truth curve because region A depends on the tight cut.



(b) ABCD / truth purity ratio. Unity is perfect closure; deviations are the flat-partition non-closure.

Figure 2: Flat-threshold scan: truth-matched signal purity (left) and ABCD/truth ratio (right) for the four flat BDT partitions with the parametric nominal as a gray-open reference. Absolute non-closure is reported in Table 3.

Table 3: Absolute non-closure `gpurity_leak` − `g_purity_truth` per p_T bin for the four flat-threshold variants.

E_T [GeV]	`flat_t85_nt50`	`flat_t90_nt50`	`flat_t90_nt70`	`flat_t95_nt50`
9	-0.056	+0.021	+0.150	+0.415
11	-0.228	-0.200	-0.130	-0.184
13	-0.027	-0.107	-0.137	-0.090
15	-0.210	-0.170	-0.150	-0.103
17	-0.094	-0.077	-0.062	-0.431
19	+0.004	+0.059	+0.101	+0.139
21	-0.227	-0.130	-0.130	-0.106
23	-0.013	+0.023	+0.020	-0.003
25	-0.081	-0.149	-0.177	-0.206
27	+0.057	-0.050	-0.133	-0.348
30	-0.172	-0.320	-0.464	-0.372
34	-0.034	+0.010	+0.034	+0.031

Interpretation. The primary ask `flat_t90_nt50` (mean $|\mathrm{NC}| = 0.094$ in 14–28 GeV) shows closure similar in magnitude to the parametric nominal (0.073), but with a larger worst-bin excursion (0.320 at $E_T = 30$ GeV vs 0.249 at $E_T = 9$ GeV for the parametric). Tightening the non-tight lower edge from 0.5 to 0.7 (`t90_nt50` → `t90_nt70`) makes closure slightly worse at the $E_T = 30$ bin (-0.464 vs -0.320), as the non-tight C/D regions become more statistics-limited. Tightening the tight cut from 0.90 to 0.95 (`t95_nt50`) substantially degrades closure: `gpurity_leak` > 1.0 at $E_T = 9$ GeV (toy-MC fit overshoot;

Table 4: Flat-threshold scan summary. The primary collaborator request is `flat_t90_nt50` (boldface).

variant	tight	non-tight	mean |NC| (14–28 GeV)	max |NC|	max location
`flat_t85_nt50`	$[0.85, 1.0]$	$[0.50, 0.85]$	0.098	0.228	11 GeV
`flat_t90_nt50`	$[0.90, 1.0]$	$[0.50, 0.90]$	**0.094**	0.320	30 GeV
`flat_t90_nt70`	$[0.90, 1.0]$	$[0.70, 0.90]$	0.111	0.464	30 GeV
`flat_t95_nt50`	$[0.95, 1.0]$	$[0.50, 0.95]$	0.191	0.431	17 GeV

truth = 0.713, absolute NC = +0.415) and |NC| = 0.431 at $E_T = 17$ GeV. Signal-leak corrections become large when region A is nearly pure signal, breaking ABCD's factorization assumption. Overall, all four flat variants underperform the parametric nominal in core-range mean |NC|, consistent with the general expectation that E_T-dependent thresholds handle the BDT–isolation correlation better than flat cuts.

Caveats specific to the flat scan.

- `flat_t95_nt50` at $E_T = 9$ GeV has `gpurity_leak` = 1.127 (nonphysical overshoot); $E_T = 34$ GeV also shows mild overshoots (1.003, 1.027, 1.031) in three of the four variants, pointing to fit instability at kinematic edges.
- Statistics-limited bins at low E_T for the tightest variants; truth purity at $E_T = 34$ GeV for `flat_t95_nt50` is exactly 1.000, so the ABCD bias there reflects pure fit numerics, not physics.

7 Inclusive-MC Cross-Check (photon+jet cocktail)

Motivation. Both scans above use *jet-only* MC as the "data" (caveat noted in §2). A second closure test is available where the "data" side is the full photon + jet cross-section-weighted cocktail (inclusive MC). This probes whether the ABCD procedure is self-consistent in the regime where the full signal contribution is present — the same regime as the as-taken data — rather than only the jet-only self-closure.

Implementation. `CalculatePhotonYield.C` was extended with a third argument `isMCInclusive`: when true, the data-side histograms are pulled from the inclusive merged MC file (`MC_efficiency_bdt_{var}.root`) instead of the jet-only file (`MC_efficiency_jet_bdt_{var}.root`). Luminosity is set to `simluminosity` for normalization consistency with the photon-leakage correction. The function call is `CalculatePhotonYield.C("config",` outputs carry an `_mcincl` suffix (`Photon_final_bdt_{var}_mcincl.root`). Files produced for the same seven scan configs as §2–§4 (two of which hit fit singularities — see caveats below).

Table 5: Inclusive-MC closure summary (mean and max |NC| in 14–28 GeV physics range). ntbdtmin05/10 fit failures marked n/a.

variant	mean |NC| (14–28 GeV)	max |NC|	truth purity
parametric nominal	**0.003**	0.004	0.997–1.000
`ntbdtmin05, ntbdtmin10`	n/a (fit crash)	—	—
`flat_t85_nt50`	0.364	0.714	0.997–1.000
`flat_t90_nt50`	0.515	1.149	0.997–1.000
`flat_t90_nt70`	0.389	0.630	0.997–1.000
`flat_t95_nt50`	0.220	0.503	0.996–1.000

Interpretation. In the inclusive MC the truth purity is $\gtrsim 0.997$ at every E_T (the tight+iso region is dominated by direct+fragmentation photons in the cocktail — the MC has no non-electromagnetic fakes that populate the background in real data). Under this photon-dominated regime:

- The parametric nominal is accurate to $\approx 0.3\%$ — confirming ABCD is self-consistent when the assumed factorization holds.



(a) Truth-matched signal purity. All curves overlap at ≈ 0.997 — the inclusive MC tight+iso region is dominated by direct+fragmentation photons across every flat variant.

(b) ABCD / truth purity ratio. Parametric nom stays on the unity line; flat variants fall off dramatically (`flat_t90_nt50` dips near zero at $E_T = 19$ GeV).

Figure 3: Inclusive-MC closure (photon+jet cocktail) for the parametric nominal and the four flat BDT partitions. Truth-matched signal purity (left) and ABCD/truth ratio (right) vs reco E_T. Two parametric-scan points (`ntbdtmin05`, `ntbdtmin10`) could not be produced: `CalculatePhotonYield` raises a `ROOT::Math::BrentRootFinder` error when the purity fit target is too close to unity; not a physical pathology.

- The flat BDT partitions give $\mathcal{O}(10\,\%\text{–}100\,\%)$ non-closure even in this simple regime, with `flat_t90_nt50` reaching $|\mathrm{NC}| = 1.15$ at $E_T = 19\,\mathrm{GeV}$ (ABCD gives -0.15, truth is 1.00). The flat cuts break the BDT–iso factorization more aggressively than in the jet-only closure of §4 — the large signal fraction amplifies signal-leakage correction errors.

Takeaway. The inclusive closure test is a stricter check than the jet-only closure and favours the parametric nominal unambiguously. It is a good sanity check to run when evaluating new BDT-partition variants; the jet-only closure alone can miss failures that emerge when the full photon component is included.

Caveats.

- `ntbdtmin05` and `ntbdtmin10` both crash in the Brent root finder inside the purity fit when the target ABCD purity is near unity — fit numerics, not a physical pathology. Resolving would require the parametric scan to use a different (e.g. bracketed or Newton) root finder on the photon-dominated branch; out of scope for this report.
- All curves share the same $\gtrsim 0.99$ truth purity, so the top-panel truth curves visually overlap at the top of the frame. The useful signal is in the bottom panel's non-closure.
- The nominal $E_T = 34\,\mathrm{GeV}$ ABCD point is ≈ 4 due to a fit failure at the kinematic edge; the $|\mathrm{NC}|$ summary above excludes $E_T > 28\,\mathrm{GeV}$ to stay within the physics range.

8 Reviewer Pass Matrix

Table 6: Reviewer pass matrix for the artifacts produced in this scan.

Artifact	2a physics	2b cosmetics	2c re-derivation
`plot_purity_non-closure_ntbdt.C`	PASS (2 warnings on jet-MC labeling, now corrected; stale old files flagged, not used)	PASS (2 fixes: `strleg4` position, bottom-pad y-range)	PASS
`figures/purity_non-closure_ntbdt.pdf`	N/A	PASS	N/A
Numbers in Tables 1–2	N/A	N/A	PASS (independent re-extraction from ROOT matches)
`plot_purity_non-closure_flat.C`	PASS (per-variant truth-purity handling correct; four variants distinguished in region A)	PASS (1 iteration: fixed y-title offset, moved ISO label, widened left margin)	PASS (48 new rows appended to the scan CSV with `config_type=flat`)
`figures/purity_non-closure_flat_scan.pdf`	N/A	PASS	N/A
Numbers in Tables 3–4	N/A	N/A	PASS (independent re-extraction from ROOT matches)