


Elective Course

Course Code: CS4103

Autumn 2025-26



## Lecture #30

# Artificial Intelligence for Data Science

## Week-8:

### Introduction to Probabilistic Reasoning [Part-II]

Reasoning Under Uncertainty and Probability Theory

Course Instructor:

Dr. Monidipa Das

Assistant Professor

Department of Computational and Data Sciences

Indian Institute of Science Education and Research Kolkata, India 741246

## Bayesian Network



- Probabilistic graphical model
- Represents **dependence/independence** via *directed acyclic graph* (DAG)
- Structure of the graph  $\Leftrightarrow$  Conditional independence relations

In general,

$$P(X_1, X_2, \dots, X_N) = \prod P(X_i | \text{parents}(X_i))$$


*The full joint distribution*

*The graph-structured approximation*

- Components of Bayesian network

- *Qualitative*: Graph structure
- *Quantitative*: Numerical probabilities

$$P(A, B, C) = P(C | A, B)P(A)P(B)$$



## Example




### Variables in the study:

1. Rush-hour (yes/no)
2. Bad-weather (yes/no)
3. Accident (yes/no)
4. Traffic-jam (heavy/light)
5. Miss-Flight (yes/no)



50 km/hr

65 km




If I leave the home at 2:30 PM  
will I be able to catch the flight at 7:30 PM?

Dr. Monidipa Das, Department of CDS, IISER Kolkata

## Example



### Qualitative Component



Node Name	Type	Values
Rush-Hour (R)	Binary	{Yes (Y), No (N)}
Bad-Weather (W)	Binary	{Yes (Y), No (N)}
Accident (A)	Binary	{Yes (Y), No (N)}
Traffic-Jam (T)	Binary	{Heavy (H), Light (L)}
Miss-Flight (M)	Binary	{Yes (Y), No (N)}

Dr. Monidipa Das, Department of CDS, IISER Kolkata

## Example




### Quantitative Component

$P(R=Y)$
0.25

$P(W=Y)$
0.5

W	$P(A=Y W)$
Y	0.75
N	0.10

T	$P(M=Y T)$
H	0.85
L	0.05



R	W	A	$P(T=H R, W, A)$
Y	Y	Y	0.99
Y	Y	N	0.92
Y	N	Y	0.96
Y	N	N	0.85
N	Y	Y	0.95
N	Y	N	0.70
N	N	Y	0.90
N	N	N	0.05

$$P(R, W, A, T, M) = P(R) \cdot P(W) \cdot P(A|W) \cdot P(T|R, W, A) \cdot P(M|T)$$

Dr. Monidipa Das, Department of CDS, IISER Kolkata

## Advantage





- If we have a Bayesian network, with a maximum of  $k$  parents for any node, then we need  $O(n \cdot 2^k)$  probabilities
  - Bayesian network
    - $n = 40, k = 3$ : need 320 probabilities
  - Full joint distribution
    - $n = 40$ : need  $10^{12}$  probabilities for full joint distribution

Dr. Monidipa Das, Department of CDS, IISER Kolkata

# Conditional Independence Relations in Bayesian Network



1. Each node is conditionally independent of its non-descendants given its parents
2. Each node is conditionally independent of all others given its **Markov blanket**: (parents + children + children's parents)



Dr. Monidipa Das, Department of CDS, IISER Kolkata


## Conditional Independence



### Conditionally independent effects:

$$\begin{aligned}
 P(A, B, C) &= P(A|C)P(B|C)P(C) \\
 P(A, B, C) &= P(A|B, C)P(B, C) \\
 &= P(A|B, C)P(B|C)P(C) \\
 &= P(A|C)P(B|C)P(C)
 \end{aligned}$$


A and B are conditionally independent Given C



### Independent Causes:


$$P(A, B, C) = P(C|A, B)P(A)P(B)$$

A and B are (marginally) independent but become dependent once C is known



### Markov dependence:

$$P(A, B, C) = P(B|C)P(C|A)P(A)$$



Dr. Monidipa Das, Department of CDS, IISER Kolkata


# Inference in Bayesian Networks



- Consider answering a query in a Bayesian Network
  - $Q$  = set of query variables
  - $e$  = evidence (set of instantiated variable-value pairs)
  - Inference = computation of conditional distribution  $P(Q|e)$
- Examples**
  - $P(\text{MissFlight} | \text{BadWeather} = Y)$
  - $P(\text{MissFlight} | \text{RushHour} = N)$
  - $P(\text{TrafficJam}, \text{Accident} | \text{BadWeather} = N, \text{RushHour} = Y)$

Dr. Monidipa Das, Department of CDS, IISER Kolkata

# Inference in Bayesian Networks




**Exact Method**

$$\begin{aligned}
 P(\text{MissFlight} | \text{BadWeather} = Y) &\equiv P(M | W = Y) \\
 &= P(M, W = Y) / P(W = Y) = \gamma \cdot P(M, W = Y) = \gamma \sum_R \sum_A \sum_T P(R, W = Y, A, T, M) \\
 &= \gamma \sum_R \sum_A \sum_T P(R) \cdot P(W = Y) \cdot P(A | W = Y) \cdot P(T | R, W = Y, A) \cdot P(M | T)
 \end{aligned}$$

Dr. Monidipa Das, Department of CDS, IISER Kolkata

# Inference in Bayesian Networks



$$P(\text{MissFlight} | \text{BadWeather} = Y, \text{Accident} = N) \equiv P(M | W = Y, A = N)$$

$$= P(M, W = Y, A = N) / P(W = Y, A = N) = \gamma \cdot P(M, W = Y, A = N) = \gamma \sum_T \sum_R P(R, W = Y, A = N, T, M)$$

$$= \gamma (P(R = Y, W = Y, A = N, T = H, M) + P(R = N, W = Y, A = N, T = H, M) + P(R = Y, W = Y, A = N, T = L, M) + P(R = N, W = Y, A = N, T = L, M))$$


Dr. Monidipa Das, Department of CDS, IISER Kolkata

# Inference in Bayesian Networks



$$P(\text{MissFlight} = Y | \text{BadWeather} = Y, \text{Accident} = N) \equiv P(M = Y | W = Y, A = N)$$

$$= \gamma (P(R = Y, W = Y, A = N, T = H, M = Y) + P(R = N, W = Y, A = N, T = H, M = Y) + P(R = Y, W = Y, A = N, T = L, M = Y) + P(R = N, W = Y, A = N, T = L, M = Y))$$



$$= \gamma (P(R = Y) \cdot P(W = Y) \cdot P(A = N | W = Y) \cdot P(T = H | R = Y, W = Y, A = N) \cdot P(M = Y | T = H) + P(R = N) \cdot P(W = Y) \cdot P(A = N | W = Y) \cdot P(T = H | R = N, W = Y, A = N) \cdot P(M = Y | T = H) + P(R = Y) \cdot P(W = Y) \cdot P(A = N | W = Y) \cdot P(T = L | R = Y, W = Y, A = N) \cdot P(M = Y | T = L) + P(R = N) \cdot P(W = Y) \cdot P(A = N | W = Y) \cdot P(T = L | R = N, W = Y, A = N) \cdot P(M = Y | T = L))$$

$$= \gamma ((0.25 \cdot 0.5 \cdot 0.25 \cdot 0.92 \cdot 0.85) + (0.75 \cdot 0.5 \cdot 0.25 \cdot 0.70 \cdot 0.85) + (0.25 \cdot 0.5 \cdot 0.25 \cdot 0.08 \cdot 0.05) + (0.75 \cdot 0.5 \cdot 0.25 \cdot 0.3 \cdot 0.05))$$

$$= \gamma (0.0244 + 0.0558 + 0.0001 + 0.0014)$$

$$= 0.082\gamma$$


Dr. Monidipa Das, Department of CDS, IISER Kolkata

# Inference in Bayesian Networks



$$P(\text{MissFlight} = N | \text{BadWeather} = Y, \text{Accident} = N) \equiv P(M = N | W = Y, A = N)$$

$$= \gamma(P(R = Y, W = Y, A = N, T = H, M = N) + \\ P(R = N, W = Y, A = N, T = H, M = N) + P(R = \\ Y, W = Y, A = N, T = L, M = N) + P(R = N, W = \\ Y, A = N, T = L, M = N))$$



$$= \gamma(P(R = Y).P(W = Y).P(A = N | W = Y).P(T = H | R = Y, W = Y, A = N).P(M = N | T = H) + \\ P(R = N).P(W = Y).P(A = N | W = Y).P(T = H | R = N, W = Y, A = N).P(M = N | T = H) + P(R = Y).P(W = \\ Y).P(A = N | W = Y).P(T = L | R = Y, W = Y, A = N).P(M = N | T = L) + P(R = N).P(W = \\ Y).P(A = N | W = Y).P(T = L | R = N, W = Y, A = N).P(M = N | T = L))$$

$$= \gamma((0.25 * 0.5 * 0.25 * 0.92 * 0.15) + \\ (0.75 * 0.5 * 0.25 * 0.70 * 0.15) + (0.25 * 0.5 * 0.25 * 0.08 * 0.95) + (0.75 * 0.5 * 0.25 * 0.3 * 0.95))$$

$$= \gamma(0.004 + 0.0098 + 0.0024 + 0.0267)$$

$$= 0.043\gamma$$

Dr. Monidipa Das, Department of CDS, IISER Kolkata

# Inference in Bayesian Networks



$$P(\text{MissFlight} = Y | \text{BadWeather} = Y, \text{Accident} = N) + \\ P(\text{MissFlight} = N | \text{BadWeather} = Y, \text{Accident} = N) = 1$$

$$\Rightarrow P(M = Y | W = Y, A = N) + P(M = N | W = Y, A = N) = 1$$

$$\Rightarrow 0.082\gamma + 0.043\gamma = 1$$

$$\Rightarrow 0.125\gamma = 1$$

$$\Rightarrow \gamma = 8$$

$$\therefore P(\text{MissFlight} = Y | \text{BadWeather} = Y, \text{Accident} = N) = 0.082 * 8 = 0.656$$


$$\therefore P(\text{MissFlight} = N | \text{BadWeather} = Y, \text{Accident} = N) = 0.043 * 8 = 0.344$$

Dr. Monidipa Das, Department of CDS, IISER Kolkata

## Do it now

15

Consider the following Bayesian Network. Given evidence about C which of the following pair of variables are conditionally independent.




- ☐ A and B
- ☐ A and D
- ☐ D and E
- ☐ A and F

Dr. Monidipa Das, Department of CDS, IISER Kolkata

## Do it now

16

Consider the following Bayesian Network.  
Which of the following variables belong to Markov Blanket of F.



- ☐ A
- ☐ B
- ☐ C
- ☐ D


Dr. Monidipa Das, Department of CDS, IISER Kolkata



## Do it now



How many minimum parameters are required to represent the following Bayesian Network, if each variable is Boolean?



Dr. Monidipa Das, Department of CDS, IISER Kolkata

## Solving Homework



- Given the full joint distribution shown below, calculate the following:
  - a.  $P(\text{toothache})$ .
  - b.  $P(\text{Cavity})$ .
  - c.  $P(\text{Cavity} \mid \text{toothache} \vee \text{catch})$ .

	<i>toothache</i>		$\neg$ <i>toothache</i>	
	<i>catch</i>	$\neg$ <i>catch</i>	<i>catch</i>	$\neg$ <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
$\neg$ <i>cavity</i>	0.016	0.064	0.144	0.576


Solved during lecture session

Dr. Monidipa Das, Department of CDS, IISER Kolkata


# Solving Homework




Given the following networks, write the graph structured approximation of the full joint distributions



(a)



(b)



(c)

Solved during lecture session

Dr. Monidipa Das, Department of CDS, IISER Kolkata



## Questions?

Slide Content taken from  
Prof. Cesare Tinelli, Prof. Stuart Russell, and  
Prof. Jim Martin

Dr. Monidipa Das, Department of CDS, IISER Kolkata