


Slowing down hydrolysis of cisplatin: Consequences!

- The hydroxo / water ligands are much more reactive to substitution than the chloride ligands
- The hydrolysis rate is mainly determined by the *trans* effect of the ligands *trans* to Cl^-
- Steric hindrance can also slow rates of ligand substitution



A undergoes hydrolysis 6x slower than **B**


The general order in bond strength is $\text{Pt-C (sp)} > \text{Pt-O} > \text{Pt-N} > \text{Pt-C (sp}^3\text{)}$

Role of metal : adding targets to existing organic drugs

Jaouen and coworkers:


S. Top et al. *Chem. Comm.* 1996, 955

S. Top et al. *J. Organometal. Chem.* 1997, 541, 355




Ferrocifen : Both effects coexist together


Anti-tumor and Anti-oestrogen properties



Role of metal is adding targets to existing organic drugs



3 (Z+E)



4-Hydroxytamoxifen

- Binding affinity < hydroxytamoxifen for **3** (sterics of ferrocenyl moiety)
- **3** > lipophilic
- Antiproliferative activity on breast cancer cells : **3** = OH-TAM for ER(+)
- Ferrocifen show remarkable **antiproliferative behaviour even against ER- tumors**


S. Top et al. *J. Organometal. Chem.* 2001, 637, 500

S. Top et al. *Chem. Eur. J.* 2003, 9, 5223

Protein Kinases

Protein Kinases:

- Phosphorylation of proteins : turn them on or off





- Due to their involvement in various forms of cancers, PTKs have become prominent targets for therapeutics
- Regulate the majority of cellular pathways e.g DNA replication, cell growth
- Most kinases contain a 250-300 amino acid domain with a conserved core structure, comprising a binding pocket for ATP
- These domains are more or less homologous

Blume-Jensen. P.; Hunter, T. *Nature*, 2002, 411, 355
Fischer, P.M. *Curr. Med. Chem.* 2004, 11, 1583


ATP Binding

- ATP-binding site is an ubiquitous “receptor” in nature
- Most kinase inhibitors mimic mainly the adenine portion of ATP
- Approach is limited in terms of selectivity



Fischer, P.M. *Curr. Med. Chem.* 2004, 11, 1583

Staurosporine metal complexes



- copying the structural features of small organic molecule inhibitors
- metal plays solely a structural role
- access to new areas of chemical space

Zhang, L. Carroll, P. Meggers, E.; *Org. Lett.* 2004, 6, 521

Bregman, H. Williams, G. S. Meggers, E.; *Synthesis*, 2005, 9, 1521


Bregman, H, Carroll, P.J. Meggers, E. *J. Am. Chem. Soc.* 2006, 128, 877

Analysis of IC₅₀ values: Ruthenium worsening kinase inhibition or enhancing selectivity

Inhibition of some protein kinases with the various compounds (in μM)


compound	Ab1	RSK1	Src	PKCa	ZAP70
staurosporine	2	<1	<1	<1	<1
7	25	30	>100	>100	>100
6	20	25	60	>100	50
1	10	8	30	>100	40
2	2	8	40	>100	30
3	5	8	30	50	40

POTENCY and SPECIFICITY




Zhang, L. Caroll, P. Meggers, E. *Org. Lett.* 2004, 6, 521



What is PDT (Photodynamic Therapy)



TOOKAD: Padeliporfin



PADE-001



Padeliporfin di-potassium

