

Multi-player minimum cost flow problems with nonconvex costs and integer flows

Shuvomoy Das Gupta Lacra Pavel

Department of Electrical and Computer Engineering
University of Toronto

55th IEEE Conference on Decision and Control, 2016

Outline

Introduction

Algorithm construction


Algorithm description

Numerical example

Network Flow Problems

- ▶ *Network flow problems*: optimization problems associated with underlying directed network.
- ▶ They arise in numerous application settings and in many forms.
- ▶ Some common application areas: communication networks, transportation system, social network, power system, electro-mechanical systems *etc.*
- ▶ *The minimum cost flow problem* is the most fundamental among network flow problems.

Minimum Cost Flow Problem




Minimum Cost Flow Problem

$$\begin{aligned} &\text{minimize}_x \quad \sum_{i \in \{1, \dots, n\}} c_i f_i(x_i) \quad \backslash * \text{Default Case: } f_i(x_i) = x_i * \backslash \\ &\text{subject to} \quad Ax = b \\ &\quad \quad \quad 0 \preceq x \preceq u \\ &\quad \quad \quad x \in \mathbf{Z}^n. \end{aligned} \tag{1}$$

- ▶ Here, the network has n arcs, $m+1$ nodes.
- ▶ A : reduced node-arc incidence matrix, dimension $m \times n$, full row rank.
- ▶ We re-index the m linearly independent columns of A as the first m columns.
- ▶ b : represents supplies to the nodes at different points.
- ▶ u : upper bound vector for the flow.

Extension of Min. Cost Flow to a Multi-player Setup

- ▶ With **each arc** of the network graph we associate **one player**.
- ▶ Each of the players is trying to minimize its **nonconvex cost function**, subject to the network flow constraints.
- ▶ Our goal is to seek an **efficient solution concept** in this multi-player problem.



Multi-player Extension

- ▶ The goal of the i th player for $i \in [n] = \{1, \dots, n\}$, given other players' strategies $x_{-i} \in \mathbf{Z}^{n-1}$, is to solve:

$$\begin{aligned} & \text{minimize}_{x_i} && f_i(x_i) \quad \text{\textit{*proper: practically can be anything*}} \\ & \text{subject to} && A(x_i, x_{-i}) = b \quad \text{\textit{*The constraints}} \\ & && 0 \preceq (x_i, x_{-i}) \preceq u \quad \text{\textit{couple the players*}} \\ & && x \in \mathbf{Z}^n. \end{aligned}$$

(2)

Solution Concepts in Consideration

- ▶ A *vector optimal solution* that minimizes all the objectives simultaneously is unlikely to exist.
- ▶ The celebrated *Nash equilibrium* is also not very efficient in our setup because:
 - the constraint set of the problem has equality constraints, thus making any feasible point a Nash equilibrium, and
 - posteriori some of the players may decide to deviate from the Nash equilibrium in order to reduce their costs even more at the expense of the rest of the players.
- ▶ A more effective solution concept is the *Pareto optimal point*.

Pareto Optimal Point

A *Pareto optimal point* is a solution concept where none of the objective functions can be improved without worsening some of the other objective values.

Definition

(Pareto Optimal Point) In problem 2, a point $x^* \in P$ is Pareto optimal if it satisfies the following: there *does not* exist another point $\tilde{x} \in P$ such that

$$(\forall i \in [n]) \quad f_i(\tilde{x}_i) \leq f_i(x_i^*),$$

with at least one index $j \in [n]$ satisfying $f_j(\tilde{x}_j) < f_j(x_j^*)$.

Pareto Optimal Point

A *Pareto optimal point* is a solution concept where none of the objective functions can be improved without worsening some of the other objective values.

Definition

(Pareto Optimal Point) In problem 2, a point $x^* \in P$ is Pareto optimal if it satisfies the following: there *does not* exist another point $\tilde{x} \in P$ such that

$$(\forall i \in [n]) \quad f_i(\tilde{x}_i) \leq f_i(x_i^*),$$

with at least one index $j \in [n]$ satisfying $f_j(\tilde{x}_j) < f_j(x_j^*)$.

Related Work

- ▶ Our problem setup does not seem to be investigated in existing literature.
- ▶ Nonconvex network flow problems for very specific cost functions (Magnanti1984, Graves1985, Daskin2011, He2015).
- ▶ Integer multi-commodity flow problems (Brunetta2000, Ozdaglar2004).

Outline

Introduction

Algorithm construction

Algorithm description

Numerical example

Decoupling the Last $n - m$ Players

Theorem

(Existence of a decoupling variable) Denote $B = [A_1 | A_2 | \cdots | A_m]$. The equality constraint set $Q = \{x \in \mathbf{Z}^n \mid Ax = b\}$ is nonempty and for any vector x , $x \in Q$ is equivalent to saying that there exists a $z \in \mathbf{Z}^{n-m}$ such that

$$x = (d_1 - h_1^T z, \dots, d_m - h_m^T z, z_1, \dots, z_{n-m}), \quad (3)$$

where d_i is the i th component of $d = B^{-1}b$, and $h_i^T \in \mathbf{Z}^{n-m}$ is the i th row of $B^{-1}A_{[1:m, m+1:n]}$.

Decoupled Problems for the $n - m$ Players

- ▶ In z we can decouple the optimization problems for players $m + 1, m + 2, \dots, n$ as follows

$$\begin{aligned} & \text{minimize}_{z_i} && f_i(z_i) \\ & \text{subject to} && 0 \leq z_i \leq u_i \\ & && z_i \in \mathbf{Z}. \end{aligned} \tag{4}$$

- ▶ Set of different optimal solutions for player $m + i$ for $i \in [n - m]$ is

$$D_i = \{z_{i,1}, z_{i,2}, \dots, z_{i,p_i}\}.$$

- ▶ Define, $D = \times_{i=1}^{n-m} D_i \neq \emptyset$

Consensus Reformulation for the First m players

- ▶ We provide each player $i \in [m]$ with its *own local copy* of z , denoted by $z^{(i)} \in \mathbf{Z}^{n-m}$, which acts as its decision variable.
- ▶ For any $i \in [m]$, $x_i = d_i - h_i^T z^{(i)}$.
- ▶ The copy $z^{(i)}$ has to be in *consensus* with the rest of the first m players:

$$z^{(i)} = z^{(j)} \quad \forall j \in [m] \setminus \{i\}.$$

- ▶ The copy $z^{(i)}$ has to satisfy the *flow bound constraints*, i.e., $0 \leq d_i - h_i^T z^{(i)} \leq u_i$ for all $i \in [m]$.
- ▶ For the last $n - m$ players $z_i \in D_i$, so:

$$z^{(i)} \in D$$

Consensus Reformulation for the First m players

For all $i \in [m]$, the i th player's optimization problem in variable $z^{(i)}$ can be written as:

$$\begin{aligned} & \text{minimize}_{z^{(i)}} \quad \bar{f}_i(z^{(i)}) = f_i(d_i - h_i^T z^{(i)}) \\ & \text{subject to} \quad z^{(i)} = z^{(j)}, \quad j \in [m] \setminus \{i\} \\ & \quad \quad \quad 0 \leq d_i - h_i^T z^{(i)} \leq u_i \\ & \quad \quad \quad z^{(i)} \in D. \end{aligned} \tag{5}$$

$$q_i(z^{(i)}) = (d_i - h_i^T z^{(i)})(d_i - h_i^T z^{(i)} - 1) \cdots (d_i - h_i^T z^{(i)} - u_i) = 0, \tag{6}$$

$$r_j(z^{(i)}) = (z_j^{(i)} - z_{j,1})(z_j^{(i)} - z_{j,2}) \cdots (z_j^{(i)} - z_{j,p_i}) = 0, \quad j \in [n-m]. \tag{7}$$

$$\mathcal{F} = \{z \in \mathbf{Z}^{n-m} \mid (\forall k \in [m]) \ q_k(z) = 0, (\forall j \in [n-m]) \ r_j(z) = 0\}.$$

Consensus Reformulation for the First m players

For all $i \in [m]$, the i th player's optimization problem in variable $z^{(i)}$ can be written as:

$$\begin{aligned} & \text{minimize}_{z^{(i)}} \quad \bar{f}_i \left(z^{(i)} \right) = f_i(d_i - h_i^T z^{(i)}) \\ & \text{subject to} \quad z^{(i)} = z^{(j)}, \quad j \in [m] \setminus \{i\} \\ & \quad \quad \quad 0 \leq d_i - h_i^T z^{(i)} \leq u_i \\ & \quad \quad \quad z^{(i)} \in D. \end{aligned} \tag{5}$$

$$q_i(z^{(i)}) = (d_i - h_i^T z^{(i)})(d_i - h_i^T z^{(i)} - 1) \cdots (d_i - h_i^T z^{(i)} - u_i) = 0, \tag{6}$$

$$r_j(z^{(i)}) = (z_j^{(i)} - z_{j,1})(z_j^{(i)} - z_{j,2}) \cdots (z_j^{(i)} - z_{j,p_i}) = 0, \quad j \in [n-m]. \tag{7}$$

$$\mathcal{F} = \{z \in \mathbb{Z}^{n-m} \mid (\forall k \in [m]) \ q_k(z) = 0, (\forall j \in [n-m]) \ r_j(z) = 0\}.$$

Consensus Reformulation for the First m players

For all $i \in [m]$, the i th player's optimization problem in variable $z^{(i)}$ can be written as:

$$\begin{aligned} & \text{minimize}_{z^{(i)}} \quad \bar{f}_i \left(z^{(i)} \right) = f_i(d_i - h_i^T z^{(i)}) \\ & \text{subject to} \quad z^{(i)} = z^{(j)}, \quad j \in [m] \setminus \{i\} \\ & \quad \quad \quad 0 \leq d_i - h_i^T z^{(i)} \leq u_i \\ & \quad \quad \quad z^{(i)} \in D. \end{aligned} \tag{5}$$

$$q_i(z^{(i)}) = (d_i - h_i^T z^{(i)})(d_i - h_i^T z^{(i)} - 1) \cdots (d_i - h_i^T z^{(i)} - u_i) = 0, \tag{6}$$

$$r_j(z^{(i)}) = (z_j^{(i)} - z_{j,1})(z_j^{(i)} - z_{j,2}) \cdots (z_j^{(i)} - z_{j,p_i}) = 0, \quad j \in [n-m]. \tag{7}$$

$$\mathcal{F} = \{z \in \mathbf{Z}^{n-m} \mid (\forall k \in [m]) \ q_k(z) = 0, (\forall j \in [n-m]) \ r_j(z) = 0\}.$$

Consensus Reformulation for First m players

For $i \in [m]$, i.e., each of these players are optimizing over a *common constraint set* \mathcal{F} :

$$\begin{aligned} & \underset{z^{(i)}}{\text{minimize}} && \bar{f}_i \left(z^{(i)} \right) \\ & \text{subject to} && z^{(i)} \in \mathcal{F}. \end{aligned} \tag{8}$$

- ▶ When \mathcal{F} is nonempty? How to find the points in it?
- ▶ A little algebraic geometry will take us a long way...

A Little Algebraic Geometry

- ▶ The *ideal* generated by polynomials $f_1, f_2, \dots, f_m \in \mathbf{C}[x]$ is the set

$$\mathbf{ideal}\{f_1, \dots, f_m\} = \left\{ \sum_{i=1}^m h_i f_i \mid (\forall i \in [m]) \ h_i \in \mathbf{C}[x] \right\}.$$

** analogous to span of vectors **

- ▶ A *Groebner basis* G_{\succ} is particular kind of generating set of an ideal I over a field $\mathbf{C}[x]$ ** analogous to basis set of a span **
- ▶ *Reduced Groebner basis* $G_{\text{reduced}, \succ}$ is the most compact Groebner basis for an ideal I . ** analogous to orthonormal basis of a span **
- ▶ There are many computer algebra packages to compute reduced Groebner basis such as Macaulry2, SINGULAR, FGb, Maple, Mathematica etc.

When is \mathcal{F} Nonempty?

Theorem

The set \mathcal{F} is nonempty if and only

$$G_{\text{reduced}, \succ} \neq \{1\},$$

*where $G_{\text{reduced}, \succ}$ is the reduced Groebner basis of
ideal $\{q_1, \dots, q_m, r_1, \dots, r_{n-m}\}$ with respect to any ordering.*

Computing Points in \mathcal{F}

► For $i = [n - m]$ calculate

$$G_{n-m-i} = G_{\text{reduced}, \succ_{\text{lex}}} \cap \mathbf{C}[z_{n-m-i+1}, z_{n-m-i+2}, \dots, z_{n-m}].$$

Theorem

$$G_0 = \mathcal{F}.$$

Outline

Introduction

Algorithm construction

Algorithm description

Numerical example

Finding the Pareto optimal points from \mathcal{F}

```

for  $i \in [m]$ 
     $X_i := d_i - h_i^T \mathcal{F}$     \* The inverse operator is denoted  $X_i^{-1}$  * \
end for

    Sort the elements of the  $\{X_i\}_{i=1}^m$ s with respect to cardinality of the elements in a
    descending order.

    Denote the index set of the sorted set by  $\{s_1, \dots, s_m\}$  such that  $|X|_{s_1} \geq \dots \geq |X|_{s_m}$ .
for  $i \in [m]$ 
     $X_{s_i}^* := \operatorname{argmin}_{x_{s_i} \in X_{s_i}} f_{s_i}(x_{s_i})$     \* Univariate optimization problem * \
     $\mathcal{F}_{s_i}^* := \bigcup_{x_{s_i} \in X_{s_i}^*} (X_{s_i}^*)^{-1}(x_{s_i})$     \* Lemma. The set  $\mathcal{F}_{s_i}^*$  is nonempty * \
    if  $i \leq m$ 
         $X_{s_{i+1}} := \left\{ d_{s_{i+1}} - h_{s_{i+1}}^T z \mid z \in \mathcal{F}_{s_i}^* \right\}.$ 
    end if
end for
return  $\mathcal{F}_{s_m}^*$     \* Theorem. Any member of  $\mathcal{F}_{s_m}^*$  is a Pareto optimal point
    * \

```


Finding the Pareto optimal points from \mathcal{F}

```

for  $i \in [m]$ 
     $X_i := d_i - h_i^T \mathcal{F}$     \* The inverse operator is denoted  $X_i^{-1}$  * \
end for

    Sort the elements of the  $\{X_i\}_{i=1}^m$ s with respect to cardinality of the elements in a
    descending order.

    Denote the index set of the sorted set by  $\{s_1, \dots, s_m\}$  such that  $|X|_{s_1} \geq \dots \geq |X|_{s_m}$ .

for  $i \in [m]$ 
     $X_{s_i}^* := \operatorname{argmin}_{X_{s_i} \in \mathcal{X}_{s_i}} f_{s_i}(X_{s_i})$     \* Univariate optimization problem * \
     $\mathcal{F}_{s_i}^* := \bigcup_{X_{s_i} \in \mathcal{X}_{s_i}^*} (X_{s_i}^*)^{-1}(X_{s_i})$     \* Lemma. The set  $\mathcal{F}_{s_i}^*$  is nonempty * \
    if  $i \leq m$ 
         $X_{s_{i+1}} := \left\{ d_{s_{i+1}} - h_{s_{i+1}}^T z \mid z \in \mathcal{F}_{s_i}^* \right\}.$ 
    end if
end for
return  $\mathcal{F}_{s_m}^*$     \* Theorem. Any member of  $\mathcal{F}_{s_m}^*$  is a Pareto optimal point
    * \

```

Finding the Pareto optimal points from \mathcal{F}

```

for  $i \in [m]$ 
     $X_i := d_i - h_i^T \mathcal{F}$     \* The inverse operator is denoted  $X_i^{-1}$  * \
end for

    Sort the elements of the  $\{X_i\}_{i=1}^m$ s with respect to cardinality of the elements in a
    descending order.

    Denote the index set of the sorted set by  $\{s_1, \dots, s_m\}$  such that  $|X|_{s_1} \geq \dots \geq |X|_{s_m}$ .

for  $i \in [m]$ 
     $X_{s_i}^* := \operatorname{argmin}_{x_{s_i} \in X_{s_i}} f_{s_i}(x_{s_i})$     \* Univariate optimization problem * \
     $\mathcal{F}_{s_i}^* := \bigcup_{x_{s_i} \in X_{s_i}^*} (X_{s_i}^*)^{-1}(x_{s_i})$     \* Lemma. The set  $\mathcal{F}_{s_i}^*$  is nonempty * \
    if  $i \leq m$ 
         $X_{s_{i+1}} := \left\{ d_{s_{i+1}} - h_{s_{i+1}}^T z \mid z \in \mathcal{F}_{s_i}^* \right\}.$ 
    end if
end for
return  $\mathcal{F}_{s_m}^*$     \* Theorem. Any member of  $\mathcal{F}_{s_m}^*$  is a Pareto optimal point
    * \

```

Outline


Introduction

Algorithm construction

Algorithm description

Numerical example

Numerical Example: 16 Players




A multi-player transportation problem

- 5 nodes, 16 arcs
- Nodes 2, 4 are retail centers
- Nodes 1, 3 are warehouses
- Node 5 transshipment node
- Products need to be shipped from warehouses to retail centers
- Arcs represent different modes of shipment from one node to another and carried out by different organization
- Cost of a certain type of shipment depends on the number of products shipped

$$A = \begin{pmatrix} 1 & 0 & 0 & 0 & -1 & -1 & -1 & 1 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 1 & 0 & 0 & -1 & -1 & -1 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & -1 & -1 & -1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & -1 & -1 & -1 \end{pmatrix}$$

$$b = (9, -13, 15, -11), \quad u = (5, 6, 6, 10, 10, 7, 11, 13, 16, 12, 4, 5, 6, 14, 13, 15)$$

Pareto Optimal Solutions



A multi-player transportation problem

- 5 nodes, 16 arcs
- Nodes 2, 4 are retail centers
- Nodes 1, 3 are warehouses
- Node 5 transshipment node
- Products need to be shipped from warehouses to retail centers
- Arcs represent different modes of shipment from one node to another and carried out by different organization
- Cost of a certain type of shipment depends on the number of products shipped

The Pareto optimal points are

$(1, 3, 5, 4, 11, 10, 2, 1, 3, 7, 7, 5)$

and

$(1, 3, 5, 6, 11, 10, 2, 1, 3, 7, 7, 6).$

Concluding Remarks

- ▶ Methodology is numerically efficient when $m < \frac{n}{2}$.
- ▶ Calculating Groebner basis can be numerically challenging for very large system.
- ▶ Based on structure of the network it may happen that \mathcal{F} is empty \Rightarrow penalty based approach.

Thank You!
Questions?

Cost Functions


Player	Cost function
1	$-\frac{x_1^4}{30} - \frac{13x_1^3}{15} + \frac{259x_1^2}{30} - \frac{263x_1}{15} + 1$
2	$\frac{77x_2^5}{120} - \frac{247x_2^4}{24} + \frac{471x_2^3}{8} - \frac{3365x_2^2}{24} + \frac{6779x_2}{60} + 1$
3	$\frac{47x_3^4}{24} - \frac{133x_3^3}{4} + \frac{4897x_3^2}{24} - \frac{2123x_3}{4} + 485$
4	$\frac{323x_4^5}{3360} - \frac{2179x_4^4}{1120} + \frac{47393x_4^3}{3360} - \frac{48709x_4^2}{1120} + \frac{7885x_4}{168} + 5$
5	$(x_5 - 1)^2$
6	$-\frac{x_6^4}{8} + \frac{25x_6^3}{12} - \frac{71x_6^2}{8} + \frac{95x_6}{12} + 10$
7	$ x_7 - 5 $
8	$\frac{11x_8^7}{1260} - \frac{7x_8^6}{36} + \frac{119x_8^5}{72} - \frac{479x_8^4}{72} + \frac{4609x_8^3}{360} - \frac{803x_8^2}{72} + \frac{155x_8}{28} + 1$
9	$-\frac{15}{16}x_9^3 + \frac{365x_9^2}{16} - \frac{2865x_9}{16} + \frac{7315}{16}$
10	$(x_{10} - 10)^2$
11	$\frac{5x_{11}^4}{6} - \frac{35x_{11}^3}{3} + \frac{355x_{11}^2}{6} - \frac{370x_{11}}{3} + 90$
12	$\frac{5x_{12}^4}{6} - \frac{25x_{12}^3}{3} + \frac{175x_{12}^2}{6} - \frac{110x_{12}}{3} + 15$
13	$\frac{5x_{13}^4}{6} - 15x_{13}^3 + \frac{595x_{13}^2}{6} - 280x_{13} + 285$
14	$\frac{5x_{14}^4}{6} - \frac{85x_{14}^3}{3} + \frac{2155x_{14}^2}{6} - \frac{6020x_{14}}{3} + 4165$

Numerical example

Cost Functions

Player	Cost function
15	$ x_{15} - 7 $
16	$\begin{cases} x_{16} + 1, & \text{if } 0 \leq x_{16} \leq 3 \\ 0, & \text{if } 4 \leq x_{16} \leq 6 \\ (x_{16} + 1)^3, & \text{if } 7 \leq x_{16} \leq 9 \\ -\frac{x_{16}^3}{6} + \frac{13x_{16}^2}{2} - \frac{244x_{16}}{3} + 330, & \text{else} \end{cases}$

A Generic Network



The Minimum Cost Flow Problem

- ▶ There is a *directed connected graph* that represents the network.
- ▶ There is *flow* of some commodity along the arcs of the graphs.
- ▶ Each arc incurs a *cost* depending on the amount of flow.
- ▶ The flow is often taken to be *integral*.
- ▶ The goal is to *minimize the total cost* of all flows subject to the *network constraints*.