




香港大學

THE UNIVERSITY OF HONG KONG

A NEW APPROACH TO THE GeV FLARE OF PSR B1259-63/LS2883




Yi Shu-Xu (易疏序), K-S Cheng

Department of Physics

University of Hong Kong


FPS6 @ Wuhan

Introduction of the system




George G. Pavlov et al. 2015

X-ray/TeV emission




PSR B1259-63
spin P: 47.76 ms,


orbit P = 1237 days




Origin of the X-ray/TeV emission





Origin of the X-ray/TeV emission



Origin of the X-ray/TeV emission



Emission in 100 MeV-100 GeV




GeV flare at un-expected orbital phase

Tam 2011, Abdo 2011, Caliendo 2015


Models for GeV emission

- Inverse Compton scattering: first come-to-mind mechanism for GeV emission




Models for GeV emission

- Inverse Compton scattering
- Problem: where comes the soft photons (target)?



Models for GeV emission





Condition of mass transfer from optical companion



Shock front should be inside the Bondi-Hoyle sphere

$$r_{\text{BH}} = \frac{2GM_{\text{p}}}{v_{\text{rel}}^2},$$

$$r_{\text{s}} = \sqrt{\frac{L_{\text{spin}}}{4\pi\rho_{\text{cd}}v_{\text{vel}}^2c}},$$



Location of the circumstellar disk, and phases of mass transfer





Condition of the formation of accretion disk

- The transferred material should have enough specific angular momenta: $r_{circ} > r_{lc}$
- $r_{circ} = \frac{l^2}{GM_p}$
- The angular momenta of the transferred material are due to the density and velocity gradient of the circumstellar disk.

$$l(t) = \frac{(GM_p)^2}{v_{rel}^3} \left(\frac{|\nabla v_{vel}|}{v_{rel}} + \frac{|\nabla \rho_{cd}|}{\rho_{cd}} \right).$$

Formation of the accretion disk

- Phase I: matter kinetic energy redistribution → torus
- Phase II: torus → accretion disk
- Phase III: inner edge of accretion disk decrease until it reaches the inner most radius
- Phase IV: mass and accretion rate decrease



Viscosity time
scale

$$\tau \approx \frac{R_{\text{circ}}}{v_r} \approx \frac{1}{2.7} \alpha^{-4/5} \dot{M}_{\text{acc},16}^{-3/10} m_p^{1/4} R_{\text{circ}}^{5/4} \times 10^6 \text{ s.}$$

Evolution of the accretion disk

$$T = 1.4 \times 10^4 \alpha^{-1/5} \dot{M}_{\text{acc},16}^{3/10} m_p^{1/4} r_{\text{in},10}^{-3/4} \text{ K},$$





Figure 3. The temperature of the inner most region of the accretion disk, as a function of the time after the formation of the accretion disk: From left to right correspond to

Evolution of SED









Figure 5. The evolution of the SED of IC:

Evolution of SED and light curve



Predication




Figure 8. IR/Optical/UV SED from the sys-

Summary of the model

- Matter from circumstellar disk captured by gravity of pulsar
- An accretion disk forms.
- Pulsar wind inverse-Compton scatter the soft photon from accretion disk

