

## Assignment-based Subjective Questions

1. **From your analysis of the categorical variables from the dataset, what could you infer about their effect on the dependent variable?**

**Answer:** The Categorical variable in the dataset after cleaning and evaluation were season, weatherise , holiday , month (mnth), year (yr) and weekday. These were curated using boxplot and later adding and removing i.e. creating combinations out of these to put into model. Following are the effect of each below:

a. **Season**: The boxplot showed that the spring season had least value on cnt whereas fall had a maximum value of cnt. Summer and winter had intermediate value of cnt.

b. **Weathersit**: The boxplot shows that heavy rains/ snow is extremely unfavourable. Highest cnt weathersit was Clear,Partly cloud, Few clouds.

c. **Holiday**: Rental reduced during the holiday

d. **Mnth**: September saw highest no of rentals while December saw least.This observation is on par with the observation made in weathersit.The weather situation in December is usually heavy snow.


e. **Yr** : The number of rental is high in 2019 as compare to 2018

2. **Why is it important to use drop\_first=True during dummy variable creation?**

**Answer:** “drop\_first =True” is important to use , as it helps in reducing the extra column created during dummy variable creation. Hence it reduces the correlations created among dummy variables. This may affect the model adversely and the effect is stronger when the cardinality is smaller. For example iterative models may have trouble converging and lists of variable importances may be distorted.Another reason is, if we have all dummy variables it leads to Multicollinearity between the dummy variables. To keep this under control, we lose one column

3. **Looking at the pair-plot among the numerical variables, which one has the highest correlation with the target variable?**


**Answer:**



“temp” and “atemp” are the two numerical variables which are highly correlated with the target variable (cnt)

#### 4. How did you validate the assumptions of Linear Regression after building the model on the training set?

Answer:



Residuals distribution should follow normal distribution and centred around 0.(mean = 0). We validate this assumption about residuals by plotting a distplot of residuals and see if residuals are following normal distribution or not. The above diagram shows that the residuals are distributed about mean = 0.

5. **Based on the final model, which are the top 3 features contributing significantly towards explaining the demand of the shared bikes?**

**Answer:**

a. temp - coefficient : 0.4709

b. yr - coefficient: 0.2334

c. Weathersit -with value snow/rain - coefficient - 0.3007

## General Subjective Questions

1. **Explain the linear regression algorithm in detail.**

**Answer:** Linear Regression is a machine learning algorithm based on supervised learning. It performs a regression task. Regression models a target prediction value based on independent variables. It is mostly used for finding out the relationship between variables and forecasting. Different regression models differ based on – the kind of relationship between dependent and independent variables, they are considering and the number of independent variables being used.

Linear regression is based on the popular equation “ $y = mx + c$ ”

It assumes that there is a linear relationship between the dependent variable(y) and the predictor(s)/ independent variable(x). In regression, we calculate the best fit line which describes the relationship between the independent and dependent variable.

Regression is performed when the dependent variable is of continuous data type and Predictors or independent variables could be of any data type like continuous, nominal/categorical etc.

Regression method tries to find the best fit line which shows the relationship between the dependent variable and predictors with least error.

In regression, the output/dependent variable is the function of an independent variable and the coefficient and the error term.

Regression is broadly divided into simple linear regression and multiple linear regression.

**1. Simple Linear Regression : SLR** is used when the dependent variable is predicted using only one independent variable.

**2. Multiple Linear Regression : MLR** is used when the dependent variable is predicted using multiple independent variables.

The equation for MLR will be:

$$Y_i = \beta_0 + \beta_1 X_{1i} + \beta_2 X_{2i} + \beta_3 X_{3i} + \dots$$

$\beta_1$  = coefficient for X1 variable  
 $\beta_2$  = coefficient for X2 variable  
 $\beta_3$  = coefficient for X3 variable and so on...


$\beta_0$  is the intercept (constant term).

## 2. Explain the Anscombe's quartet in detail.

**Answer:** Anscombe's Quartet was developed by statistician Francis Anscombe. It includes four data sets that have almost identical statistical features, but they have a very different distribution and look totally different when plotted on a graph. It was developed to emphasise both the importance of

$$Y_i = \beta_0 + \beta_1 X_{1i} + \beta_2 X_{2i} + \beta_3 X_{3i} + \dots$$

graphing data before analysing it and the effect of **outliers** and other **influential observations** on statistical properties



- The first scatter plot (top left) appears to be a simple linear relationship.
- The second graph (top right) is not distributed normally; while there is a relation between them, it's not linear.
- In the third graph (bottom left), the distribution is linear, but should have a different regression line. The calculated regression is offset by the one outlier which exerts enough influence to lower the correlation coefficient from 1 to 0.816.
- Finally, the fourth graph (bottom right) shows an example when one high-leverage point is enough to produce a high correlation coefficient, even though the other data points do not indicate any relationship between the variables.

### 3. What is Pearson's R?

**Answer:** Pearson's  $r$  is a numerical summary of the strength of the linear association between the variables. Its value ranges between  $-1$  to  $+1$ . It shows the linear relationship between two sets of data. In simple terms, it tells us can we *draw a line graph to represent the data*?

$r = 1$  means the data is perfectly linear with a positive slope

$r = -1$  means the data is perfectly linear with a negative slope  $r = 0$  means there is no linear association

### 4. What is scaling? Why is scaling performed? What is the difference between normalized scaling and standardized scaling?

**Answer:** Feature scaling is a method used to normalize or standardize the range of independent variables or features of data. It is performed during the data preprocessing stage to deal with varying values in the dataset. If feature scaling is not done, then a machine learning algorithm tends to weigh greater values, higher and consider smaller values as the lower values, irrespective of the units of the values.

- Normalization is generally used when you know that the distribution of your data does not follow a Gaussian distribution. This can be useful in algorithms that do not assume any distribution of the data like K-Nearest Neighbors and Neural Networks.
- Standardization, on the other hand, can be helpful in cases where the data follows a Gaussian distribution. However, this does not have to be necessarily true. Also, unlike normalization, standardization does not have a bounding range. So, even if you have outliers in your data, they will not be affected by standardization.

### 5. You might have observed that sometimes the value of VIF is infinite. Why does this happen?

**Answer:** **VIF - the variance inflation factor** - The VIF gives how much the variance of the coefficient estimate is being inflated by collinearity.  $(VIF) = 1/(1-R_1^2)$ . If there is perfect correlation, then  $VIF = \text{infinity}$ . Where  $R_1^2$  is the R-square value of that independent variable which we want to check how well this independent variable is explained well by other independent variables- If that independent variable can be explained perfectly by other independent variables, then it will have perfect correlation and its R-squared value will be equal to 1. So,  $VIF = 1/(1-1)$  which gives  $VIF = 1/0$  which results in "infinity".

### 6. What is a Q-Q plot? Explain the use and importance of a Q-Q plot in linear regression.

**Answer:** A q-q plot is a plot of the quantiles of the first data set against the quantiles of the second data set. It is used to compare the shapes of distributions. A Q-Q plot is a scatterplot created by plotting two sets of quantiles against one another. If both sets of quantiles came from the same distribution, we should see the points forming a line that's roughly straight.

The q-q plot is used to answer the following questions:

- Do two data sets come from populations with a common distribution?
- Do two data sets have common location and scale?
- Do two data sets have similar distributional shapes?
- Do two data sets have similar tail behaviour?