



SIR PRATEEK JAIN

- . Founder @ Physicsaholics
- . Top Physics Faculty on Unacademy (IIT JEE & NEET)
- . 8+ years of teaching experience in top institutes like FIITJEE (Delhi, Indore) , CP (KOTA) etc.
- . Produced multiple Top ranks.
- . Research work with HC Verma sir at IIT Kanpur
- . Interviewed by International media.

PLUS

ICONIC ^{**}

✓ India's Best Educators

✓ Interactive Live Classes

✓ Structured Courses & PDFs

✓ Live Tests & Quizzes

✗ Personal Coach

✗ Study Planner

24 months

No cost EMI

₹2,333/mo

₹56,000

>

18 months

No cost EMI

₹2,625/mo

₹47,250

>

12 months

No cost EMI

₹3,208/mo

₹38,500

>

6 months

No cost EMI

₹4,667/mo

₹28,000

>

To be paid as a one-time payment

View all plans

 Add a referral code

APPLY

PHYSICSLIVE

PLUS

ICONIC ^{**}

✓ India's Best Educators

✓ Interactive Live Classes

✓ Structured Courses & PDFs

✓ Live Tests & Quizzes

✗ Personal Coach

✗ Study Planner

24 months

No cost EMI

₹2,100/mo

+10% OFF ₹50,400

>

18 months

No cost EMI

₹2,363/mo

+10% OFF ₹42,525

>

12 months

No cost EMI

₹2,888/mo

+10% OFF ₹34,650

>

6 months

No cost EMI


₹4,200/mo

+10% OFF ₹25,200

>

To be paid as a one-time payment

View all plans

 Awesome! PHYSICSLIVE code applied

×

Use code **PHYSICSLIVE** to get 10% OFF on Unacademy PLUS.

For Video Solution of this DPP, Click on below link

Solution on
Website:-

<https://physicsaholics.com/home/courseDetails/41>

Solution on
YouTube:-

<https://youtu.be/IHAly8GLkms>

Physics DPP

DPP-1 Kinematics: Speed, Velocity, Distance and Displacement

By Physicsaholics Team

Q) A Body moves 6 m north. 8 m east and 10m vertically upwards, what is its resultant displacement from initial position:

(a) $10\sqrt{2}$ m

(b) 10 m


(c) $\frac{10}{\sqrt{2}}$ m

(d) 20 m


Join Unacademy PLUS Referral Code :

Physicslive

Ans. a



N 00 ; 10m in Horizontal plane
 & 10 m Vertically upwards



$$d = \sqrt{10^2 + 10^2}$$

$$\boxed{d = 10\sqrt{2} \text{ m}}$$

Q) An athlete completes one round of a circular track of radius R in 40 sec with uniform speed. What will be his displacement at the end of 2 min. 30 sec?

(a) zero

(b) $\sqrt{2}R$

(c) $\frac{5}{2}\pi R$

(d) $\frac{15}{2}\pi R$

Join Unacademy PLUS Referral Code :

Physicslive

Ans. b

Q) A car covers the first half of the distance between two places at 40 kmph and the other half at 60 kmph. The average speed of the car is:

(a) 40 kmph

(b) 48 kmph


(c) 50 kmph

(d) 60 kmph

Join Unacademy PLUS Referral Code :

Physicslive

Ans. b



$$t_1 = \frac{d/2}{40} = \frac{d}{80} \text{ h.}$$

$$t_2 = \frac{d/2}{60} = \frac{d}{120} \text{ h.}$$

$$V = \frac{d}{t} = \frac{d}{t_1 + t_2}$$

$$V = \frac{d}{\frac{d}{80} + \frac{d}{120}} = \frac{1}{\frac{1}{80} + \frac{1}{120}}$$

$$V = \frac{1}{\frac{1}{40}(\frac{1}{2} + \frac{1}{3})} = \frac{40}{\frac{1}{2} + \frac{1}{3}}$$

$$V = \frac{40}{\frac{5}{6}} = 40 \times \frac{6}{5}$$

$$V = 48 \text{ kmph}$$


Q) A particle is constrained to move on a straight line path. It returns to the starting point after 10 sec. The total distance covered by the particle during this time is 30 m. Which of the following statements about the motion of the particle is false?

- (a) Displacement of the particle is zero
- (b) Average speed of the particle is 3 m/s
- (c) Displacement of the particle is 30 m
- (d) Average velocity of the particle is zero.

Join Unacademy PLUS Referral Code :

Physicslive

Ans. c



$$t = 10 \text{ sec.}$$

$$\boxed{\text{Distance} = 30 \text{ m.}}$$

\therefore initial position = final position

$$\therefore \boxed{\text{Displacement} = \text{Zero}}$$

$$\text{Avg. speed} = \frac{\text{Distance}}{\text{time}} = \frac{30 \text{ m}}{10 \text{ sec.}}$$

$$\boxed{\text{Avg speed} = 3 \text{ m/sec}}$$

$$\text{Avg. Velocity} = \frac{\text{Displacement}}{\text{time}} = \frac{0 \text{ m}}{10 \text{ sec.}}$$

$$\boxed{\text{Avg. Velocity} = \text{Zero}}$$


Q) A particle moves along a semicircle of radius 10m from A to B in 5 seconds. The average velocity of the particle is:

(a) $2\pi \text{ m/s}^{-1}$

(b) $4\pi \text{ m/s}^{-1}$

(c) 2 m/s^{-1}


(d) 4 m/s^{-1}



Join Unacademy PLUS Referral Code :

Physicslive

Ans. d



$$d = 2R$$

$$\text{displacement} = 2R$$

$$= 2 \times 10 = 20 \text{ m}$$

$$\text{Time} = 5 \text{ sec}$$

$$\text{Avg. Velocity} = \frac{\text{Displacement}}{\text{Time}}$$

$$= \frac{20}{5}$$

$$\boxed{\text{Avg. Velocity} = 4 \text{ m/s}}$$

Q) A passenger travels along a straight line with velocity V_1 for first half time and with velocity V_2 for next half time, then the mean speed v is given by -

(a) $v = \frac{v_1 + v_2}{2}$

(b) $v = \sqrt{v_1 v_2}$


(c) $v = \sqrt{\frac{v_2}{v_1}}$

(d) $\frac{2}{v} = \frac{1}{v_1} + \frac{1}{v_2}$

Join Unacademy PLUS Referral Code :

Physicslive

Ans. a



$$d_1 = v_1 t/2 \quad d_2 = v_2 t/2$$

$$d = d_1 + d_2$$

$$\text{Avg. speed} = \frac{v_1 t/2 + v_2 t/2}{t/2 + t/2}$$

$$= \frac{v_1 + v_2}{1 + 1}$$

$$\boxed{\text{Avg. Speed} = \frac{v_1 + v_2}{2}}$$

Q) A particle's position as a function of time is described as $y = 2t^2 + 3t + 4$. What is the average velocity of the particle from $t = 0$ to $t = 3$ sec?

(a) 3 m/s

(b) 6 m/s

(c) 9 m/s

(d) 12 m/s

Join Unacademy PLUS Referral Code :

Physicslive

Ans. c

$$x = 2t^2 + 3t + 4$$

$$\text{at } t=0; x = 4 \text{ m,}$$

$$\begin{aligned} \text{at } t=3 \text{ sec, } x &= 2(3)^2 + 3(3) + 4 \\ &= 18 + 9 + 4 \end{aligned}$$

$$x = 31 \text{ m,}$$

$$\begin{aligned} \text{Displacement} &= x_2 - x_1 \\ &= 27 \text{ m,} \end{aligned}$$

$$\text{Avg velocity} = \frac{27}{3} = 9 \text{ m/s}$$

$$\boxed{\text{Avg. Velocity} = 9 \text{ m/s}}$$


Q) Position-time graph of a particle is shown below. What is the average velocity of the particle between the times $t = 0 \text{ s}$ to $t = 12 \text{ s}$?

(a) 1.33 m/s

(b) zero

(c) 12 m/s

(d) -01.33 m/s



Join Unacademy PLUS Referral Code :

Physicslive

Ans. b

$$\text{at } t=0; \quad u_1 = 0$$

$$\text{at } t=12\text{ s}; \quad u_2 = 0$$

$$\therefore \text{displacement} = \text{Zero}$$

$$\therefore \text{Avg. Velocity} = \frac{\text{Zero}}{12}$$

$$\boxed{\text{Avg. Velocity} = 0 \text{ m/s}}$$


Q) Position-time graph of a particle is shown below. What is the average speed of the particle between the times $t = 8 \text{ s}$ to $t = 12 \text{ s}$?

(a) 0.5 m/s

(b) -0.5 m/s

(c) zero

(d) 2 m/s



Join Unacademy PLUS Referral Code :

Physicslive

Ans. a

$$\text{at } t = 8 \text{ sec}; u_1 = 2 \text{ m/s}$$

$$\text{at } t = 12 \text{ sec}; u_2 = 0 \text{ m/s}$$

$$\begin{aligned} \text{distance} &= |u_2 - u_1| \\ &= |0 - 2| = |-2| \end{aligned}$$

$$\text{distance} = 2 \text{ m/s}$$


$$\therefore \text{Avg speed} = \frac{\text{distance}}{\text{time}} = \frac{2 \text{ m}}{4 \text{ sec}}$$

$$\boxed{\text{Avg speed} = 0.5 \text{ m/s}}$$

Q) Velocity-time graph of a particle is shown below. What is the average velocity of the particle between the times $t = 2 \text{ s}$ to $t = 6 \text{ s}$?

- (a) 0.5 m/s
(c) -3.5 m/s


- (b) 3.5 m/s
(d) 5 m/s



Join Unacademy PLUS Referral Code :

Physicslive

Ans. b



from $t = 2 \text{ s}$ to $t = 6 \text{ s}$

displacement $= A_1 + A_2 + A_3$

$$= (2 \times 3) + (2 \times 3) + \frac{1}{2} \times 2 \times 3$$

$$= 6 + 6 + 3$$

$$= 15 \text{ m}$$

$$\text{Avg. Velocity} = \frac{15}{4} = 3.75 \text{ m/s}$$


Q) Velocity-time graph of a particle is shown below. What is the average speed of the particle between the times $t = 0 \text{ s}$ to $t = 10 \text{ s}$?

(a) 3.5 m/s

(b) -3.5 m/s

(c) 3


(d) -3 m/s



Join Unacademy PLUS Referral Code :

Physicslive

Ans. c



$$\text{distance covered} = A_1 + A_2 + A_3 + A_4 + A_5 + |A_6|$$

$$= \left(\frac{1}{2} \times 2 \times 3\right) + (2 \times 3) + (2 \times 3) + \left(\frac{1}{2} \times 2 \times 3\right) + \left(\frac{1}{2} \times 2 \times 6\right) + \left|\left(\frac{1}{2} \times 2 \times 3\right)\right|$$

$$= 3 + 6 + 6 + 3 + 6 + 3$$

$$d = 30 \text{ m}$$


$$\text{Avg. speed} = \frac{d}{t}$$

$$= \frac{30}{10}$$

$$\boxed{\text{Avg. speed} = 3 \text{ m/s}}$$

Q) Velocity-time graph of a particle is shown below. What is the instantaneous velocity of the particle at $t = 5 \text{ s}$?


- (a) 4 m/s (b) -4 m/s
(c) -4.5 m/s (d) 4.5 m/s



Join Unacademy PLUS Referral Code :

Physicslive

Ans. d



$$\text{slope of line AB} = \frac{6-3}{6-4}$$

$$= \frac{3}{2} \text{ m/s}$$

at $t = 5 \text{ sec}$

$$v = 3 \text{ m/s}$$

in 1 sec.

from A \rightarrow B

velocity increases $= \frac{3}{2} \text{ m/s}$

\therefore Velocity at $t = 5 \text{ sec}$

$$v = (v_{\text{at } t=4s}) + \frac{3}{2}$$

$$v = 3 + \frac{3}{2} = 3 + 1.5$$

$$\boxed{v = 4.5 \text{ m/s}}$$

For Video Solution of this DPP, Click on below link

Solution on
Website:-

<https://physicsaholics.com/home/courseDetails/41>

Solution on
YouTube:-

<https://youtu.be/IHAly8GLkms>

Chalo Niklo