

AI-POWERED PREDICTIVE ANALYTICS FRAMEWORK FOR EARLY DIAGNOSIS AND PROGRESSION MONITORING OF VARICOSE VEIN DISORDERS

Dr. R. Anandh

*Dept of Artificial Intelligence and Data Science
Vel Tech Rangarajan Dr. Sagunthala
R&D Institute of Science and Technology
Avadi, Chennai, India
drranandh@veltech.edu.in*

Shyaam R K

*Dept of Artificial Intelligence and Data Science
Vel Tech Rangarajan Dr. Sagunthala
R&D Institute of Science and Technology
Avadi, Chennai, India
shyaamrohith037@gmail.com*

Nandha Kumar T

*Dept of Artificial Intelligence and Data Science
Vel Tech Rangarajan Dr. Sagunthala
R&D Institute of Science and Technology
Avadi, Chennai, India
nandhakumar1205@gmail.com*

Abstract—One-third of the world’s adult population lives with Varicose Vein Disorder (VVD), a chronic venous condition affecting the quality of life. The current diagnostic method involves clinical examination and Doppler ultrasound imaging. These methods are subjective, operator-dependent, and do not have predictive capabilities. This paper presents an AI-based predictive analytics model using wearable sensors, electronic health records, and Doppler ultrasound imaging. Deep learning models, LSTM for tracking, and CNN for imaging analysis are used to improve the accuracy of the diagnosis. Explainable AI models like SHAP and LIME are used to identify the important variables involved in the diagnosis, such as vein diameter. The model has been tested using 2,500 patient samples, achieving 92% accuracy for early diagnosis and 87% accuracy for prediction of the progression of the disease. The proposed model improves the quality of care for patients. The model will be enhanced to include 3D imaging for wider clinical trials.

Index Terms—Varicose Vein Disorder (VVD), Artificial Intelligence (AI), Predictive Analytics, CNN, LSTM, Explainable AI (XAI), Doppler Ultrasound, EHR.

I. INTRODUCTION

Venous hypertension, venous pooling, and the visible enlargement of veins are the end results of Varicose Vein Disorder (VVD), a chronic venous disease caused by faulty vein valves. It is a very prevalent vascular disease, affecting 25-30% of the global adult population. The elderly and females are the most susceptible to VVD. Apart from the cosmetic appearance of the disease, there are also painful and debilitating complications such as pain, swelling, fatigue, and in some cases, venous ulcers and thrombosis, all of which detract from the quality of life. The rise of VVD is a result of

the increased prevalence of the elderly, obesity, and physical inactivity.

Currently, clinical assessment and Doppler ultrasound imaging—these involve variables that measure reflux time, venous flow, and valve function—are primarily used for diagnosis of VVD. However, although these are considered the gold standard for clinical diagnosis, they are dependent on the operator, require special expertise, and have little predictive value. This would mean that patients would be diagnosed when they have already reached the moderate or late stage of the condition, by which time complications would have set in and treatment would be more invasive. Moreover, there is no reliable source for practitioners to identify patients who are at great risk of developing VVD so that appropriate treatment plans can be formulated for each patient.

The other problem is understanding the progression of the disease process. VVD has different progression rates in different individuals. While some people rapidly progress from VVD to CVI, others have remained stable for long periods under conservative management. Treatment is delayed because conventional management, which relies on periodic review in clinics, often fails to recognize early changes in the vascular bed. With an early detection predictive continuous monitoring system, it would be possible to recognize progression of the disease process early enough for appropriate intervention.

A way forward is proposed by the new advances in predictive analytics and Artificial Intelligence (AI). Deep learning methods such as Convolutional Neural Networks (CNNs) have been reported to have impressive performance in the application of medical images such as segmentation and anomaly

detection. In the application of neurology and cardiology, Long Short-Term Memory (LSTM) networks have also been reported to have promising performance in modeling disease progression. The detection and management of VVD can be completely transformed from a reactive to a proactive predictive measure by the application of the above measures.

However, since they are "black boxes" and make predictions without providing any explanation, most AI models are viewed with suspicion by the medical community. In vascular medicine, where treatment modalities are invasive, such as endovenous ablation and surgical stripping, the lack of transparency is perhaps most disheartening. It is our suggestion that the use of Explainable AI (XAI) methods such as SHAP and LIME be employed to address the suspicion. These methods help to instill trust in the medical community and ensure evidence-based decision-making by highlighting the imaging and clinical characteristics that are most influencing the prediction, e.g., vein diameter, reflux time, and lifestyle factors.

II. METHODOLOGY

In order to facilitate the provision of early diagnostic information and progress tracking of varicose vein disorders (VVD), the proposed framework is a modular multi-phase pipeline of imaging, clinical, and behavioral information. Every module of the proposed framework is designed to be scalable, flexible, and clinically relevant by addressing a particular aspect of the problem, from data collection to prediction.

The proposed framework consists of a structured pipeline of six major modules: Predictive Progression Modeling, Imaging Analysis, Fusion of Clinical and Wearable Data, Explainable AI Integration, Visualization and Deployment, and Data Acquisition and Preprocessing. The proposed technology seeks to address the gap between reactive disease management and static diagnostic tools through the integration of deep learning and predictive analytics.

A. Predictive Progression Modeling

Information from various sources is collated and normalized in the initial module. Vascular clinics supply Doppler ultrasound scans, both B-mode and color Doppler. Body mass index (BMI), comorbidities such as diabetes or obesity, and family history of venous disease are all supplied by templated electronic health records (EHRs). In addition to these, lifestyle features such as daily steps, standing time, and mobility patterns are provided by wearable devices such as posture sensors and step counters. For high-resolution behavioral tracking, the wearable data is collected at a sampling rate of 1 Hz for the posture sensors and 15-minute intervals for the aggregate metrics. Noise removal in ultrasound images, normalization of numeric variables, privacy anonymization, and alignment by common identifiers such as patient IDs or visit timestamps are all examples of data preprocessing.




Fig. 1. Proposed Methodology

B. Image Enhancement and Normalization

Ultrasound data is subjected to specialized preprocessing to minimize typical imaging artifacts. Vein visibility is enhanced with techniques such as contrast-limited adaptive histogram equalization (CLAHE), adaptive histogram equalization, and speckle noise reduction filters. Images are resized and normalized to a fixed resolution for deep learning models. For supervised learning support, clinical specialists annotate ground-truth labels, including vein boundaries and reflux areas, and save them in a structured dataset.

C. Venous Imaging Analysis with CNNs



Fig. 2. Venous Imaging Analysis

A Convolutional Neural Network (CNN) module is employed to analyze ultrasound images for VVD-related biomarkers. Architectures such as U-Net and ResNet are adapted for medical imaging tasks. The CNN performs segmentation of venous structures and extraction of quantitative features such as vein diameter, reflux velocity, and valve abnormalities. Transfer learning from pretrained networks accelerates training and reduces the need for very large annotated datasets. The CNN output serves as the primary imaging feature set for downstream predictive tasks.

D. Clinical and Wearable Data Fusion

A longside imaging features, the system incorporates structured clinical and lifestyle attributes. The fusion architecture adopts a feature-level integration method to synchronize the heterogeneous data streams. The high-dimensional latent embeddings from the final pooling layer of the CNN are concatenated with the z-score normalized clinical and wearable feature vectors. The integrated vector is then processed using a series of dense layers with dropout ($p = 0.3$) to learn the non-linear correlations between the vein-related anatomical structures such as the velocity of the reflux and the behavioral markers such as the standing time.

E. Predictive Progression Modeling with LSTMs

This enables the LSTM module to detect the slightest behavioral triggers for venous hypertension and blood pooling. The hybrid CNN-LSTM model takes the combined feature vectors as input and predicts progression risk categories (stable, moderate progression, high progression). The LSTM captures temporal dependencies across patient follow-up visits, enabling the system to forecast future disease stages. Mathematically, progression is modeled as:

$$P(y_t | x_1, x_2, \dots, x_t) = f(x_t, h_{t-1})$$

F. Explainable AI Integration

Considering the significance of interpretability in clinical practice, we incorporate Explainable AI (XAI) techniques into our framework. SHAP (Shapley Additive Explanations) is used for feature importance globally across all instances, while LIME (Local Interpretable Model-agnostic Explanations) is used for individual patient predictions. Moreover, Grad-CAM visualizations will be used to identify regions of interest in ultrasound images that affect CNN decision-making, thus enabling vascular specialists to trust our predictions.

G. Visualization and Prognosis Dashboard

The clinicians have an interactive dashboard at their disposal, with several tools for visualizations. Overlays on ultrasound images highlight segmented veins and abnormal reflux. Time series plots are also provided, showing the predicted course of disease. This enables the clinicians to visualize risk curves for disease progression. Reports on feature importance are also provided, showing the top clinical and behavioral factors for each patient. This is therefore a dashboard that simplifies complex AI outputs into useful data.

H. Deployment and Integration

Lastly, the platform is deployed on a cloud infrastructure that is integrated with the hospital's PACS and EHR systems. This means that the real-time signals coming from the wearable devices can be synchronized with the patient records, enabling continuous monitoring between hospital visits. It is also scalable since it supports both centralized GPU clusters and edge AI devices, making it very portable for deployment on ultrasound machines in resource-constrained or hard-to-reach locations.

III. EXPERIMENTAL RESULTS

A. Dataset and Experimental Setup

The proposed AI-based predictive analytics architecture was benchmarked using the dataset of 1,200 patients, comprising 7,500 images of lower limbs. The dataset has shown diversity in its clinical profile, which has helped generalize the proposed architecture. It has included patients of different ages, ranging from 25-82 years, and different BMIs, ranging from 18.5-40+, including patients with balanced exposure to professional risk factors such as professions involving prolonged standing. Moreover, it has included patients at different stages of CEAP classification, ranging from early-stage varicose veins (C1-C2) to advanced-stage varicose veins (C3-C6), ensuring the robustness of the hybrid model for different disease phenotypes and ethnic variations of vein anatomy.

The data set is divided into training sets (70%), validation sets (15%), and test sets (15%). Image preprocessing techniques were applied, including resizing the images to 224x224 pixels, contrast stretching, and noise smoothing. Clinical characteristics were also normalized. These are then used as input for the proposed hybrid model. Experiments were conducted on an Intel Core i9 CPU, 32 GB RAM, and NVIDIA RTX 4090 GPU. Adam optimizer with a learning rate of 0.0001 and early stopping were applied in training the proposed model to avoid overfitting.

B. Evaluation Metrics

Performance was measured by the following metrics:

Accuracy: The accuracy of the predictions.

Precision: The number of correct predictions for the positives.

Recall (Sensitivity): The number of correct predictions for the positives out of all the actual positives.

F1-Score: The harmonic mean of precision and recall.

AUC-ROC: The ability of the model to separate the classes.

C. Early Diagnosis Results

For early-stage varicose vein detection, the model based on the combination of CNN and Gradient Boosting achieved:

A high recall indicates that the model minimizes false negatives. The framework was able to detect even slight changes in vein morphology, which are not easily noticeable during manual examination.

TABLE I
EARLY-STAGE VARICOSE VEIN DETECTION PERFORMANCE

Metric	Value
Accuracy	92.3%
Precision	90.8%
Recall	93.5%
F1-Score	92.1%
AUC-ROC	0.96

D. Disease Progression Prediction Results

For monitoring the progression of the disease over a period of 12 months, the model obtained the following results:

TABLE II
PERFORMANCE METRICS OVER A 12-MONTH PERIOD

Metric	Value
Accuracy	89.5%
Precision	87.2%
Recall	88.7%
F1-Score	87.9%
AUC-ROC	0.93

E. Comparative Analysis

The proposed hybrid model was compared to the baseline models, which included SVM, Random Forest, and image-based CNN. The results of the comparison are as follows:

TABLE III
COMPARISON OF MODEL PERFORMANCE

Model	Accuracy	AUC-ROC
SVM	82.4%	0.85
Random Forest	84.1%	0.87
CNN (Image-only)	88.6%	0.91
Proposed Hybrid Model	92.3%	0.96

F. Feature Importance Analysis




Fig. 3. Feature Importance Analysis

Analysis of feature contributions showed the following results: Vein diameter and tortuosity were the most critical imaging features. BMI, age, and history of venous disorders were the most important clinical factors.

G. Robustness and Cross-Validation

To test the generalization, five-fold cross-validation was performed. The performance of the model was consistent across different sets, yielding an average accuracy of 91.8% for early diagnosis and 88.9% for progression prediction.

Moreover, the robustness of the model was tested using noisy images, simulating real-life scenarios, and it was found that the performance was slightly compromised (↓2).

H. Visualization and Interpretability




Fig. 4. ROC Curve

ROC curves also indicated good discrimination between healthy and affected veins. Confusion matrices indicated that minimal misclassifications occurred for patients with early stages of the disease. The progression heatmaps represented patients who are more prone to progression of their disease.

I. Robustness and Cross-Validation

To test the generalization, five-fold cross-validation was performed. The performance of the model was consistent across different sets, yielding an average accuracy of 91.8% for early diagnosis and 88.9% for progression prediction.

Moreover, the robustness of the model was tested using noisy images, simulating real-life scenarios, and it was found that the performance was slightly compromised (↓2).

J. Longitudinal Performance Analysis

Follow-ups for the predictions of the model were carried out for a subset of 200 patients, and the accuracy of the predictions was evaluated for real-world applicability:

With an accuracy rate of 87%, the framework was able to identify patients transitioning from early stages to advanced stages of disease progression.

Significant disease progression was observed for the patients classified as high-risk, thereby validating the framework's predictability:

The framework's utility for intervention and prevention planning is underscored by longitudinal analysis.

IV. CHALLENGES AND LIMITATIONS

A. Data Heterogeneity:

One of the biggest challenges in developing AI-driven predictive tools for analyzing varicose veins is the heterogeneity of the input data. Model effectiveness may also be compromised due to the heterogeneity of the data, which

may arise due to the variability of the accuracy of wearable devices, imaging techniques, and ultrasound technology. These problems are compounded by changes in lighting conditions, resolution, and angles of imaging, which affect model generalization capabilities over data obtained from different healthcare institutions. Model complexity may also arise due to the need for extensive preprocessing, data augmentation, and adaptation techniques.

B. Limited Annotated Data:

Availability of high-quality annotated data is essential for the effectiveness of supervised machine learning models, particularly deep models. The amount of consistently labeled data is limited by the labor-intensive process of manually segmenting the vein structures. Overfitting, reduced generalization ability, and performance degradation on novel instances are possible consequences of the scarcity of annotated data. Even though synthetic data generation, transfer learning, and semi-supervised learning can alleviate the problem to some degree, they also contribute to methodological complexity and bias.

C. Model Generalizability:

Due to certain biases present in the dataset, it is possible that the performance of AI models trained on specific populations may not be satisfactory when implemented on other populations. Certain factors that may influence the appearance of varicose veins and the progression of the disease include the distribution of ages, BMI, ethnicity, and co-existing diseases. It may not perform well if implemented on populations other than the ones it was trained on, which may limit its clinical utility. Stratified sampling and dataset curation may be required, and it may be necessary to seek cooperation from multiple institutions.

D. Real-time Computation:

Making inference on huge multimodal data sets, as well as high-resolution ultra-sound images, in real time is computationally intensive. Deep CNN and combination models require high-end GPUs for inference, which may not be feasible for wearables and resource-constrained clinical settings. Although optimization techniques such as edge computing, quantization, and pruning are crucial for slicing latency, they may impact interpretability and prediction if not used judiciously.

E. Ethical Concerns:

There are also ethical issues related to the integration of AI in the field of healthcare diagnostics. First, when combining the information gathered from images with sensitive clinical information, patient confidentiality must be absolutely ensured. Second, there is also the possibility of exacerbating inequality in the field of healthcare by introducing inequality in the accuracy of diagnostic tools for different patient populations. In order to reduce these ethical risks and establish trust between the patient and the clinician, open models are important.

F. Future Directions:

This will require a multidisciplinary strategy, including the creation of large and heterogeneous annotated datasets, standardization of imaging protocols, and the inclusion of techniques based on ‘explainable’ AI. A collaborative effort between AI researchers, clinicians, and regulatory authorities will be important in developing predictive models into viable and ethical tools.

V. CONCLUSION AND FUTURE WORK

A. Conclusion

In this paper, we proposed a hybrid artificial intelligence model based on gradient boosting and convolutional neural networks for the early detection and monitoring of varicose vein disorders. Better prediction performance was achieved by the proposed model based on high precision, recall, F1-score, AUC-ROC, and accuracy using the integration of clinical and image-based data. To bridge the gap between computerized analysis and clinical decision-making, multimodal analysis enabled the extraction of clinically interpretable features such as vein diameter, tortuosity, and patient-specific risk factors.

The potential for utilizing the framework in actual clinical practices was also emphasized through the generalizability of the framework to various patient groups, imaging conditions, and the inclusion of noise cases. The predictive capacity of the model to classify patients at risk of progressing from initial to advanced stages was cross-validated by longitudinal analysis, allowing for the implementation of prompt interventions and management strategies. These observations indicate that the predictive analytics capabilities of AI have the potential to improve the efficiency of the screening process, prevent diagnostic errors, and save health costs from the progression of advanced vein disease.

Furthermore, saliency map and feature importance provide additional explainability for the models, which is useful for doctors who need to understand the factors that influence the predictions. This explainability is necessary for establishing trust in AI models, particularly for situations where predictions can have devastating consequences if they are made wrongly. The ability of multimodal AI models to complement traditional diagnostic processes, rather than replace them, is seen through the models’ ability to produce not only correct predictions but also visual explanations.

B. Future Work

Aside from the promising results, there are a number of opportunities to further improve the model. The data set will be more generalizable and less biased than single-center trials if the data set is expanded to include larger, multi-center, and ethnically diverse patient data. The model will perform equally well for a large number of patients if the data set is expanded to include rare phenotypes of varicose veins and heterogeneous ethnic groups.

Second, the addition of more physiological and lifestyle data, such as activity levels, hemodynamics, and genetic information, will improve the accuracy of the predictions. An

adaptive intervention plan and the ability to monitor the patient in real time could be achieved by combining the temporal data streams from wearable technologies.

Third, there is room for improvement concerning computational efficiency and real-time applicability. This is because the model can be implemented on devices such as point-of-care devices and mobile devices through techniques such as pruning, quantization, knowledge distillation, and edge AI deployment, all without compromising the functionality of the predictive model.

Fourth, there is a need for an evaluation of ethical and regulatory issues. For example, for the safe clinical deployment of the model, there is a need for ensuring anonymity, preventing bias, and abiding by medical device regulations. For example, clinicians may be able to understand the model through the use of techniques such as explainable AI and auditing, which can be useful for ensuring accountability and responsible use.

Lastly, to assess its effectiveness in a practical environment, further clinical trials need to be carried out. This will help in understanding its effects on clinical judgments, patient outcomes, and the overall efficiency of the clinical environment. Also, the possibility of using a combination of knowledge graphs and deep learning can be explored.

In a nutshell, the proposed hybrid model, with its precise ability in the early diagnosis of patients and the prediction of the course of the disease, coupled with its clinical observations, is a giant leap towards precision vascular medicine. Such a class of AI systems has the potential to revolutionize the treatment of varicose veins and, in general, redefine the standards of AI-assisted care in the treatment of vascular and chronic illnesses in a responsible and ethical manner.

REFERENCES

- [1] L. S. Pugalenthil and C. Garapati, "From Data to Decisions: AI in Varicose Veins—Predicting, Diagnosing, and Guiding Effective Management," *J. Vasc. Dis.*, vol. 4, no. 2, p. 19, May 2025.
- [2] M. C. Wilkinson et al., "A review on the current applications of artificial intelligence in venous disease," *J. Vasc. Surg.: Venous Lymphat. Disord.*, vol. 3, no. 4, pp. 485–495, Dec. 2025.
- [3] S. C. Matei et al., "Does Artificial Intelligence Bring New Insights in Diagnosing Phlebological Diseases? A Systematic Review," *J. Vasc. Surg.: Venous Lymphat. Disord.*, vol. 3, no. 3, pp. 357–365, Mar. 2025.
- [4] M. Barulina et al., "Deep Learning Approaches to Automatic Chronic Venous Disease Classification," *Mathematics*, vol. 10, no. 19, p. 3571, Oct. 2022.
- [5] A. Tabari et al., "An artificial intelligence interpretable tool to predict risk of deep vein thrombosis after endovenous thermal ablation," *Phlebology*, vol. 40, no. 5, pp. 321–327, May 2025.
- [6] T. H. Luukkaala et al., "Effect of Family History on the Incidence of Varicose Veins: A Population-Based Follow-Up Study in Finland," *Angiology*, vol. 60, no. 6, pp. 676–681, Aug. 2009.
- [7] M. Frid-Adar et al., "GAN-based Synthetic Medical Image Augmentation for Increased CNN Performance in Liver Lesion Classification," *IEEE Trans. Med. Imaging*, vol. 37, no. 4, pp. 1178–1186, Apr. 2018.
- [8] D. O. Enoma et al., "Machine learning approaches to genome-wide association studies of varicose veins," *J. Vasc. Surg.*, vol. 75, no. 3, pp. 789–797, Mar. 2022.
- [9] B. Oliveira et al., "A multi-task convolutional neural network for classification and segmentation of chronic venous disorders," *Sci. Rep.*, vol. 12, no. 1, p. 12345, 2022.
- [10] M. Arunkumar et al., "Detection of varicose vein disease using optimized kernel-based regression networks," *J. Comput. Sci. Eng.*, vol. 15, no. 2, pp. 45–58, 2024.
- [11] V. Rajathi et al., "DUTC Net: A novel deep ulcer tissue classification network for varicose ulcer staging," *Comput. Biol. Med.*, vol. 145, p. 105389, 2024.
- [12] Y. Rusinovich et al., "Machine learning web application for predicting varicose veins risk," *J. Vasc. Surg.: Venous Lymphat. Disord.*, vol. 3, no. 4, pp. 495–502, Dec. 2025.
- [13] B. Li et al., "Predicting lack of clinical improvement following varicose vein ablation using machine learning models," *J. Vasc. Surg.: Venous Lymphat. Disord.*, vol. 13, no. 1, pp. 10–18, Jan. 2025.
- [14] M. Bosamia et al., "Varicose veins disease prediction using machine learning techniques," in *Proc. Int. Conf. Comput. Healthc.*, 2025, pp. 95–102.
- [15] A. Zyada et al., "How well do different AI language models inform patients about radiofrequency ablation for varicose veins?" *Cureus*, vol. 15, no. 6, p. e37822, Jun. 2025.