

Table II:

	sandwich	brickwork
excitation energy E_a	$2.522 \pm 0.014 \text{ eV}$	$2.565 \pm 0.010 \text{ eV}$
excitation energy E_b	$2.523 \pm 0.014 \text{ eV}$	$2.561 \pm 0.010 \text{ eV}$
excitonic coupling V_{exc}	$0.25 \pm 0.017 \text{ eV}$	$-0.064 \pm 0.008 \text{ eV}$
center-center distance R_{ab}	$4.25 \pm 0.08 \text{ \AA}$	$8.33 \pm 0.24 \text{ \AA}$
angle between long axes	$6.0 \pm 1.7^\circ$	$34.0 \pm 4.0^\circ$
angle between short axes	$174.0 \pm 2.6^\circ$	$122.0 \pm 3.9^\circ$
distance of charge centers R_{cc}	$5.42 \text{ \AA}$	$7.57 \text{ \AA}$

V. FIGURE CAPTIONS:

Figure 1:

The atom numbering for PIC is shown as it is used to tabulate the parameters

Figure 2:

The experimental absorption spectrum⁴ of monomeric PIC (dots) is compared with a calculated spectrum (full line) from the displaced oscillator model. The calculated spectrum was shifted to reproduce the absorption maximum at 19100 cm^{-1}

Figure 3:

Starting from a sandwich (a) or brickwork (b) structure relative stable dimer configurations evolved. The figure shows representative snapshots.