



FIGURE 9. Γ_5 FIGURE 10. Γ_6

- (3) $\sigma_1(\vec{e}_i) = \vec{e}_i$ and $\sigma_1(\vec{f}_j) = \vec{f}_j$; $1 \leq i \leq 3$ and $1 \leq j \leq 7$.
 (4) $\sigma_0 = C_1 \dots C_5$, where

$$C_1 = (\vec{e}_1, \vec{f}_1, \vec{e}_3, \vec{f}_1),$$

$$C_2 = (\vec{e}_2, \vec{f}_3, \vec{e}_1, \vec{f}_2),$$

$$C_3 = (\vec{e}_3, \vec{f}_6, \vec{e}_2, \vec{f}_5),$$

$$C_4 = (\vec{f}_2, \vec{f}_4, \vec{f}_3, \vec{f}_4),$$

$$C_5 = (\vec{f}_5, \vec{f}_7, \vec{f}_6, \vec{f}_7).$$

Now, we count the boundary components of Γ_5 using the permutation σ_∞ . The boundary components of the graph Γ_5 given by:

$$\partial_1 = \vec{e}_1 \vec{f}_2 \vec{f}_4 \vec{f}_2 \vec{e}_2 \vec{f}_5 \vec{f}_7 \vec{f}_5 \vec{e}_3 \vec{f}_1 \vec{e}_3 \vec{f}_6 \vec{f}_7 \vec{f}_6 \vec{e}_2 \vec{f}_3 \vec{f}_4 \vec{f}_3 \vec{e}_1 \vec{f}_1.$$

So, there is only one boundary component.

4. UPPER BOUND ON MAPPING CLASS ORBITS OF MAXIMUM SIZE FILLING

Let $I(S_g)$ represent the set of all isotopy classes of simple closed curves on S_g . $I(S_g)^n$ denotes the Cartesian product of $n \in \mathbb{N}$ copies of $I(S_g)$. Define an equivalence relation $\sim$ on $I(S_g)^n$ by