

- Drahl, C., Cravatt, B.F., and Sorensen, E.J. (2005). Protein-reactive natural products. *Angew. Chem. Int. Ed.* **44**, 5788–5809.
- Heyninck, K., Lahtela-Kakkonen, M., Van der Veken, P., Haegeman, G., and Vanden Berghe, W. (2014). Withaferin A inhibits NF-kappaB activation by targeting cysteine 179 in IKK β . *Biochem. Pharmacol.* **91**, 501–509.
- Heyninck, K., Sabbe, L., Chirumamilla, C.S., Szarc Vel Szic, K., Vander Veken, P., Lemmens, K.J.A., Lahtela-Kakkonen, M., Naulaerts, S., Op de Beeck, K., Laukens, K., et al. (2016). Withaferin A induces heme oxygenase (HO-1) expression in endothelial cells via activation of the Keap1/Nrf2 pathway. *Biochem. Pharmacol.* **109**, 48–61.
- Jessani, N., Humphrey, M., McDonald, W.H., Niessen, S., Masuda, K., Gangadharan, B., Yates, J.R., Mueller, B.M., and Cravatt, B.F. (2004). Carcinoma and stromal enzyme activity profiles associated with breast tumor growth in vivo. *Proc. Natl. Acad. Sci. USA* **101**, 13756–13761.
- Jessani, N., Niessen, S., Mueller, B.M., and Cravatt, B.F. (2005). Breast cancer cell lines grown in vivo: what goes in isn't always the same as what comes out. *Cell Cycle* **4**, 253–255.
- Lee, I.-C., and Choi, B.Y. (2016). Withaferin-A—a natural anticancer agent with pleiotropic mechanisms of action. *Int. J. Mol. Sci.* **17**, 290.
- Lee, J., Liu, J., Feng, X., Salazar Hernández, M.A., Mucka, P., Ibi, D., Choi, J.W., and Ozcan, U. (2016). Withaferin A is a leptin sensitizer with strong anti-diabetic properties in mice. *Nat. Med.* **22**, 1023–1032.
- Louie, S.M., Grossman, E.A., Crawford, L.A., Ding, L., Camarda, R., Huffman, T.R., Miyamoto, D.K., Goga, A., Weerapana, E., and Nomura, D.K. (2016). GSTP1 is a driver of triple-negative breast cancer cell metabolism and pathogenicity. *Cell Chem. Biol.* **23**, 567–578.
- M, M., Zameer, F., Naidu, A., M N, N.P., Dhananjaya, B.L., and Hegdekatte, R. (2016). Evaluating the inhibitory potential of *Withania somnifera* on platelet aggregation and inflammation enzymes: an in vitro and in silico study. *Pharm. Biol.* **54**, 1936–1941.
- Medina-Cleghorn, D., Bateman, L.A., Ford, B., Heslin, A., Fisher, K.J., Dalvie, E.D., and Nomura, D.K. (2015). Mapping proteome-wide targets of environmental chemicals using reactivity-based chemoproteomic platforms. *Chem. Biol.* **22**, 1394–1405.
- Nomura, D.K., Long, J.Z., Niessen, S., Hoover, H.S., Ng, S.-W., and Cravatt, B.F. (2010). Monoacylglycerol lipase regulates a fatty acid network that promotes cancer pathogenesis. *Cell* **140**, 49–61.
- Roberts, A.M., Ward, C.C., and Nomura, D.K. (2016). Activity-based protein profiling for mapping and pharmacologically interrogating proteome-wide ligandable hotspots. *Curr. Opin. Biotechnol.* **43**, 25–33.
- Roberts, A.M., Miyamoto, D.K., Huffman, T.R., Bateman, L.A., Ives, A.N., Akopian, D., Heslin, M.J., Contreras, C.M., Rape, M., Skibola, C.F., et al. (2017a). Chemoproteomic screening of covalent ligands reveals UBA5 as a novel pancreatic cancer target. *ACS Chem. Biol.* **12**, 899–904.
- Roberts, L.S., Yan, P., Bateman, L.A., and Nomura, D.K. (2017b). Mapping novel metabolic nodes targeted by anti-cancer drugs that impair triple-negative breast cancer pathogenicity. *ACS Chem. Biol.* **12**, 1133–1140.
- Sangodkar, J., Farrington, C.C., McClinch, K., Galsky, M.D., Kastrinsky, D.B., and Narla, G. (2016). All roads lead to PP2A: exploiting the therapeutic potential of this phosphatase. *FEBS J.* **283**, 1004–1024.
- Tao, S., Tillotson, J., Wijeratne, E.M.K., Xu, Y., Kang, M., Wu, T., Lau, E.C., Mesa, C., Mason, D.J., Brown, R.V., et al. (2015). Withaferin A analogs that target the AAA+ chaperone p97. *ACS Chem. Biol.* **10**, 1916–1924.
- Wang, C., Weerapana, E., Blewett, M.M., and Cravatt, B.F. (2014). A chemoproteomic platform to quantitatively map targets of lipid-derived electrophiles. *Nat. Methods* **11**, 79–85.
- Weerapana, E., Wang, C., Simon, G.M., Richter, F., Khare, S., Dillon, M.B.D., Bachovchin, D.A., Mowen, K., Baker, D., and Cravatt, B.F. (2010). Quantitative reactivity profiling predicts functional cysteines in proteomes. *Nature* **468**, 790–795.
- Xu, T., Park, S.K., Venable, J.D., Wohlschlegel, J.A., Diedrich, J.K., Cociorva, D., Lu, B., Liao, L., Hewel, J., Han, X., et al. (2015). ProLuCID: an improved SEQUEST-like algorithm with enhanced sensitivity and specificity. *J. Proteomics* **129**, 16–24.
- Yu, Y., Hamza, A., Zhang, T., Gu, M., Zou, P., Newman, B., Li, Y., Gunatilaka, A.A.L., Zhan, C.-G., and Sun, D. (2010). Withaferin A targets heat shock protein 90 in pancreatic cancer cells. *Biochem. Pharmacol.* **79**, 542–551.
- Zhou, Y., Li, W., Wang, M., Zhang, X., Zhang, H., Tong, X., and Xiao, Y. (2016). Competitive profiling of celastrol targets in human cervical cancer HeLa cells via quantitative chemical proteomics. *Mol. Biosyst.* **13**, 83–91.