


Table 4 Fumigant and contact toxicity of essential oil of *A. stolonifera* aerial parts and its main components against *L. serricorne* adults.

Treatment	Fumigant toxicity			Contact toxicity		
	LC ₅₀ ^a (μg/mL air)	95% FL ^c	Chi square (χ ²)	LD ₅₀ ^b (μg/adult)	95% FL ^c	Chi square (χ ²)
Essential oil	0.99	1.58- 3.37	9.38	12.68	11.14-14.16	14.16
Eucalyptol*	5.18	4.63- 5.70	16.79	15.58	12.88-18.02	15.18
β-Pinene	29.03	26.38-31.79	17.48	65.55	58.13-76.09	21.62
Camphor	2.91	2.57- 3.26	13.11	11.30	7.78-14.07	16.13
Terpinen-4-ol	1.30	0.75- 1.70	12.44	5.42	3.99- 6.57	11.63
Phosphine**	9.23 × 10 ⁻³	7.13 × 10 ⁻³ -11.37 × 10 ⁻³	11.96	—	—	—
Pyrethrins**	—	—	—	0.24	0.16- 0.35	17.36

*Data from Zhang *et al.*¹⁾; ** data from Yang *et al.*³¹⁾; ^a 50% of lethal concentration; ^b 50% of lethal dose; ^c fiducial limits.

**Fig. 2** Percentage repellency (PR) of the essential oil from *A. stolonifera* aerial parts and its constituents against *T. castaneum* at 2 h (A) and 4 h (B) after exposure^a.

^ameans in the same column followed by the same letters do not differ significantly ($p > 0.05$) in ANOVA and Tukey's tests. PR was subjected to an arcsine square-root transformation before ANOVA and Tukey's tests.