

Table 1. Number of species and zoogeographical endemisms in the nine zoological regions (endemisms are the ratios of endemics to total species)

	All classes	Amphibian	Bird	Mammal	Reptile
Totals					
Madagascar	853	143	240	136	334
Hawaii	87	0	86	1	0
New Zealand	122	3	82	2	35
Australasia	4015	526	1610	674	1205
Sub-Sahara	5043	629	2046	1038	1330
Indo-Malaysia	4572	710	1780	830	1252
Neotropics	9259	2168	3687	1229	2175
Palearctic	3638	376	1568	905	789
Nearctic	1950	267	728	481	474
Endemics					
Madagascar	709	140	122	126	321
Hawaii	36	0	36	0	0
New Zealand	80	3	40	2	35
Australasia	3532	515	1285	607	1125
Sub-Sahara	4277	618	1536	919	1204
Indo-Malaysia	2497	565	555	438	939
Neotropics	8127	2092	3110	1023	1902
Palearctic	1353	231	211	471	440
Nearctic	758	192	95	258	213
Ratios					
Madagascar	0.83	0.98	0.51	0.93	0.96
Hawaii	0.41	–	0.42	0.00	–
New Zealand	0.66	1.00	0.49	1.00	1.00
Australasia	0.88	0.98	0.80	0.90	0.93
Sub-Sahara	0.85	0.98	0.75	0.89	0.91
Indo-Malaysia	0.55	0.80	0.31	0.53	0.75
Neotropics	0.88	0.96	0.84	0.83	0.87
Palearctic	0.37	0.61	0.13	0.52	0.56
Nearctic	0.39	0.72	0.13	0.54	0.45

occurs in more than one zoological region, we count it as present in each region in which it lives.

Statistical methods

We transformed the variables S and AE to common logarithms. Then we performed univariate and bivariate linear regressions and calculated the usual statistics: intercepts, coefficients, probabilities, and R^2 -values. We excluded Hawaii from the regression analyses for Amphibia and Reptilia because these classes are not found in that zoological region.