

Table 4. Intestinal microbiota test results of the control group (CG).

Microbiota	Mean (log DNA copies/g) ± SD			Δ Average %		p Value*	
	Pre-Test (D-0)	Post Test I (D-35)	Post Test II (D-42)	Day 0 and 35	Day 35 and 42	Day 0 and 35	Day 35 and 42
<i>Bifidobacterium</i>	4.79±0.37	4.81±0.31	4.77±0.51	↑0.5%	↓0.8%	0.8	0.84
<i>Bacteroides</i>	0.00±0.00	3.03±1.50	2.45±0.44	↑3.0%	↓19%	0.043	0.345
<i>Clostridium</i>	6.73±0.46	6.59±0.60	6.60±0.56	↓2%	↑0.1%	0.432	0.822
<i>Lactobacillus</i>	6.79±0.54	6.83±0.57	6.85±0.30	↑0.6%	↑0.3%	0.884	0.914
<i>E. coli</i>	5.48±1.22	5.63±2.30	5.48±1.97	↑2.8%	↓2.7%	0.774	0.38

CG: Control Group – standard feed only

*p<0.05 significantly different.

Table 5. Intestinal microbiota statistics test results treatment group I.

Microbiota	Mean (log DNA copies/g) ± SD			Δ Average %		p Value*	
	Pre-Test (D-0)	Post Test I (D-35)	Post Test II (D-42)	Day 0 and 35	Day 35 and 42	Day 0 and 35	Day 35 and 42
<i>Bifidobacterium</i>	5.36±0.53	4.15±0.22	4.20±0.27	↓22.5%	↑1.2%	0.003	0.665
<i>Bacteroides</i>	0.53±0.73	0.00±0.00	0.00±0.00	↓100%	0%	0.180	1.000
<i>Clostridium</i>	6.17±0.81	7.40±0.32	7.74±0.17	↑19.9%	↑4.6%	0.013	0.095
<i>Lactobacillus</i>	6.89±0.71	7.83±0.25	7.64±0.54	↑13.7%	↓2.5%	0.034	0.480
<i>E. coli</i>	3.71±1.24	6.91±2.56	6.54±2.70	↑86.4%	↓5.4%	0.043	0.138

TG I: Treatment Group I — standard feed and HFD intervention

*p<0.05 significantly different

Table 6. Statistical test results of intestinal microbiota treatment group II.

Microbiota	Mean (log DNA copies/g) ± SD			Δ Average %		p Value*	
	Pre-Test (D-0)	Post Test I (D-35)	Post Test II (D-42)	Day 0 and 35	Day 35 and 42	Day 0 and 35	Day 35 and 42
<i>Bifidobacterium</i>	5.07±0.33	4.25±0.31	5.99±0.98	↓16.2%	↑41%	0.011	0.013
<i>Bacteroides</i>	1.19±1.54	0.00±0.00	1.41±0.90	↓100%	↑100%	0.068	0.043
<i>Clostridium</i>	6.16±0.81	8.03±0.71	7.51±0.66	↑30.4%	↓6.3%	0.012	0.036
<i>Lactobacillus</i>	7.74±0.72	8.23±0.61	8.86±0.40	↑6.3%	↑7.6%	0.006	0.043
<i>E. coli</i>	3.64±1.10	7.22±2.42	5.60±2.32	↑98.4	↓22.5%	0.043	0.043

TG II: Treatment Group II — standard feed, HFD, and ADE intervention.

*p<0.05 significantly different




Figure 1. Microbiota profile of all groups.