

Table 4. Clinical Trials of Pre-Exposure Prophylaxis

Trial	Study Population	Location	Intervention	Outcome	Comments
TDF2	1,219 sexually active adults; 55% male, 45% female; 94% unmarried; approximately 90% aged 21–29	Botswana	Daily oral TDF/FTC	63% protection	>30% did not complete study; cannot draw definitive conclusions for women and men separately.
PIP	4,758 serodiscordant heterosexual couples; 38% HIV-negative female, 68% HIV-negative male partner; 98% married; median age 33	Botswana, Kenya, Rwanda, South Africa, Tanzania, Uganda, Zambia	Daily oral TDF or TDF/FTC	67% protection with TDF alone; 75% protection with TDF/FTC	Serodiscordant couples may be a distinct, unique population.
FEM-PrEP	1,951 heterosexual women aged 18–35 at high risk of infection	Kenya, South Africa, Tanzania	Daily oral TDF/FTC	Trial discontinued for futility in April 2011.	Adherence assessment with monthly clinical samples to measure drug concentration is pending.
VOICE MTN-003	5,029 heterosexual women aged 18–45 in high-prevalence areas	Uganda, South Africa, Zimbabwe	Daily oral TDF or daily oral TDF/FTC or daily topical TFV gel	No study drug significantly reduced the risk of HIV acquisition. Estimates of effectiveness were less than 0 for TDF and TDF/FTC daily oral dosing (negative 48.8% and negative 4.2% TDF/FTC respectively), and reduced risk of HIV infection of 14.7% for TDF gel.	Adherence to study drugs was low; TFV was detected in 30% of the oral TDF arm, 29% in the oral TDF/FTC arm, and 25% in the TDF gel arm.

Key to Acronyms: FTC = emtricitabine; TDF = tenofovir disoproxil fumarate; TFV = tenofovir

Source: Adapted from: Kashuba et al., Pre-exposure prophylaxis for HIV prevention: how to predict success: Table Antiretroviral-based HIV prevention studies. *Lancet*. 2012;379(9835): 2409-2411.