


Partenaire de vos innovations

Use of inedible feed in the pork industry, feed efficiency: sustainability metrics, calculating tools, including C footprint

Animal Task Force, Brussels – 18 November 2021


Sandrine ESPAGNOL

Expert in environmental assessment of pig farms






Introduction




The issues: produce food with less environmental impacts






Efficiency at animal scale









Institut du porc


Efficiency at life cycle assessment level

change

phication



Institut du porc



cation


occupation

consumption

Efficiency at life cycle assessment level



Land use by the French pig production



1 million of hectares (// 3,5 % utilised agricultural land)

Today

Free hectares to produce something else


122,000 additionnal ha

20 years ago



Protein efficiency






The competition only concerns the human-edible fraction that is consumed by animals!



Net protein efficiency



Interpretation:






Net protein efficiency



Effect of pig
feeding on
protein
apparent
and net protein
efficiency






Net protein efficiency



1. Human-edible fraction of protein in animal feed ingredients


	Protein	
Pasture	0	
Maize Silage	10	
Maize grain	15	
Wheat grain	66	
Peas grais	74	
Rapeseed grain	eed grain 0	
Wheat co-products	90	
Soybean meal 60		
Rapeseed, sunflower meal 0		
Other co-products	0	

2. Proteins from a pig





Complementarity of efficiencies at different levels





24/11/2021



- Assessment of environmental performances of pig units
 - Since 2014
 - References
 - Benchmarking
- Deployment
 - 611 farmers
 - 749 environmental assessments
 - 115 advisors trained to support farmers
- Tool in the case of Label Bas
 Carbone in France

Indicators		
Natural ressources consumption	Water	Consumed water (I/kg of growth)
	Energy	Nonrenewable energy consumption (kWh/kg of growth)
Manure	Nitrogen	N excreted (g N/kg of growth) N at pig unit gate (g N/kg of growth)
	Phosphorous	Phosphorous excreted (g P ₂ O ₅ /kg of growth)
Gaseous emissions	NH ₃	Direct emissions of ammonia (g NH ₃ /kg of growth)
	GES	Emissions of greenhouse gazes (kg eq CO ₂ /kg live weight pig at farm gate)
Waste production	Waste	Waste produced (g waste/kg of growth)



Optimization of the pig system

- Efficiency and that's all?
 - Central in the environmental assessment of pig production
 - Different from cattle production where an extensification could be associated to more carbon storage
- In pig production the intensification is often associated to more efficiency
 - No correlation with more impacts per ha because in regulation the surface of spreading is adapted to the size of the pig unit
- The limit in pig production comes more from the animal welfare



■ Take-home messages

- Efficiency is good for the environment
- New challenges for the future
 - Diversified stakes to consider (welfare, environment, quality of life, profitability)
 - Changing context: Adaptation to climate change, new farmers, new consumers
 - Searching for compromise: multicriteria optimization
- Place and relevance of having a diversity of livestock systems (some based on industrial ecology, others based on agroecology)
- Choice should be made at macroscopic level of territorial agricultural systems linked to food systems
 - Global environmental results
 - Ability to feed people





Thanks for your attention

<u>sandrine.espagnol@ifip.asso.fr</u> +(33) 07 62 53 63 74



The case of Label Bas Carbone



GOALS

Helping to invest in projects reducing the carbon impact

Offset companies' polluting emissions by financing low-carbon projects

Creation of a virtuous circular economy on French territory

