

Modelling salmon spawning surveys as arrivals and residence, an advance from area under the curve estimates.

Scott A. Akenhead^{a,1}, Patrick L. Thompson^{b,*}

^a*Pacific Biological Station, Fisheries and Oceans Canada, 3190 Hammond Bay Road, Nanaimo, V9T 6N7*

^b*Pacific Science Enterprise Center, Fisheries and Oceans Canada, 4160 Marine Dr, West Vancouver, V7V 1H2*

Abstract

The abundance of spawning salmon is a fundamental variable for salmon management, but current methods provide estimates without uncertainty and biased from assumptions of residence time (to correct for observing a fish in repeated surveys). We used a new model – coded as a multi-level, non-linear, Bayesian regression – to reveal annual abundance, the timing and spread of entry to the spawning grounds, and spawner residence time. The model was challenged with simulated data to demonstrate robustness, and challenged with data from 291 surveys (25 years) of wild Sockeye Salmon (*Oncorhynchus nerka*) spawning in the Okanagan River upstream of Osoyoos Lake. Abundance estimates appear to be more accurate than conventional. While entry timing and spread varied considerably between years, residence was relatively constant. We also discovered that exit spread varies by year. We foresee this model including more ecology, such as effects from spawner biometrics and habitat variables.

Keywords: salmon, *Oncorhynchus*, spawner, spawner survey, Osoyoos Lake, Okanagan River, Columbia River

1. Introduction

1.1. The importance of spawner surveys

Spawner abundance is a key variable in salmon ecology. Functions relating spawners to their offspring that return as adults (“stock” and “recruits”) are

*Corresponding author

Email addresses: scott@s4s.com (Scott A. Akenhead),
patrick.thompson@dfo-mpo.gc.ca (Patrick L. Thompson)

¹(retired).

fundamental for management of salmon fisheries and salmon habitats (Ricker, 1975; Hilborn and Walters, 1992).

Estimates of the number of Sockeye Salmon (*Oncorhynchus nerka* Walbaum 1792) spawners in a river or a lake beach involve repeated surveys during the spawning period. Methods vary; conventionally: visual counts while boating, snorkeling, walking, or flying in a helicopter or small airplane; and recently, analyzing video from drones. There are many other methods to estimate spawner abundance: mark-recapture, conductivity sensors, conductivity sensors, counting fences, eDNA, and combinations of methods. Complications include distinguishing hatchery fish from wild, and separating anadromous Sockeye from non-anadromous Kokanee (sub-species of *O. nerka*).

1.2. AUC: area under the curve

The conventional analysis of a survey of sockeye spawners determines **AUC**, the integral of spawner count over a number of surveys (units: fish $\times$ days). Spawners have a **residence** time on spawning grounds (aka, survey life), so a spawner will be observed more than once in a series of surveys. The estimate for abundance of spawners divides AUC by residence to correct for repeated observations. That estimate of abundance is then proportional to 1/residence. Using an assumed value for residence time changes the absolute number of spawners in each survey to an index of relative abundance Askey et al. (2023). There are two approaches to estimating AUC:

1. **non-parametric**: where “curve” means a trapezoid of linear interpolations of counts between survey dates [English et al. (1992); Irvine et al. (1992);]. Prior knowledge about first and last presence of spawners (assumed dates for bounding zeros) is required. This is a robust estimator (ref?) and efforts have been made to estimate the precision of AUC from precision of observations (Askey et al., 2023; Korman et al., 2002; Millar et al., 2012; Parken et al., 2003; Parsons and Skalski, 2010). Essentially, AUC determines the maximum of a cumulative probability distribution of spawner number estimated by linear segments.
2. **parametric**: a probability density distribution, typically Gaussian, is fitted to the time series of survey estimates in a year Hilborn et al. (1999). This has been extended by multi-level regression to analyze multiple years simultaneously (Adkison and Su, 2001; Labelle et al., 2021; Su et al., 2001). Then AUC is the area under that curve. Additional parameters and different distributions can provide truncated, skewed, or flattened variants.

Both require an assumed value for residence to estimate spawner abundance. Estimates for residence are determined independently of spawner surveys, using methods such as a spawner fence or recapture of fish (after death) tagged upon entry (Askey et al., 2023; Berghe and Gross, 1986; Cormack, 1964; Fukushima and Smoker, 1997; Korman et al., 2002; Lady and Skalski, 1998; Liao, 1994;

Manske and Schwarz, 2000; Neilson and Geen, 1981; Perrin and Irvine, 1990; Semenchenko, 1993; Zaporozhets and Zaporozhets, 2021). also Bue et al. (1998) and English et al. (1992) and Thomason and Jones (1984).

1.3. Objectives

An advance from AUC by a new model will estimate **residence** directly, as well as **timing** (mean day) and **spread** (standard deviation) of spawner entry. This will improve accuracy of **run** estimates (spawner abundance) and yield precisions for those estimates. Analyzing multiple years of surveys simultaneously will provide annual estimates of parameters, improved by using groups in a multilevel regression.

2. Background

Our analysis was confined to the population of wild Sockeye Salmon that spawn in October and November in the Okanagan River (Figure 1) upstream of Osoyoos Lake. After emerging from spawning gravel, mid-April to mid-May, they spend one or two summers (rarely three) as parr in the North Basin of Osoyoos Lake before an ontological shift (near autumnal equinox) to pre-smolts and subsequent migration the following spring as smolts through the reservoirs (with abundant predators), spillways, and smolt passage tunnels of ten hydroelectric dams to reach the Pacific Ocean via the Columbia River.

After one to three years (rarely more) of marine growth, adults return to the Columbia River in late summer and attempt the transit of ten dams in a 950 km freshwater migration, before entering Osoyoos Lake and subsequently spawning yet further upstream. Ascending the Okanagan River from the Columbia River is increasingly compromised by high river temperatures (Isaak et al., 2011; Hyatt et al., 2020; Bailey and Freshwater, 2023; Martins et al., 2012; Major and Mighell, 1967; Quinn et al., 1996).

Spawner estimates for Osoyoos sockeye have historically been based on trapezoidal AUC, with an assumed residence time of 11 days (?). However, residence time was never assessed for this population directly. Therefore, the selected residence time was based on the estimate for the early Stuart sockeye stock (Perrin and Irvine, 1990) because that stock has similar average body size, migration distance, and migratory timing (?).

3. Data

Surveys of live spawners have been conducted in the main sockeye spawning area—also known as the index section—of the Okanagan river between Osoyoos and Vaseau lakes (Figure 1) since 2000. All together, surveys have been conducted on 291 days, ranging between 9 and 19 surveys per year. All surveys occurred between August 28 and November 24.




Figure 1: The spawning grounds for Sockeye Salmon population that rear as parr in the North Basin of Osoyoos Lake are ~20 to 30 km upstream, in the Okanagan River below McIntyre Dam at the outflow of Vaseux Lake. Parr are predominantly in the North Basin. River flows that affect spawners, eggs, and parr are monitored by Environment Canada (EC) at the site indicated. Map by Braden Judson, Pacific Biological Station.

3.0.1. Survey methods

JLT, SVP. Survey methods and dataset preparations are those of REFERENCE.

3.0.2. problems re dead counts

JLT, SVP. I think our adventure from undocumented (?) survey methods is a valuable warning.

4. Methods

4.1. Problem: residence time

Spawners reside on spawning ground between repeated surveys, so a spawner may be counted repeatedly. The conventional estimate of abundance uses a value of residence to correct AUC for repeated observations of spawners. Residence is an assumed value, however, derived independently of spawner surveys. Can we do better?

4.2. Solution: spawner entry and exit

The timing and spread of spawners' entry to a spawning ground is reasonably modeled as a Gaussian (normal) probability density distribution (PDD). If residence is constant (ignoring sex, age, size, condition, injury, history, and genetics), then the PDD of spawner exits is the same as the PDD of entry, just delayed. The number of live spawners in a survey day is then the cumulative distribution of entries to that day minus the cumulative distribution of exits. This model is sufficient to estimate runs and residence simultaneously, but assumes all spawners are identical. In what follows, the assumption of fixed residence is relaxed to allow exit spread differ from entry spread. for a distribution of residence times, but the model is still based on the assumption that all spawners are identical.

If available, counts of newly dead spawners provide further information to estimate residence. Newly dead spawners accumulate between surveys, so the counts depend on survey intervals: the number of newly dead spawners is exits (deaths) cumulative to this survey day, minus deaths cumulative to the preceding survey. This assumes that dead spawners are marked by tags, by cutting off tails, or "dead pitch" (removal from the river bank) to ensure they are only counted once. Some dead spawners will drift away, not counted, necessitating a factor to estimate fraction of dead observed.

4.3. Model

An annual run of spawners enters a spawning ground with a Gaussian pattern for cumulative entry by day. They reside there for some days of spawning activity. Then spawner exits have the same distribution as entry, but lagged by residence. From this, the number of live spawners present on a survey day (Figure 2) is

$$\begin{aligned}\text{live}_{\text{true}}[t] &= \text{run} \ (\text{entered} - \text{exited}) \\ &= \text{run} \left(\int_0^t \mathcal{N}(t, \mu, \sigma) dt - \int_0^t \mathcal{N}(t, \mu + \lambda, \sigma) dt \right) \\ &= \text{run} \ (\text{pnorm}(t, \mu, \sigma) - \text{pnorm}(t, \mu + \lambda, \sigma))\end{aligned}$$

with *run*: spawner abundance (run size), *t*: survey day of the year, μ : timing, σ : spread, λ : residence, and *pnorm*: the R function for cumulative Gaussian.

The resulting temporal pattern for number of spawners present is close to a simple Gaussian when $\lambda < 4\sigma$ (exit overlaps entry) but platykurtic (flat-topped) for protracted residence. Additional parameters for skew and kurtosis could be used in the entry and exit pattern.

Counts of newly dead (Figure S3) depend on survey intervals: exits (deaths) cumulative to some survey day, minus exits cumulative to the preceding survey,

$$\text{dead}_{[t+n]} = \text{run} \cdot \phi \cdot \int_t^{t+n} \mathcal{N}(t, \mu + \lambda, \sigma) dt$$

with ϕ : the fraction of dead observed, and *t* and *t* + *n*: dates for consecutive surveys separated by *n* days. The first survey of dead spawners in a year observes an accumulation over an unlimited preceding interval; a small number or zero if surveys bracket spawning. The fraction of dead observed (ϕ) is likely to change within and between surveys, perhaps from varying flow. We note the potential for bias if ϕ varies by type of spawner (age, sex) when dead spawners are the source of biometrics data.

The model parameters are further described in the simulation model (Supplementary), including prior distributions and parameters discovered as analysis progressed.

4.4. Simulations

We challenged this model with simulated observations to ensure robust and reliable results. Parameters were sampled randomly from prior PDDs. Simulated surveys were deliberately poor: the number of surveys each year was random uniform between 6 and 10, and survey dates within years random uniform but separated by at least 1 day. The resulting “true” data, but with multiplicative normal sampling error (% accuracy), was the “observed” data subsequently fitted by the multi-level regression coded in Stan. The scenarios explored fits (see




Figure 2: Example of how the model estimates total spawner abundance, arrival timing, arrival spread, mean residence, and death spread based on the observed counts (points) of spawners on multiple days. The number of spawners present on a given day is the difference between accumulated arrivals and accumulated deaths. Mean residence time is estimated as a parameter which defines the mean difference in timing of arrivals and death. We assume that spawner counts are imperfect observations of the true number, and so individual points vary from the estimated presence value. The total abundance of spawners is estimated as the total number that have arrived. All model parameters are estimated with uncertainty — lines show the posterior median and bands show the 95% credible intervals. In the case of arrived and dead numbers, uncertainty increases with abundance because uncertainty from the timing and spread of arrival or death is multiplied by uncertainty from abundance.

Stan Model) with and without surveys of newly dead spawners. Despite noisy observations and gaps between surveys, parameters were recovered accurately from Stan model, with the caveat that the model used to fit the simulated data was the model that created it. Specifically, this procedure correctly estimated residence over a wide range. The simulations also generated indices of relative abundance of spawners, such as AUC as Gaussian and as linear interpolations (a trapezoid), peak live, and peak live puls dead (PLD). These were compared to the fitted run size to indicate reliability.

4.5. Stan Models

We used the Stan programming language (?) to code series of increasingly complicated series of non-linear, multi-level Bayesian regressions to challenge this model with data. We explored which parameters were important and the trade-off of complication with precision. Our initial fits used explicit and tightly bounded prior PDD (informed priors), and initiated Monte Carlo sampling chains with parameter estimates derived from the data, but this was not necessary, the default procedures successfully initiated sampling chains. See Supplementary Materials for the Stan code that fits these models, plus the R code run it. includes this code, and details regarding convergence and model comparisons.

Differences between the models are summarised in Figure 3 A.

outline: describe the different multi-level Stan models (which, why)
 details of Stan models: how and why: priors, inits, convergence,

4.5.1. Likelihood Model

JLT, please. About sampling negative binomial residuals to fit. Compared to lognormal:

```
target <- normal_lpdf(log_run_predicted|log_run_predicted, residuals)
```

Our negative binomial distribution accounts for overdispersion in observed counts as live_ϕ , which includes both observation error as well as unmodelled variation in spawner abundance.

The observed (dependent) variable, y_{obs} , is predicted, y_{pred} , by a sample of the parameters from the chosen model. An assumed PDD for the residuals, $y_{pred} - y_{obs}$, provides the likelihood of each observation given the model and a sample from its parameters space. There is a prior for the residuals PDD. That likelihood guides subsequent sampling to converge on a “best fit” multivariate PDD for the parameters [Gelman et al. \(2013\)](#).

4.5.2. Diagnostics

An essential concept in Bayesian statistics is that the data is known but the best model for that data is unknown [McElreath \(2016\)](#) and always a vast simplification of reality [Jaynes \(2003\)](#). Metrics to compare and rank competing models are required. In ecological systems, a superabundance of habitat indicators as covariates means over-fitting is a problem. Leave-one-out (LOO) statistics², where each point is predicted while omitted from fitting (cross-validations), provides diagnostics for over-fitting. We used **ELPD**, the sum of expected log predictive probability density for each of n omitted observation's prediction, and $ELPD_{SE} = \text{sd}(ELPD)/\sqrt{n}$ the standard error of those probabilities [Vehtari et al. \(2017\)](#).

5. Results

outline: Simulations recapture residence. why critical. Tables *
descriptions and fits (ELPD, AIC,) comparison of Stan models
* parameter estimates and statistics for best model (so far).
Plots
* example of model with CLs to entry and exit CPDs, observed
datapoints, and CLs for predicted curve. Just one year. * timing,
spread, residence by years, with CLs. Fancier the better.

5.1. Model selection and convergence

Based on ELPD statistics (**?@fig-mod_compare B**) we selected model 7 as the best from those explored. This model included year-to-year variation in spawner abundance, arrival timing, arrival spread, and death spread, but assumed that residence time and observation error is constant across years. Model 8, which also allowed residence time to vary across years had an equivalent ELPD ($\Delta\text{ELPD} = -1.6$, $\text{SE} = 2.1$).

The effective number of parameters (p_{loo}) increased when arrival timing variation was introduced (model 1 vs. 2), and also when any of arrival spread, death spread, or residence were allowed to vary (models 3 to 8; **?@fig-mod_compare C**). However, all models with variation in any or all of these parameters (models 3 to 8) had similar numbers of effective parameters.

Convergence was good for model 7; all parameters had $\hat{R}$ values ≤ 1.006 and effective sample sizes ≥ 804 . Allowing residence time to vary across years (model 8), resulted in some minor convergence issues; 3.1% of parameters had $\hat{R} > 1.01$,

²see R package LOO <https://cran.r-project.org/web/packages/loo/loo.pdf> and LOO Package Glossary <https://mc-stan.org/loo/reference/loo-glossary.html>.

and the highest $\hat{R}$ was 1.03. Model 8 also had 1.3% of its parameters with effective sample sizes < 400 , the lowest being 210. The reason that model 8 exhibited minor convergence issues is likely because of a posterior correlation between residence time and the other timing parameters, which are only partially identifiable from the observed spawner counts. Thus, we have selected the simpler model (model 7), which has equivalent predictive power and which does not have convergence issues, but we contrast the results from both where appropriate.




Figure 3: Model summary and comparison. *Panel A* shows which parameters were held constant (white) versus varied (black) across years in each of the candidate models (rows). *Panel B* shows the relative expected log predictive density (ΔELPD) compared to the model with the highest out-of-sample predictive accuracy. *Panel C* shows the effective number of parameters (p_{100}). The model with the highest out-of-sample predictive accuracy is highlighted in red in panels B and C. Points and error bars (B and C) represent means and standard errors.

5.2. Model fit to observed counts

Our expected counts correspond well with the observed counts Figure 4 and Figure S2. Variation between the observed and expected is quantified in our estimate of spawner count overdispersion (live_ϕ), which had a posterior median of 6.9 (95% CI: 5.6–8.5).

5.3. Spawning timing

The average arrival day (arrival_μ) across years was estimated to be 285.4 (posterior median; 95% CI: 283.7–287) which corresponds approximately to October 12. Interannual variation in arrival date (arrival_σ) was estimated as a standard deviation of 1.6 days. As a result, the median estimated arrival date ranged from October 10 to October 15 (Figure 5 A).




Figure 4: Observed versus estimated spawners present over the spawning period in each year (panels). Points show the observed counts, the lines and shaded bands show the posterior median the 95% credible intervals.

The average arrival spread ($e^{\text{arrival_spread}_\mu}$) across years was estimated to be 7.6 days (posterior median; 95% CI: 6.7–8.5). This implies that 95% of spawners enter within a span of 29.7 days. Arrival spread varied across years by a multiplicative factor ($e^{\text{arrival_spread}_\sigma}$) of 1.2. As a result, the median estimated arrival spread ranged from 5 to 7 days (Figure 5 B).

The average death spread ($e^{\text{death_spread}_\mu}$) across years was estimated to be 6.7 days (posterior median; 95% CI: 5.9–7.6). Death spread varied across years by a multiplicative factor ($e^{\text{death_spread}_\sigma}$) of 1.2. As a result, the median estimated death spread ranged from 4 to 8 days (Figure 5 C).

Residence was estimated as 8.8 days (95% CI: 6.4–11.6; Figure 6). In the alternative model (m8) where residence is allowed to vary across years, the median residence time across years was 9.3 days (Figure S1 D; inter-annual range: 7–13.3).

5.4. Total spawner abundance

Estimated posterior median annual spawner abundances vary from as low as 3700 in 2002 to as high as 176,000 in 2010. Since 2008, spawner abundance has alternated between low and high levels in a two-year cycle.

Our estimated annual abundances are, on average, slightly higher than those from the traditional trapezoidal AUC estimate, with an assumed 11 day residence time. The median difference across all years is 3.9 % (inter-annual range: -10.8–11.6), which is a smaller difference compared to the 19.9 increase that we would expect if we were to apply our estimated residence time (8.8 days) to the traditional method. In addition, while our spawner abundance estimate tends to be higher, this is only the case in 16% of the 25 years and the traditional estimate falls within the 66% credible interval of our model prediction in all years.

6. Discussion

outline: General- new model is successful.

Run size estimates are more accurate than AUC.

Which parameters (models) were important (annual spread and timing), which not (varying residence, autocorrelation of surveys)? Residence varies little, so conventional is a reliable relative index (?) (but I need to understand how you changed shape of exit).

How to accommodate biometrics: if early entries (big females?) stay longer,. Scope for covariates: biometrics and habitat, parameter correlations,. Events like Testalinden. scope for more detail: model to predict counts by segment. Value of (good) surveys of newly dead. (?? Value of analyzing all data simultaneously. possible extensions: * autocorrelation in spawner runs (ages, SAR,); * models for entire life history (add parr, smolts, returns (SAR), prespawning mort)




Figure 5: Posterior distributions of timing, entry_spread, and exit_spread. Points and error bars show the posterior median and the 66% and 95% credible intervals.




Figure 6: Posterior distribution of the estimated mean residence time. Points and error bars show the posterior median and the 66% and 95% credible intervals. The vertical dashed line shows the 11 day residence time that has previously been assumed for this system.




Figure 7: Posterior distributions of total spawning abundance by year (black) compared to the point estimate based on traditional trapezoidal AUC with an assumed 11 day residence (red). Points and error bars show the posterior median and the 66% and 95% credible intervals.

6.1. Residence and exit spread.

The attempt to estimate residence directly from spawner surveys was successful. For this species and place, the estimate was ?? days ($\$ = ??$). This is approximately twice the spread of entry, implying nearly half of the spawners enter before newly-dead are abundant. Our estimate suggests that residence time is likely lower than the value (11 days) previously used to correct for AUC (?), which sits at the 94th percentile of our distribution. Furthermore, we identified annual variability in exit spread that is related to changes in residence. Our interpretation is that residence time is a function of date of entry, likely because early entrants stay longer than late entrants, a pattern identified by REFERENCES and ascribed to age and sex. Residence as a linear function of entry changes exit spread compared to entry spread:

$$\mathcal{N}(x, \mu, \sigma) + (a + bx) = \mathcal{N}(x, \mu + a, b\sigma)$$

where a is the mean residence time (constant), and b where ($0 < b < 1$) describes how much the exit spread shrinks from entry spread. If entry spread = 5 days and $b = 0.8$ then exit spread = 4. We determined exit spread directly rather than determining b because the underlying ecology will be more complicated than a single, linear, parameter. Inspection of Figure 5 reveals exit spread is not correlated with entry spread, but does share a common temporal trend with entry timing (recently later and longer). That exit spread varies is an intriguing discovery awaiting explanation.

6.2. Annual run size

Our model estimates precision for annual run size (spawner abundance). This is valuable as regression weights for subsequent application (parr from spawners, returns from spawners). Estimated as a negative binomial coefficient, it is a multiplicative factor (% accuracy) of $??(\sigma = ??)$. As a coefficient of variation (σ/μ), this varied from $??(\sigma = ??)$ in 2000 to $??(\sigma = ??)$ in 2000 (Figure 7). Note this precision includes model error as well as observation error.

Because our estimate of mean residence time is similar to value used to correct conventional AUC estimates, the 95% uncertainty intervals for run estimates from this analysis overlap most of the conventional estimates. The exception is determining that conventional estimates (red dots in Figure 7) for years 2016 to 2018 might be substantial underestimates. A severe overestimate in 2011 was noted, but investigation revealed the conventional estimate had been increased, arbitrarily, to better match spawner passage estimates from Columbia River dams. Conventional estimates could not be accessed for years after 2019. Although the results for Osoyoos Lake Sockeye, and the results from simulations, provide initial confidence in the accuracy of these run estimates, complete confidence awaits resolution of the 2016-2018 discrepancies and results from different salmon populations.

6.3. Entry timing and spread

Timing and spread of spawner entry are ecologically informative parameters not previously extracted from spawner surveys. Timing overall was day 285($\sigma = ??$) and ranged from day 282 in 2013 to 288 in 2002, less than a week. Although recent timings (2016–2024) do not have any of the earliest timings (2000–2015) this is not a linear trend ($r^2 = 0.00$).

Entry spread was 285($\sigma = ??$) days, ranged from 6 days (2013) to 11 days (2002). Although the estimate of 12.7 days for 2010 is the widest estimate, this is associated with the 2010 June 13 Testalinden Creek dam failure that disgorged silt and toxins to the Okanagan River downstream of the spawning region, but affecting Osoyoos Lake and the Okanagan River below it. Entry spread is the only parameter apparently affected by that event; how that affected spawners, if it did, has yet to be determined.

6.4. Relative importance of parameters

Which parameters (models) were important: annual spread and timing; and which not: varying residence, autocorrelation of surveys,?

6.5. Adding covariates: biometrics and habitat indices

This model is the basis for expanding the information applied to analyses of spawner surveys. Perrin and Irvine [Perrin and Irvine \(1990\)](#) identified a suite of potential covariates for residence, including spawner density, sex, body size, body size by sex (females stay longer, to guard redds, than males), sex ratio, run timing (a year with late timing may have shorter residence), time of arrival, flow, and water temperature; concluding *Since residence can vary among years within a stream, and among streams within a year, residence estimates should not be transferred among streams or years. This can produce serious errors in the escapement estimate. Residence should be determined on a site specific basis each time the AUC estimate is used to estimate escapement* [replacing ‘survey life’ with ‘residence’].

Spawner biometrics affect entry timing, entry spread, and residence (thus exit spread): *In general, older and larger wild Atlantic salmon return to freshwater to spawn earlier in the season than younger and smaller salmon, and female salmon return earlier in the season than males* [Miettinen et al. \(2024\)](#).

EXACTLY HOW SHOULD BIOMETRIC AND HABITAT DATA BE INCLUDED, AND VALUE THEREOF BE DETERMINED?

Spawner surveys collect length, sex, and age (otoliths and/or scales) from dead salmon, with varying precisions for mean length at age by sex. Typically this can be augmented with biometrics from catches, counting fences, and dam passage. Relevant to this population, [Bailey et al. \(2025\)](#) (their appendix table 5) identified large proportions of ocean age 1 spawners (jacks) in 2019, 2007, 2021, and 2003; respectively 55.8%, 48.2%, 41.9%, and 38.3%. These years are all low runs (< 2,000), and do not have unusual entry timing, entry spread, or exit spread.

6.6. Multiple rivers

This analysis of multiple years together might be extended to include multiple rivers if spawning groups in those rivers covary consistently as parts of population or possibly a conservation unit (metapopulation). As an equation,

$$N_{tjk} = N_j P_k (T_0[\mu, \sigma]_t + T_j[\mu, \sigma]_t + T_k[\mu, \sigma]_t) + \epsilon_{tjk}$$

where N_{tjk} is the number of spawners entering river k on day t in year j , estimated with error ϵ from the proportion P_k of all spawners that year N_j that enter river k according to how the background temporal pattern T_0 is affected by the pattern for that year T_j and that river T_k . Terms $T_x[\]_t$ are temporal distributions, with μ_x and σ_x for timing and spread if Gaussian.

6.7. *Residuals: lognormal versus negative binomial*

Weights are changed by log transform and/or choice of residuals model. Do you have anything to say about that changes results, or advice to users of this model? If no, delete this section.

Fitting a model to the log of abundance observations changes the relative importance (effect) of the data points, it emphasizes the contribution to the sum of squares from the residuals at small abundance estimates compared to large. The transform involves asking (a) are small abundance estimates more, or less, precise than large? and (b) which abundances are more important to estimate accurately? Minimizing residuals from log values, as done here, implies sampling error is proportional to the observation (to the true abundance): it is multiplicative as in % accuracy, rather than additive. If the precision of each abundance estimate can be quantified, w , then a weighted regression is appropriate $\hat{y} = \mathcal{N}(a + bx, \sigma w)$. Note that ordinal weights (ranked, not quantified) can be used by fitting an additional parameter, $\hat{y} = \mathcal{N}(a + bx, \sigma w^{-\lambda})$.

6.8. *Further sources of uncertainty*

Again, if you want to say something about methods, that will be valuable. Else, delete this section.

Observer accuracy is a calibration problem, recognized as efficiency, the fraction of fish observed (bias), but includes uncertainty. This varies by method. Comparison of observers may involve factors including time of day, weather conditions, and fish density. For instance, counts while snorkeling are not possible during high flows or turbid water, and are truncated at high spawner densities.

6.9. *Value of surveys for newly dead spawners*

Awaits comparisons from simulations. Could be important.
I think readers of this paper would benefit from your experience, discovering counts of dead spawners were misleading, dead pitch events were few compared to boat surveys of live.

7. Conclusions

outline: Overall: This new approach delivered more accurate runs and estimated ecologically important variables. Runs – estimates with precision. Accuracy is better than AUC. Newly Dead – these data would have been informative. Residence * precise, CV 5%. * Did this identify an important bias in AUC? * Did not vary greatly between years (unexpected.) Timing, Spread – precise estimates, opportunity to add more ecology.

1. Residence time, a key parameter, was successfully estimated by taking advantage of newly dead spawners surveyed in multiple years spawner surveys.
The value obtained is twice the assumed value, suggesting spawner abundance from AUC was previously overestimated.
2. The fraction of dead spawners observable (not drifted) was estimated as $\sim 2/3$, but refinement is warranted.
3. more
4. more

8. Recommendations

We propose two groups of suggestions.

As a generalization, salmon datasets are not maintained following best practices ([Diack et al. \(2024\)](#)). As we discovered during this analysis, datasets from spawner surveys warrant attention to:

1. Documenting survey methods, including data processing. This is essential to avoid erroneous assumptions by analysts. We point to the website [Monitoring Resources](#) maintained by the [Pacific Northwest Aquatic Monitoring Partnership](#).
2. Datasets that are F.A.I.R. ([Wilkinson et al. \(2016\)](#)). Cataloging the existence of proprietary datasets is rewarded from judicious sharing.

Improving the accuracy of spawner run size estimates appears possible by using more of the information collected in spawner surveys but not used. This model can be expanded with further ecological realism. Simulations showed how surveys of newly dead spawners are useful to that end. With the caveat that adding complications and noisy datasets will have diminishing returns, we foresee including:

1. Habitat variables (river flow, temperature,) applied to improve the spawner entry and models; likely asymmetric or discontinuous in response to flow events;
2. Covariates from the biometrics of spawners: age, sex, age*sex, genomics, and genetics;
3. Including correlations between annual parameters (noted) might improve accuracies;
4. Temporal trends (runs, timings) are valid covariates *per se*, but ideally related to habitat variables;
5. Surveys in a year are autocorrelated, an important consideration; and
6. Sockeye spawners observed in specific year produce spawners returning three to six years later; information from a salmon's life-cycle is not extraneous to spawner surveys.

9. Acknowledgements

Diligence, care, and persistence by survey field teams from the Okanagan Nations Alliance (ONA, sylix.org) resulted in a spawners dataset based on unusually thorough and consistent methods; their commitment to *cause to come back* is inspiring. This paper commemorates accomplishments of the late Dr. Kim Hyatt who designed, implemented, and supported this survey program. We thank Karilyn Alex (ONA) and Collette Louie (Osoyoos Indian Band, oib.ca) for access and permission regarding this proprietary data, possible through the data sharing agreement of the Canadian Okanagan Basin Technical Working Group (COBTWG). We thank Howard Stiff and Athena Ogden (Pacific Biological Station, Fisheries and Oceans Canada) and Elizabeth Ng (Four Peaks Environmental Ltd.) for the work required to reconcile and summarize 25 years of field notes and spreadsheets into the quality controlled and standardized dataset we received.

* I can't find a reference to the data-sharing agreement *per se*.

Dr. James R. Irvine (DFO Pacific, retired) improved text based on his extensive knowledge of salmon surveys and related analyses.

9.1. Competing Interests

The authors declares no competing interests.

9.2. Funding Statement

The authors declares no specific funding for this work.

9.3. Data Availability

The data used in this study are the property of the Okanagan Nations Alliance. Please contact them directly concerning availability and acknowledgements: Sylix.org.

Figures

not used

Tables

not used




Figure S1: Posterior distributions of timing, entry_spread, exit_spread, and residence from the alternative model that also includes year-to-year variation in residence. Points and error bars show the posterior median and the 66% and 95% credible intervals.




Figure S2: Observed survey counts versus predicted. Points and error bars show the posterior median and the 66% and 95% credible intervals.




Figure S3: Newly dead spawners each day (blue) is the rate of exits. Newly dead accumulate between surveys, here with alternating 1 day and 4 day separations, leading to an irregular pattern of counts (red). Parameters as in preceding figure.

10. Supplementary Materials

The following documents are available for public access on [GitHub](#) and [Zenodo](#):
FIX LINKS

1. spawners.stan – Stan code for the models
2. Spawners_est.R – This is the R code to run the Stan model and plot results.
3. Newly Dead Spawners.pdf – Simulations report as PDF.
4. Newly Dead Spawners.qmd – Simulation code to describe, run, and report the model.
5. newly dead spawners.stan – Stan code for simulations.

References

- Adkison, M., Su, Z., 2001. A comparison of salmon escapement estimates using a hierarchical bayesian approach versus separate maximum likelihood estimation of each year's return. *Canadian Journal of Fisheries and Aquatic Sciences* 58, 1663–1671. doi:[10.1139/cjfas-58-8-1663](#).
- Askey, P.J., Ward, H.G., Weir, T., King, K., 2023. Comparison of known spawner abundance from fence counts to visual counts for simplified spawner estimation methods. *Canadian Journal of Fisheries and Aquatic Sciences* 80, 1935–1948. doi:[10.1139/cjfas-2023-0111](#).

- Bailey, C., Freshwater, C., 2023. State of the physical, biological and selected fishery resources of pacific canadian marine ecosystems in 2022, Fisheries and Oceans Canada. pp. 133–140. URL: <https://waves-vagues.dfo-mpo.gc.ca/library-bibliotheque/41199248.pdf>.
- Bailey, C., Stiff, H., Thompson, P., Judson, B., Ogden, A., 2025. Osoyoos lake sockeye salmon return abundance (1977-2023), adult age composition (1985-2023), and marine survival (1998-2021). Canadian Manuscript Report of Fisheries and Aquatic Sciences 3282, vi+49. URL: <https://waves-vagues.dfo-mpo.gc.ca/library-bibliotheque/41273965.pdf>, doi:10.60825/v3g8-jw96.
- Berghe, E., Gross, M., 1986. Length of breeding life of coho salmon (*Oncorhynchus kisutch*). Can. J. Zool 64, 1482–1486. doi:10.1139/z86-221.
- Bue, B., Fried, S., Sharr, S., Sharp, D., Wilcock, J., Gieger, H., 1998. Estimating salmon escapement using the area-under-the-curve, aerial observer efficiency, and stream-life estimates: The prince william sound pink salmon example. North Pacific Anadromous Fish Commission Bulletin 1, 240–245.
- Cormack, R., 1964. Estimates of survival from the sighting of marked animals. Biometrika 51, 429–438.
- Diack, G., Bird, T., Akenhead, S., Bayer, J.M., Brophy, D., Bull, C., Eyto, E.d., Hanson, N., Johnson, B., Jones, M.B., Knight, A., Nevoux, M., Stap, T.C.A.v.d., Walker, A., 2024. Salmon data mobilization. N. Pac. Anadr. Fish Comm. Bull. 7, 61–76. URL: <https://osf.io/hk4gu>, doi:<https://doi.org/10.23849/npafcb7/x3rlpo23a>. publisher: OSF.
- English, K., Bocking, R.C., Irvine, J.W., 1992. A robust procedure for estimating salmon escapement based on the area-under-the-curve method. Can. J. Fish. Aquat. Sci 49, 1982–1989. URL: https://www.researchgate.net/publication/233865122_A_Robust_Procedure_for_Estimating_Salmon_Escapement_based_on_the_Area-Under-the-Curve_Method, doi:10.1139/f92-220. publication Title: Canadian Journal of Fisheries and Aquatic Sciences DOI: 10.1139/f92-220.
- Fukushima, M., Smoker, W., 1997. Determinants of stream life, spawning efficiency, and spawning habitat in pink salmon in the auke lake system, alaska. Canadian Journal of Fisheries and Aquatic Sciences 54, 96–104.
- Gelman, A., Carlin, J., Stern, H., Dunson, D., Vehtari, A., Rubin, D., 2013. Bayesian data analysis. Third ed., Chapman and Hall, New York.
- Hilborn, R., Bue, B., Sharr, S., 1999. Estimating spawning escapements from periodic counts: a comparison of methods. Canadian Journal of Fisheries and Aquatic Sciences 56, 888–896.
- Hilborn, R., Walters, C.J., 1992. Quantitative Fisheries Stock Assessment: Choice, Dynamics and Uncertainty. Springer US, Boston, MA. URL: <http://>

[//link.springer.com/10.1007/978-1-4615-3598-0](https://link.springer.com/10.1007/978-1-4615-3598-0), doi:[10.1007/978-1-4615-3598-0](https://doi.org/10.1007/978-1-4615-3598-0).

- Hyatt, K., Stiff, H., Stockwell, M., 2020. Historic water temperature (1924-2018), river discharge (1929-2018), and adult Sockeye Salmon migration (1937-2018) observations in the Columbia, Okanogan, and Okanogan rivers. Technical Report.
- Irvine, J.R., Bocking, R.C., English, K.K., Labelle, M., 1992. Estimating coho salmon (*Oncorhynchus kisutch*) spawning escapements by conducting visual surveys in areas selected using stratified random and stratified index sampling designs. *Can J Fish Aquat Sci* 49, 1972–1981.
- Isaak, D., Wollrab, S., Horan, D., Chandler, G., 2011. Climate change effects on stream and river temperatures across the northwest us from 1980-2009 and implications for salmonid fishes. *Climatic Change* 113, 499–524. doi:[10.1007/s10584-011-0326-z](https://doi.org/10.1007/s10584-011-0326-z).
- Jaynes, E.T., 2003. Probability Theory, The Logic Of Science. Cambridge University Press. ISBN-13 978-0-511-06589-7. Cambridge University Press.
- Korman, J., Ahrens, R., Higgins, P., Walters, C., 2002. Effects of observer efficiency, arrival timing, and survey life on estimates of escapement for steelhead trout (*Oncorhynchus mykiss*) derived from repeat mark recapture experiments. *Can J Fish Aquat Sci* 59, 1116–1131.
- Labelle, M., Bussanich, A., Benson, R., 2021. Maximum likelihood estimation of kokanee and sockeye salmon spawners in a stream using visual survey data, 2003-2017. *Fisheries and Aquaculture Journal* 12. URL: <https://www.longdom.org/open-access/maximum-likelihood-estimation-of-kokanee-and-sockeye-salmon-spawners-in-a-stream-using-visual-survey-data-20032017-70830.html>, doi:[10.35248/2150-3508.21.12.277](https://doi.org/10.35248/2150-3508.21.12.277).
- Lady, J., Skalski, J., 1998. Estimators of stream residence time of pacific salmon (*Oncorhynchus* spp.) based on release-recapture data. *Can. J. Fish. Aquat. Sci* 55, 2580–2587.
- Liao, S., 1994. Statistical models for estimating salmon escapement and stream residence time based on stream survey data. Ph.D. thesis. Seattle.
- Major, R., Mighell, J., 1967. The influence of rocky reach dam and the temperature of the okanogan river on the upstream migration of sockeye salmon. *Fishery Bulletin* 66, 131–147. URL: https://fisherybulletin.nmfs.noaa.gov/sites/default/files/pdf-content/fish-bull/major_0.pdf.
- Manske, M., Schwarz, C., 2000. Estimates of stream residence time and escapement based on capture- recapture data. *Canadian Journal of Fisheries and Aquatic Sciences* 57, 241–246.

- Martins, E.G., Hinch, S.G., Patterson, D.A., Hague, M.J., Cooke, S.J., Miller, K.M., Robichaud, D., English, K.K., Farrell, A.P., 2012. High river temperature reduces survival of sockeye salmon (*oncorhynchus nerka*) approaching spawning grounds and exacerbates female mortality. *Canadian Journal of Fisheries and Aquatic Sciences* 69, 330–342. URL: <https://cdnsiencepub.com/doi/10.1139/f2011-154>, doi:10.1139/f2011-154. publisher: NRC Research Press.
- McElreath, R., 2016. Statistical rethinking: A Bayesian course with examples in R and Stan. CRC Press. Taylor and Francis, Boca Raton.
- Miettinen, A., Romakkaniemi, A., Dannewitz, J., Pakarinen, T., Palm, S., Persson, L., Östergren, J., Primmer, C.R., Pritchard, V.L., 2024. Temporal allele frequency changes in large-effect loci reveal potential fishing impacts on salmon life-history diversity. *Evolutionary Applications* 17, e13690. URL: <https://onlinelibrary.wiley.com/doi/abs/10.1111/eva.13690>, doi:10.1111/eva.13690.
- Millar, R., McKechnie, S., Jordan, C., 2012. Simple estimators of salmonid escapement and its variance using a new area-under-the-curve method. *Canadian Journal of Fisheries and Aquatic Sciences* 69, 1002–1015. URL: <https://cdnsiencepub.com/doi/10.1139/f2012-034>, doi:10.1139/f2012-034. publisher: NRC Research Press.
- Neilson, J., Geen, G., 1981. Enumeration of spawning salmon from spawner residence time and aerial counts. *Trans. Am. Fish. Soc* 110, 554–556. doi:10.1577/1548-8659(1981)110.
- Parken, C.K., Bailey, R.E., Irvine, J.R., 2003. Incorporating uncertainty into area-under-the-curve and peak count salmon escapement estimation. *North American Journal of Fisheries Management* 23, 78–90. URL: <https://onlinelibrary.wiley.com/doi/abs/10.1577/1548-8675%282003%29023%3C0078%3AIUIAUT%3E2.0.CO%3B2>, doi:10.1577/1548-8675(2003)023<0078:IUIAUT>2.0.CO;2.
- Parsons, A.L., Skalski, J.R., 2010. Quantitative assessment of salmonid escapement techniques. *Reviews in Fisheries Science* 18, 301–314. URL: <https://doi.org/10.1080/10641262.2010.513020>, doi:10.1080/10641262.2010.513020.
- Perrin, C., Irvine, J.R., 1990. A review of survey life estimates as they apply to the area-under-the-curve method for estimating the spawning escapement of pacific salmon. Technical Report. Pacific Biological Station, Nanaimo BC. URL: https://publications.gc.ca/collections/collection_2014/mpo-dfo/Fs97-6-1733-eng.pdf. iSSN 1488-5379.
- Quinn, T.P., Adkison, M.D., Ward, M.B., 1996. Behavioral tactics of male sockeye salmon (*oncorhynchus nerka*) under varying operational sex ratios. *Ethology* 102, 304–322. URL: <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1439-0310.1996.tb01127.x>, doi:10.1111/j.1439-0310.1996.tb01127.x.

- Ricker, W.E., 1975. Computation and interpretation of biological statistics of fish populations. *Fish. Res. Board Can. Bull.* 191, 1–382.
- Semenchenko, N., 1993. Life expectancy of spawning grounds for spawning grounds of sockeye salmon *oncorhynchus nerka*. *Vopr. Ikhtiol* 33, 550–555.
- Su, Z., Adkison, M.D., Van Alen, B.W., 2001. A hierarchical bayesian model for estimating historical salmon escapement and escapement timing. *Canadian Journal of Fisheries and Aquatic Sciences* 58, 1648–1662. URL: <https://cdnsiencepub.com/doi/10.1139/f01-099>, doi:10.1139/f01-099. publisher: NRC Research Press.
- Thomason, G., Jones, J., 1984. Southeastern alaska pink salmon (*oncorhynchus gorbuscha*) stream life studies .
- Vehtari, A., Gelman, A., Gabry, J., 2017. Practical bayesian model evaluation using leave-one-out cross-validation and waic. *Statistics and Computing* 27, 1413–1432. URL: <https://arxiv.org/abs/1507.04544>, doi:10.1007/s11222-016-9696-4. <https://mc-stan.org/loo/index.html>.
- Wilkinson, M.D., Dumontier, M., Aalbersberg, I.J., Appleton, G., Axton, M., Baak, A., Blomberg, N., Boiten, J.W., da Silva Santos, L.B., Bourne, P.E., Bouwman, J., Brookes, A.J., Clark, T., Crosas, M., Dillo, I., Dumon, O., Edmunds, S., Evelo, C.T., Finkers, R., Gonzalez-Beltran, A., Gray, A.J., Groth, P., Goble, C., Grethe, J.S., Heringa, J., 't Hoen, P.A., Hooft, R., Kuhn, T., Kok, R., Kok, J., Lusher, S.J., Martone, M.E., Mons, A., Packer, A.L., Persson, B., Rocca-Serra, P., Roos, M., van Schaik, R., Sansone, S.A., Schultes, E., Sengstag, T., Slater, T., Strawn, G., Swertz, M.A., Thompson, M., van der Lei, J., van Mulligen, E., Velterop, J., Waagmeester, A., Wittenburg, P., Wolstencroft, K., Zhao, J., Mons, B., 2016. The fair guiding principles for scientific data management and stewardship. *Scientific Data* 3, 160018. URL: <https://doi.org/10.1038/sdata.2016.18>, doi:10.1038/sdata.2016.18.
- Zaporozhets, O., Zaporozhets, G., 2021. Experimental determination of the lifespan of sockeye salmon spawners at the spawning ground in the littoral of lake nachikinskoye (kamchatka. *TINRO News* 201, 313–323. doi:10.26428/1606-9919-2021-201-313-323.