# Greedy Algorithms: Main Ideas

Michael Levin

Higher School of Economics

# Algorithmic Toolbox Data Structures and Algorithms

#### Outline

- 1 Largest Number
- 2 Car Fueling
- 3 Implementation and Analysis
- 4 Main Ingredients

#### Learning objectives

Come up with a greedy algorithm yourself











3 5 9 1 7 9

#### Largest Number

#### Toy problem

What is the largest number that consists of digits 3, 9, 5, 9, 7, 1? Use all the digits.

#### Largest Number

#### Toy problem

What is the largest number that consists of digits 3, 9, 5, 9, 7, 1? Use all the digits.


#### Examples

359179, 537991, 913579, . . .


#### Correct answer

#### 


5 7 3 9 1 9 →



■ Find max digit



- Find max digit
- Append it to the number



- Find max digit
- Append it to the number
- Remove it from the list of digits

Find max

Remove


- Find max digit
- Append it to the number
- Remove it from the list of digits

Find max

$$\boxed{5 \ 7 \ 3} \boxed{1 \ 9} \rightarrow \boxed{9}$$


#### Remove

- Find max digit
- Append it to the number
- Remove it from the list of digits
- Repeat while there are digits in the list




#### Remove

- Find max digit
- Append it to the number
- Remove it from the list of digits
- Repeat while there are digits in the list



- Find max digit
- Append it to the number
- Remove it from the list of digits
- Repeat while there are digits in the list



- Find max digit
- Append it to the number
- Remove it from the list of digits
- Repeat while there are digits in the list

Find max


#### Remove

- Find max digit
- Append it to the number
- Remove it from the list of digits
- Repeat while there are digits in the list


# Find max $\downarrow$ $5 7 3 1 \rightarrow 9 9$

#### Remove

- Find max digit
- Append it to the number
- Remove it from the list of digits
- Repeat while there are digits in the list



- Find max digit
- Append it to the number
- Remove it from the list of digits
- Repeat while there are digits in the list



- Find max digit
- Append it to the number
- Remove it from the list of digits
- Repeat while there are digits in the list


Find max

```
\boxed{5} \boxed{3} \boxed{1} \rightarrow \boxed{9} \boxed{9} \boxed{7}
```

#### Remove


- Find max digit
- Append it to the number
- Remove it from the list of digits
- Repeat while there are digits in the list

#### Find max




#### Remove

- Find max digit
- Append it to the number
- Remove it from the list of digits
- Repeat while there are digits in the list



- Find max digit
- Append it to the number
- Remove it from the list of digits
- Repeat while there are digits in the list



- Find max digit
- Append it to the number
- Remove it from the list of digits
- Repeat while there are digits in the list


#### Find max



#### Remove


- Find max digit
- Append it to the number
- Remove it from the list of digits
- Repeat while there are digits in the list






#### Remove

- Find max digit
- Append it to the number
- Remove it from the list of digits
- Repeat while there are digits in the list




- Find max digit
- Append it to the number
- Remove it from the list of digits
- Repeat while there are digits in the list



- Find max digit
- Append it to the number
- Remove it from the list of digits
- Repeat while there are digits in the list


#### Find max



#### Remove

- Find max digit
- Append it to the number
- Remove it from the list of digits
- Repeat while there are digits in the list

#### Find max



#### Remove

- Find max digit
- Append it to the number
- Remove it from the list of digits
- Repeat while there are digits in the list

## 

- Find max digit
- Append it to the number
- Remove it from the list of digits
- Repeat while there are digits in the list

# 

- Find max digit
- Append it to the number
- Remove it from the list of digits
- Repeat while there are digits in the list

#### Find max



#### Remove

- Find max digit
- Append it to the number
- Remove it from the list of digits
- Repeat while there are digits in the list

# Greedy Strategy

- Find max digit
- Append it to the number
- Remove it from the list of digits
- Repeat while there are digits in the list


### Outline

- 1 Largest Number
- 2 Car Fueling
- 3 Implementation and Analysis
- 4 Main Ingredients

Distance with full tank = 400km

Distance with full tank = 400km

950km



0km

Distance with full tank = 400km

0km 200km 375km 550km 750km 950km

Distance with full tank = 400km



Distance with full tank = 400km



Distance with full tank = 400km

Minimum number of refills = 2



A car which can travel at most L kilometers with full tank, a source point A, a destination point B and n gas stations at distances  $x_1 < x_2 < x_3 < \cdots < x_n$  in kilometers from A along the path from A to B

Output: The minimum number of refills to get from A to B, besides refill at A.

# Greedy Strategy

- Make some greedy choice
- Reduce to a smaller problem
- Iterate

# Greedy Choice

- Refill at the the closest gas station
- Refill at the farthest reachable gas station
- Go until there is no fuel

# Greedy Choice

- Refill at the the closest gas station
- Refill at the farthest reachable gas station
- Go until there is no fuel

Start at A

- Start at A
- Refill at the farthest reachable gas station G

- Start at A
- Refill at the farthest reachable gas station G
- Make *G* the new *A*

- Start at A
- Refill at the farthest reachable gas station G
- Make G the new A
- Get from new A to B with minimum number of refills

### Definition

Subproblem is a similar problem of smaller size.

### Examples

• LargestNumber(3, 9, 5, 9, 7, 1) =

### Examples

■ LargestNumber(3, 9, 5, 9, 7, 1) = "9" +

### Examples

■ LargestNumber(3, 9, 5, 9, 7, 1) = "'9" + LargestNumber(3, 5, 9, 7, 1)

### Examples

- LargestNumber(3, 9, 5, 9, 7, 1) =''9'' + LargestNumber(3, 5, 9, 7, 1)
- Min number of refills from A to B =

### Examples

- LargestNumber(3, 9, 5, 9, 7, 1) = "9" + LargestNumber(3, 5, 9, 7, 1)
- Min number of refills from A to B = first refill at G +

### Examples

- LargestNumber(3, 9, 5, 9, 7, 1) ="'9" + LargestNumber(3, 5, 9, 7, 1)
- Min number of refills from A to B = first refill at G + min number of refills from G to B


### Safe Move


### Definition

A greedy choice is called <u>safe move</u> if there is an optimal solution consistent with this first move.

### Lemma


To refill at the farthest reachable gas station is a safe move.









First case: G is closer than  $G_2$ 



First case: G is closer than  $G_2$ 




First case: G is closer than  $G_2$ Refill at G instead of  $G_1$ 




First case: G is closer than  $G_2$ Refill at G instead of  $G_1$ 




Second case:  $G_2$  is closer than G




Second case:  $G_2$  is closer than G



Second case:  $G_2$  is closer than GAvoid refill at  $G_1$ 



Second case:  $G_2$  is closer than GAvoid refill at  $G_1$ 



Second case:  $G_2$  is closer than GAvoid refill at  $G_1$ 

Route R with the minimum number of refills

- Route R with the minimum number of refills
- $lue{G}_1$  position of first refill in R

- Route R with the minimum number of refills
- $G_1$  position of first refill in R
- $G_2$  next stop in R (refill or B)

- Route R with the minimum number of refills
- $G_1$  position of first refill in R
- $G_2$  next stop in R (refill or B)
- G farthest refill reachable from A

- Route R with the minimum number of refills
- $lue{G}_1$  position of first refill in R
- $G_2$  next stop in R (refill or B)
- G farthest refill reachable from A
- If G is closer than  $G_2$ , refill at G instead of  $G_1$

- Route R with the minimum number of refills
- $G_1$  position of first refill in R
- $G_2$  next stop in R (refill or B)
- lacksquare G farthest refill reachable from A
- If G is closer than  $G_2$ , refill at G instead of  $G_1$
- Otherwise, avoid refill at  $G_1$

## Outline

- 1 Largest Number
- 2 Car Fueling
- 3 Implementation and Analysis
- 4 Main Ingredients

$$A = x_0 \le x_1 \le x_2 \le \cdots \le x_n \le x_{n+1} = B$$

# MinRefills(x, n, L)

 $numRefills \leftarrow 0$ ,  $currentRefill \leftarrow 0$ while *currentRefill* < n:  $lastRefill \leftarrow currentRefill$ 

while (currentRefill  $\leq n$  and

if currentRefill == lastRefill: return IMPOSSIBLE

 $numRefills \leftarrow numRefills + 1$ 

if *currentRefill* < *n*:

return numRefills

 $x[currentRefill + 1] - x[lastRefill] \leq L$ :  $currentRefill \leftarrow currentRefill + 1$ 

The running time of MinRefills(x, n, L) is O(n).

The running time of MinRefills(x, n, L) is O(n).

### Proof

• currentRefill changes from 0 to n + 1, one-by-one

The running time of MinRefills(x, n, L) is O(n).

- currentRefill changes from 0 to n + 1, one-by-one
- **numRefills** changes from 0 to at most n, one-by-one

The running time of MinRefills(x, n, L) is O(n).

- currentRefill changes from 0 to n + 1, one-by-one
- numRefills changes from 0 to at most n, one-by-one
- Thus, O(n) iterations

## Outline

- 1 Largest Number
- 2 Car Fueling
- 3 Implementation and Analysis
- 4 Main Ingredients

## Reduction to Subproblem

- Make a first move
- Then solve a problem of the same kind
- Smaller: fewer digits, fewer fuel stations
- This is called a "subproblem"

## Safe move

A move is called safe if there is an optimal solution consistent with this first move

### Safe move

- A move is called safe if there is an optimal solution consistent with this first move
- Not all first moves are safe

### Safe move

- A move is called safe if there is an optimal solution consistent with this first move
- Not all first moves are safe
- Often greedy moves are not safe

Problem


Problem greedy choice

Make a greedy choice


Problem greedy choice

Safe move

- Make a greedy choice
- Prove that it is a safe move



- Make a greedy choice
- Prove that it is a safe move
- Reduce to a subproblem



- Make a greedy choice
- Prove that it is a safe move
- Reduce to a subproblem
- Solve the subproblem