MedAssist AI

Recommended Blood Tests

For: Hepatocellular Carcinoma (C22.0)

⚠️ Educational use only. AI-generated recommendations — always consult a qualified doctor before ordering any tests.

1 Liver Function Tests

Essential

LFTs

To assess liver damage and detect abnormalities in liver enzymes, bilirubin, and protein levels common in liver cancer.

Normal range: ALT: 7-56 U/L; AST: 8-48 U/L; ALP: 44-147 U/L; Total Bilirubin: 0.1-1.2 mg/dL; Albumin: 3.5-5.0 g/dL

Fasting for 8–12 hours may be required. Blood is drawn from your arm.

2 Alpha-Fetoprotein

Essential

AFP

AFP is a tumor marker often elevated in hepatocellular carcinoma; used to help diagnose and monitor treatment response.

Normal range: Normal: <10 ng/mL; Elevated levels suggest liver cancer or other conditions

No special preparation needed, but your doctor may ask for fasting.

MedAssist AI

To check for anemia, infection, or low platelet count, which can occur due to liver disease or cancer spread.

Normal range: Hemoglobin: 12-16 g/dL; WBC: 4,500-11,000 cells/mcL; Platelets: 150,000-450,000/mcL

Simple blood draw; no fasting usually required.

4

Prothrombin Time

PT/INR

Recommended

Measures how long it takes blood to clot; liver dysfunction can impair clotting factor production.

Normal range: PT: 11-13 seconds; INR: 0.8-1.1

Blood sample taken from arm; inform doctor of any blood thinners you're taking.

5

Serum Creatinine and Blood Urea Nitrogen

Cr, BUN

Recommended

To evaluate kidney function, especially important if imaging with contrast is planned or in advanced liver disease.

Normal range: Creatinine: 0.6-1.1 mg/dL; BUN: 7-20 mg/dL

Fasting may be required; blood drawn from arm.

6

Hepatitis B and C Serology

HBsAg, anti-HCV

Recommended

Chronic hepatitis B or C infection is a major risk factor for hepatocellular carcinoma; testing helps identify underlying cause.

Normal range: Negative for HBsAg and anti-HCV is normal

Blood test to check for past or current viral hepatitis infection.

MedAssist AI

Gives an overview of liver and kidney function, electrolyte balance, and blood sugar, useful for overall assessment.

Normal range: Glucose: 70–99 mg/dL; Electrolytes and organ function values vary; see lab reference

Fasting for 8–12 hours is often required for accurate results.

8

Ferritin Test

Ferritin

Optional

To evaluate iron stores, which can be depleted in patients with liver disease and liver cancer.

Normal range: 15-200 ng/mL

No special preparation is required.