MedAssist AI

Recommended Blood Tests

For: Tuberculosis (A15.0)

⚠️ Educational use only. AI-generated recommendations — always consult a qualified doctor before ordering any tests.

1 Complete Blood Count

CBC

Essential

Checks for signs of infection like anemia or high white blood cell count common in TB

Normal range: Hb: 12-16 g/dL, WBC: 4,500-11,000 cells/μL

Simple blood draw; no special preparation

2 Erythrocyte Sedimentation Rate

ESR

Essential

Measures inflammation in the body, often elevated in active TB

Normal range: 0-20 mm/hr (may be higher in older adults)

No fasting required; part of routine blood work

3 C-Reactive Protein

CRP

Recommended

Another marker of inflammation that can be elevated in active TB infection

Normal range: Normal: <10 mg/L

Fasting not required; measures body inflammation

MedAssist AI

LFTs

Monitors liver health since TB medications can cause liver damage

Normal range: `ALT: 7-56 U/L, AST: 10-40 U/L`

Part of CMP; monitored regularly during treatment

5

Tuberculosis Blood Test (Interferon-Gamma Release Assay)

IGRA

Essential

Helps detect latent or active TB by measuring immune response to TB bacteria

Normal range: `Negative result (varies by lab)`

No fasting needed; blood drawn in one visit

6

Renal Function Tests

RFTs

Recommended

To assess kidney function, as TB treatment can affect kidney health

Normal range: `Varies by component (e.g., creatinine: 0.6-1.2 mg/dL)`

No fasting required, standard blood draw

7

Kidney Function Tests

KFTs

Recommended

To assess kidney function and detect any potential damage from TB treatment

Normal range: `Creatinine: 0.6 - 1.2 mg/dL, Urea: 2.5 - 7.5 mmol/L`

A blood test that measures kidney function. You may need to fast for 8-12 hours beforehand.

MedAssist AI

Evaluates organ function and electrolyte balance, important before starting TB drugs

Normal range: `Varies by component (e.g., creatinine: 0.6-1.1 mg/dL)`

Fasting for 8–12 hours may be needed