MedAssist AI

Recommended Blood Tests

For: Pneumonia (J18.9)

⚠️ Educational use only. AI-generated recommendations — always consult a qualified doctor before ordering any tests.

1 Complete Blood Count

CBC

Essential

To check for signs of infection, inflammation, and anemia; assess overall blood health

Normal range: WBC: 4,500-11,000 cells/μL; Hemoglobin: 13.8-17.2 g/dL; Platelets: 150,000-450,000/μL

A small blood sample is taken from your arm. No fasting needed.

2 C-Reactive Protein

CRP

Essential

To measure inflammation in the body, which increases during infections like pneumonia; assess infection severity

Normal range: Normal: <10 mg/L; Elevated in infection or inflammation

Blood is drawn from your arm. No special preparation needed.

3 Blood Culture

BC

Essential

To detect bacteria or fungi in the blood, especially if the infection is severe; identify the cause of pneumonia

Normal range: No growth of pathogens after 5 days

Two blood samples taken from different sites; no fasting needed.

MedAssist AI

Helps distinguish bacterial pneumonia from viral infections and guides antibiotic use

Normal range: `Normal: <0.05 ng/mL; Levels rise significantly in bacterial infections`

Simple blood draw; no fasting required.

5

Blood Glucose

BG

Recommended

To monitor diabetes control, as infection can cause blood sugar levels to rise; important in patients with diabetes

Normal range: `Fasting: 70-99 mg/dL; Random: <140 mg/dL`

Blood sample taken at any time; important to track if diabetic.

6

Arterial Blood Gas

ABG

Recommended

To check oxygen and carbon dioxide levels in blood, especially if breathing is difficult; assess respiratory function

Normal range: `pH: 7.35-7.45; PaO2: 75-100 mmHg; PaCO2: 35-45 mmHg`

Blood taken from an artery, usually in the wrist; may cause brief discomfort.

7

Electrolyte Panel

EP

Recommended

To monitor electrolyte balance, especially in patients with asthma and diabetes; part of metabolic evaluation

Normal range: `Varies by component (e.g., sodium: 135-145 mmol/L)`

No fasting required, standard blood draw

MedAssist AI

To evaluate kidney and liver function, electrolyte balance, and blood glucose levels

Normal range:	Includes 14 tests; e.g., Sodium: 135–145 mEq/L; Creatinine: 0.7–1.3 mg/dL

Fasting for 8–12 hours may be required for accurate results.