

MedAssist AI

Recommended Blood Tests

For: [Urinary Tract Infection](#) (N39.0)

⚠ Educational use only. AI-generated recommendations — always consult a qualified doctor before ordering any tests.

1

Complete Blood Count

Essential

CBC

To check for signs of infection, inflammation, or anemia; elevated white blood cell count indicates immune response.

Normal range: WBC: 4,500–11,000 cells/ μ L; Hemoglobin: 12–16 g/dL (female)

No fasting needed. A small blood sample is taken from your arm.

2

Blood Urea Nitrogen

Essential

BUN

Assesses kidney function; important in patients with existing kidney disease and suspected UTI complications.

Normal range: 6–24 mg/dL

Fasting may be preferred; blood drawn from arm.

3

Creatinine

Essential

Cr

Evaluates kidney function and detects potential kidney damage from infection or disease progression.

Normal range: 0.5–1.2 mg/dL (varies by age, gender, and kidney health)

No fasting required; part of routine blood work.

MedAssist AI

CRP

Measures systemic inflammation; helps confirm infection and assess severity, particularly in elderly or complex cases.

Normal range: <10 mg/L

Simple blood draw; no special preparation needed.

5

Electrolyte Panel

Recommended

Electrolytes

Checks sodium, potassium, chloride, and other electrolytes; imbalances can occur with kidney disease, infection, or dehydration.

Normal range: Sodium: 135–145 mEq/L; Potassium: 3.5–5.0 mEq/L

Simple blood test; often done with other labs.

6

Blood Culture

Recommended

BCx

Detects bacteremia; critical if sepsis is suspected from a spreading urinary tract infection.

Normal range: No growth after 5 days

Two samples taken from different sites to reduce contamination risk; no fasting needed.

7

Urinalysis

Recommended

UA

Detects bacteria, white blood cells, red blood cells, nitrites, and other markers of urinary tract infection.

Normal range: Varies by parameter

A clean catch urine sample is usually required.

MedAssist AI

~~

Identifies specific pathogen causing UTI and guides targeted antibiotic therapy.

Normal range: No growth after 24-48 hours

A clean catch urine sample is needed; may take a few days for results.